

Kumares Ghosh

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7773003/kumares-ghosh-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

162
papers

2,875
citations

28
h-index

40
g-index

172
ext. papers

3,117
ext. citations

2.7
avg, IF

5.94
L-index

#	Paper	IF	Citations
162	Progress of 3-aminopyridine-based amide, urea, imine and azo derivatives in supramolecular gelation. <i>Journal of the Indian Chemical Society</i> , 2022 , 100462		0
161	Silver-Ion-Selective Gelation of Simple Pyridine-Naphthalimide Conjugates with Multiple Applications: Sensing, Drug Delivery, Dye Adsorption and Ion Conductivity. <i>ChemistrySelect</i> , 2021 , 6, 11696-11705	1.8	1
160	Naphthalene-Coupled Pyridinium Urea Salt in Fluorometric Sensing of Iodide. <i>ChemistrySelect</i> , 2021 , 6, 6353-6359	1.8	1
159	Supramolecular gels in cyanide sensing: a review. <i>Materials Chemistry Frontiers</i> , 2021 , 5, 584-602	7.8	18
158	Naphthalimide-decorated imino-phenol: supramolecular gelation and selective sensing of Fe ³⁺ and Cu ²⁺ ions under different experimental conditions. <i>New Journal of Chemistry</i> , 2021 , 45, 5213-5220	3.6	2
157	A naphthalimide-linked new pyridylazo phenol derivative for selective sensing of cyanide ions (CN ⁻) in sol-gel medium. <i>Analytical Methods</i> , 2021 , 13, 695-702	3.2	1
156	A new 1,2,3-triazole-decorated imino-phenol: selective sensing of Zn ²⁺ , Cu ²⁺ and picric acid under different experimental conditions. <i>New Journal of Chemistry</i> , 2021 , 45, 10923-10929	3.6	4
155	Dipyrromethane-Based Receptor for Fluorometric Sensing of Hydrogenpyrophosphate. <i>ChemistrySelect</i> , 2021 , 6, 8932-8937	1.8	0
154	Anthracene labeled poly(pyridine methacrylamide) as a polymer-based chemosensor for detection of pyrophosphate (PO) in semi-aqueous media. <i>Analytical Methods</i> , 2020 , 12, 5699-5708	3.2	1
153	Selective Dosimetric Sensing of Hg ²⁺ Ions by Design-Based Small Molecular Gelator. <i>ChemistrySelect</i> , 2020 , 5, 5099-5108	1.8	3
152	Diamino malenitrile-linked naphthalimide in selective sensing of F ⁻ , CN ⁻ , Hg ²⁺ and Cu ²⁺ under different experimental conditions. <i>Supramolecular Chemistry</i> , 2020 , 32, 403-413	1.8	4
151	Anthraimidazoledione Derivatives in Fluoride Sensing Ensuing Si-O Bond Cleavage in Organic and Aqueous Medium. <i>ChemistrySelect</i> , 2020 , 5, 5595-5603	1.8	2
150	Effect of Substitution at Amine Functionality of 2,6-Diaminopyridine-Coupled Rhodamine on Metal-Ion Interaction and Self-Assembly. <i>ACS Omega</i> , 2020 , 5, 13984-13993	3.9	10
149	Copillar[5]arene-rhodamine conjugate as a selective sensor for Hg ²⁺ ions. <i>New Journal of Chemistry</i> , 2020 , 44, 5921-5928	3.6	8
148	Design and synthesis of azaindole heterocycle decorated new scaffold in fluorometric sensing of F ⁻ and H ₂ PO ₄ ⁻ . <i>Journal of Heterocyclic Chemistry</i> , 2020 , 57, 3558-3565	1.9	2
147	Progress in Benzimidazole/Benzimidazolium-Derived Supramolecular Gelators in Ion Recognition. <i>Mini-Reviews in Organic Chemistry</i> , 2020 , 17, 1042-1055	1.7	4
146	Diaminomaleonitrile-functionalized gelators in F ⁻ /CN ⁻ sensing, phase-selective gelation, oil spill recovery and dye removal from water. <i>New Journal of Chemistry</i> , 2020 , 44, 10275-10285	3.6	9

145	Adenine-linked naphthalimide: A case of selective colorimetric as well as fluorometric sensing of F ⁻ and anion-activated moisture detection in organic solvents and CO-sensing. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2020 , 229, 117910	4.4	17
144	Naphthalimide-Benzothiazole Conjugate: A Dosimetric Probe for Colorimetric and Fluorometric Detection of Palladium. <i>ChemistrySelect</i> , 2020 , 5, 8363-8369	1.8	2
143	Cholesterol-Coupled Diazine-Phenol Gelator: Cyanide Sensing, Phase-Selective Gelation in Oil Spill Recovery and Dye Adsorption. <i>ChemistrySelect</i> , 2020 , 5, 11874-11881	1.8	3
142	Dipyrromethane Decorated Gelator in Anion Recognition and Solvent-Dependent Aggregation-Induced Emission. <i>ChemistrySelect</i> , 2020 , 5, 9635-9640	1.8	2
141	Triazole-amide isosteric pyridine-based supramolecular gelators in metal ion and biothiol sensing with excellent performance in adsorption of heavy metal ions and picric acid from water. <i>New Journal of Chemistry</i> , 2019 , 43, 934-945	3.6	17
140	A 1,8-naphthalimide-pyridoxal conjugate as a supramolecular gelator for colorimetric read out of F ⁻ ions in solution, gel and solid states. <i>New Journal of Chemistry</i> , 2019 , 43, 2718-2725	3.6	19
139	Cholesterol linked benzothiazole: a versatile gelator for detection of picric acid and metal ions such as Ag ⁺ , Hg ²⁺ , Fe ³⁺ and Al ³⁺ under different conditions. <i>New Journal of Chemistry</i> , 2019 , 43, 10509-10516	3.6	9
138	A sulfonyl hydrazone cholesterol conjugate: gelation, anion interaction and its application in dye adsorption. <i>New Journal of Chemistry</i> , 2019 , 43, 10270-10277	3.6	9
137	Perimedine-linked rhodamine dye in visual sensing of Al ³⁺ , Fe ³⁺ and Fe ²⁺ ions in aqueous organic medium under different experimental conditions. <i>Supramolecular Chemistry</i> , 2019 , 31, 645-652	1.8	1
136	Pyridyl Azo-Based Progelator in Selective Sensing of Hg ²⁺ and Ag ⁺ Ions via Sol to Gel Conversion. <i>ChemistrySelect</i> , 2019 , 4, 11564-11571	1.8	6
135	4-Hydroxybenzaldehyde derived Schiff base gelators: case of the sustainability or rupturing of imine bonds towards the selective sensing of Ag ⁺ and Hg ²⁺ ions via sol-gel methodology. <i>New Journal of Chemistry</i> , 2019 , 43, 5139-5149	3.6	18
134	Aryl ethers decorated gallic acid-naphthalimide conjugate: aggregation and sensing towards amines and F ⁻ . <i>Supramolecular Chemistry</i> , 2019 , 31, 732-744	1.8	3
133	Pyridine-Based Macrocyclic and Open Receptors for Urea. <i>ChemistrySelect</i> , 2019 , 4, 12825-12831	1.8	3
132	Naphthalimide-linked bispyridinium clefts in selective aqueous sensing of triphosphate and triphosphate-based biomolecules. <i>Analytical Methods</i> , 2019 , 11, 5864-5871	3.2	4
131	Cholesterol-based simple supramolecular gelators: an approach to selective sensing of CN ⁻ ion with application in dye adsorption. <i>Supramolecular Chemistry</i> , 2019 , 31, 239-250	1.8	12
130	Chromenone-rhodamine conjugate for naked eye detection of Al ³⁺ and Hg ²⁺ ions in semi aqueous medium. <i>Supramolecular Chemistry</i> , 2019 , 31, 1-8	1.8	9
129	Pyridine/pyridinium symmetrical bisamides as functional materials: aggregation, selective sensing and drug release. <i>New Journal of Chemistry</i> , 2018 , 42, 6488-6497	3.6	36
128	Pyridine coupled mono and bisbenzimidazoles as supramolecular gelators: selective metal ion sensing and ionic conductivity. <i>Materials Chemistry Frontiers</i> , 2018 , 2, 385-395	7.8	28

127	Pyridylazo Derivatives with Dicyanovinyl Appendage in Selective Sensing of CN^- in Sol-Gel Medium. <i>ChemistrySelect</i> , 2018 , 3, 1809-1814	1.8	25
126	Azo and imine functionalized 2-naphthols: promising supramolecular gelators for selective detection of Fe^{3+} and Cu^{2+} , reactive oxygen species and halides. <i>Materials Chemistry Frontiers</i> , 2018 , 2, 1866-1875	7.8	29
125	Selective sensing of Hg^{2+} via sol-gel transformation of a cholesterol-based compound. <i>Supramolecular Chemistry</i> , 2018 , 30, 722-729	1.8	21
124	Cholesterol-based diazine derivative: selective sensing of Ag^+ and Fe^{3+} ions through gelation and the performance of metallogels in dye and picric acid adsorption from water. <i>Materials Chemistry Frontiers</i> , 2018 , 2, 2286-2296	7.8	24
123	Dosimetric Chromogenic Probe for Selective Detection of Sulfide via Sol-Gel Methodology. <i>ACS Omega</i> , 2018 , 3, 17319-17325	3.9	7
122	Pyridyl Azo-Based Naphthyl Acetate for Sensing of Hydrazine and Perborate in Sol-Gel Medium. <i>ChemistrySelect</i> , 2018 , 3, 9448-9453	1.8	14
121	Diaminomaleonitrile-decorated cholesterol-based supramolecular gelator: aggregation, multiple analyte (hydrazine, Hg^{2+} and Cu^{2+}) detection and dye adsorption. <i>New Journal of Chemistry</i> , 2018 , 42, 13718-13725	3.6	22
120	Nano-Pelargonidin Protects Hyperglycemic-Induced L6 Cells against Mitochondrial Dysfunction. <i>Planta Medica</i> , 2017 , 83, 468-475	3.1	19
119	Visual Sensing of Ag^+ Ions through Gelation of Cholesterol- Appended Benzimidazole and Associated Ion Conducting Behaviour. <i>ChemistrySelect</i> , 2017 , 2, 959-966	1.8	15
118	1,4-Disubstituted 1,2,3-Triazole- and 1,5-Disubstituted 1,2,3-Triazole-Based Bis-Sulfonamides in Selective Fluorescence Sensing of ATP. <i>ChemistrySelect</i> , 2017 , 2, 2034-2038	1.8	8
117	Naphthalene and pyrrole substituted guanidine in selective sensing of Cu^{2+} , Hg^{2+} , Pb^{2+} and CN^- ions under different conditions. <i>Supramolecular Chemistry</i> , 2017 , 29, 528-535	1.8	7
116	Triphenylamine-Based Open and Macrocyclic Receptors: A Study Towards Selective Recognition of Aliphatic Dicarboxylates. <i>ChemistrySelect</i> , 2017 , 2, 4794-4799	1.8	4
115	Anthraquinone Derived Cholesterol Linked Imidazole Gelator in Visual Sensing of Picric Acid. <i>ChemistrySelect</i> , 2017 , 2, 4800-4806	1.8	15
114	Coumarin-based urea-amide scaffold in ratiometric fluorescence sensing of CN^- . <i>Tetrahedron Letters</i> , 2017 , 58, 2038-2043	2	13
113	Fluorophore inserted bisbenzimidazole clefts in selective sensing of Ag^+ and Cu^{2+} ions. <i>Journal of Photochemistry and Photobiology A: Chemistry</i> , 2017 , 348, 110-117	4.7	8
112	α -Amino Acid Derived Benzimidazole-Linked Rhodamines: A Case of Substitution Effect at the Amino Acid Site toward Spiro Ring Opening for Selective Sensing of Al Ions. <i>Inorganic Chemistry</i> , 2017 , 56, 8889-8899	5.1	14
111	Cholesterol-Appended Benzimidazolium Salts: Synthesis, Aggregation, Sensing, Dye Adsorption, and Semiconducting Properties. <i>Langmuir</i> , 2017 , 33, 8277-8288	4	27
110	Pyrrole-based tetra-amide for hydrogen pyrophosphate ($\text{HP}2\text{O}_7^{3-}$) and F^- ions in sol-gel medium. <i>Supramolecular Chemistry</i> , 2017 , 29, 946-952	1.8	6

109	A Pyridinium Urea-Coupled Polyether Receptor for the Selective Sensing of Lysine and Cell Imaging. <i>European Journal of Organic Chemistry</i> , 2017 , 2017, 355-362	3.2	2
108	Naphthalene-cholesterol conjugate as simple gelator for selective sensing of CN^- ion. <i>Supramolecular Chemistry</i> , 2017 , 29, 350-359	1.8	28
107	New Six-Membered pH-Insensitive Rhodamine Spirocyclic in Selective Sensing of Cu through C-C Bond Cleavage and Its Application in Cell Imaging. <i>ACS Omega</i> , 2017 , 2, 8167-8176	3.9	19
106	Isomeric chiral pyrrole diamides and their efficacy in enantioselective sensing of tartrate in sol-gel medium. <i>Tetrahedron Letters</i> , 2016 , 57, 3629-3634	2	11
105	Aryl ethers coupled pyridoxal as supramolecular gelator for selective sensing of F^- <i>Tetrahedron Letters</i> , 2016 , 57, 5469-5474	2	23
104	Cholesterol appended bis-1,2,3-triazoles as simple supramolecular gelators for the naked eye detection of Ag^+ , Cu^{2+} and Hg^{2+} ions. <i>New Journal of Chemistry</i> , 2016 , 40, 3476-3483	3.6	36
103	Dipicolylamine coupled rhodamine dyes: new clefts for highly selective naked eye sensing of Cu^{2+} and CN^- ions. <i>RSC Advances</i> , 2016 , 6, 47802-47812	3.7	15
102	Cholesterol Based Bisamides on Biphenyl Backbone: A Case of Selective Visual Sensing of F^- and H_2PO_4^- through Breaking and Making of Gels. <i>ChemistrySelect</i> , 2016 , 1, 3667-3674	1.8	13
101	Benzimidazolium-based chemosensors: selective recognition of H_2PO_4^- , $\text{HP}_2\text{O}_7^{3-}$, F^- and ATP through fluorescence and gelation studies. <i>RSC Advances</i> , 2015 , 5, 46608-46616	3.7	16
100	Pyridinium-based flexible tripodal cleft: a case of fluorescence sensing of ATP and dihydrogenphosphate under different conditions and cell imaging. <i>RSC Advances</i> , 2015 , 5, 35175-35180	3.7	12
99	Naphthalene linked pyridyl urea as a supramolecular gelator: a new insight into naked eye detection of I^- in the gel state with semiconducting behaviour. <i>RSC Advances</i> , 2015 , 5, 72772-72779	3.7	21
98	L-Amino acid derived pyridinium-based chiral compounds and their efficacy in chiral recognition of lactate. <i>RSC Advances</i> , 2015 , 5, 24499-24506	3.7	7
97	Piperazine-based new sensor: selective ratiometric sensing of Fe^{3+} , logic gate construction and cell imaging. <i>Supramolecular Chemistry</i> , 2015 , 27, 224-232	1.8	10
96	Rhodamine-linked pyridyl thiourea as a receptor for selective recognition of F^- , Al^{3+} and Ag^+ under different conditions. <i>Supramolecular Chemistry</i> , 2015 , 27, 490-500	1.8	14
95	Coumarin-based supramolecular gelator: a case of selective detection of F^- and $\text{HP}_2\text{O}_7^{3-}$ <i>RSC Advances</i> , 2015 , 5, 12094-12099	3.7	32
94	Rhodamine-labelled simple architectures for fluorometric and colorimetric sensing of Hg^{2+} and Pb^{2+} ions in semi-aqueous and aqueous environments. <i>Analytical Methods</i> , 2014 , 6, 2648-2654	3.2	12
93	Progress of 3-aminopyridinium-based synthetic receptors in anion recognition. <i>RSC Advances</i> , 2014 , 4, 20114-20130	3.7	20
92	Azaindole-1,2,3-triazole conjugate in a tripod for selective sensing of Cl^- , H_2PO_4^- and ATP under different conditions. <i>RSC Advances</i> , 2014 , 4, 11590	3.7	12

91	Fluoride-responsive hydrogel of cholesterol appended pyridinium urea and its metal detecting ability and semi-conducting behaviour. <i>Supramolecular Chemistry</i> , 2014 , 26, 313-320	1.8	12
90	Ion conducting cholesterol appended pyridinium bisamide-based gel for the selective detection of Ag ⁺ and Cl ⁻ ions. <i>RSC Advances</i> , 2014 , 4, 3732-3737	3.7	56
89	Selective sensing of Al ³⁺ by naphthyridine coupled rhodamine chemosensors. <i>RSC Advances</i> , 2014 , 4, 23428-23432	3.7	18
88	Benzimidazolium-based new simple ratiometric fluorescent sensor for selective detection of dihydrogenphosphate. <i>Supramolecular Chemistry</i> , 2014 , 26, 856-863	1.8	9
87	A benzimidazolium-based new flexible cleft built on the piperazine unit: a case of selective fluorometric sensing of ATP. <i>RSC Advances</i> , 2014 , 4, 58530-58535	3.7	6
86	Anthracene-labeled pyridinium-based symmetrical chiral chemosensor for enantioselective recognition of l-tartrate. <i>Tetrahedron Letters</i> , 2014 , 55, 1342-1346	2	6
85	Azaindole-1,2,3-triazole conjugate as selective fluorometric sensor for dihydrogenphosphate. <i>RSC Advances</i> , 2013 , 3, 16144	3.7	11
84	l-Valine derived benzimidazole based bis-urea in enantioselective fluorescence sensing of L-tartrate. <i>Tetrahedron Letters</i> , 2013 , 54, 4568-4573	2	10
83	Pyridinium-based tripodal chemosensor in visual sensing of AMP in water by indicator displacement assay (IDA). <i>Organic and Biomolecular Chemistry</i> , 2013 , 11, 5666-72	3.9	19
82	Anthraquinone coupled benzothiazole-based receptor for selective sensing of Cu ²⁺ . <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2013 , 77, 67-74	1.7	19
81	Enantioselective sensing of lactate by pyridinium-based chiral receptor. <i>Tetrahedron Letters</i> , 2013 , 54, 5686-5689	2	8
80	Benzimidazolium-based receptors: Case of iodide/water cluster induced supramolecular chain and improved fluorometric binding of iodide involving alcoholic group. <i>Journal of Molecular Structure</i> , 2013 , 1042, 57-65	3.4	5
79	Rhodamine-labelled new architecture for dual sensing of Co ²⁺ and Hg ²⁺ ions. <i>Tetrahedron Letters</i> , 2013 , 54, 6464-6468	2	17
78	Rhodamine-labeled Sensor Bead as a Colorimetric and Fluorometric Dual Assay for Hg ²⁺ Ions in Water. <i>Asian Journal of Organic Chemistry</i> , 2013 , 2, 157-163	3	9
77	O-tert-butyldiphenylsilyl coumarin and dicoumarol: a case toward selective sensing of F ⁻ ions in organic and aqueous environments. <i>Analyst</i> , 2013 , 138, 3038-45	5	27
76	A new quinoline-based chemosensor in ratiometric sensing of Hg ²⁺ ions. <i>Supramolecular Chemistry</i> , 2013 , 25, 127-132	1.8	13
75	Piperazine-based simple structure for selective sensing of Hg ²⁺ and glutathione and construction of a logic circuit mimicking an INHIBIT gate. <i>New Journal of Chemistry</i> , 2013 , 37, 4206	3.6	20
74	Eneidyne scaffold-based highly selective chemosensor for ratiometric sensing of H ₂ PO ₄ ⁻ ions. <i>Tetrahedron Letters</i> , 2012 , 53, 2054-2058	2	20

73	Anthracene-Labeled 1,2,3-Triazole-Linked Bispyridinium Amide for Selective Sensing of H ₂ PO ₄ ³⁻ by Fluorescence and Gel Formation. <i>European Journal of Organic Chemistry</i> , 2012 , 2012, 1311-1317	3.2	27
72	Coumarin-based symmetrical bisamide as fluorescent and colorimetric probes for copper ions. <i>Supramolecular Chemistry</i> , 2012 , 24, 197-203	1.8	7
71	Pyridinium-based fluororeceptors as practical chemosensors for hydrogen pyrophosphate (HP ₂ O ₇ ³⁻) in semiaqueous and aqueous environments. <i>Organic Letters</i> , 2012 , 14, 4314-7	6.2	58
70	Ortho-phenylenediamine-based open and macrocyclic receptors in selective sensing of H ₂ PO ₄ ⁻ , ATP and ADP under different conditions. <i>Organic and Biomolecular Chemistry</i> , 2012 , 10, 9383-92	3.9	21
69	A rhodamine appended tripodal receptor as a ratiometric probe for Hg ²⁺ ions. <i>Organic and Biomolecular Chemistry</i> , 2012 , 10, 3236-43	3.9	46
68	L-Valine-derived simple benzimidazole-based host in selective sensing of Hg(II) ions. <i>Supramolecular Chemistry</i> , 2012 , 24, 748-754	1.8	6
67	Rhodamine-based bis-sulfonamide as a sensing probe for Cu ²⁺ and Hg ²⁺ ions. <i>New Journal of Chemistry</i> , 2012 , 36, 2121	3.6	49
66	Cholesterol appended pyridinium ureas: a case of gel making and breaking for selective visual readout of F ⁻ . <i>Organic and Biomolecular Chemistry</i> , 2012 , 10, 8800-7	3.9	23
65	Design and synthesis of anthracene-based bispyridinium amides: anion binding, cell staining and DNA interaction studies. <i>New Journal of Chemistry</i> , 2012 , 36, 1231	3.6	23
64	Pyrene-based simple new hetero bis amide pyridinium salt for selective sensing of benzoate and hydrogen sulphate. <i>Supramolecular Chemistry</i> , 2011 , 23, 365-371	1.8	8
63	Selective sensing of Cu (II) by a simple anthracene-based tripodal chemosensor. <i>Supramolecular Chemistry</i> , 2011 , 23, 435-440	1.8	19
62	Pyridinium-based symmetrical diamides as chemosensors in visual sensing of citrate through indicator displacement assay (IDA) and gel formation. <i>Organic and Biomolecular Chemistry</i> , 2011 , 9, 6551-389	3.9	43
61	Fluorometric recognition of both dihydrogen phosphate and iodide by a new flexible anthracene linked benzimidazolium-based receptor. <i>Beilstein Journal of Organic Chemistry</i> , 2011 , 7, 254-64	2.5	24
60	Selective Sensing of Fumarate Over Maleate by Benzimidazolium Based Fluororeceptors. <i>Mini-Reviews in Organic Chemistry</i> , 2011 , 8, 31-37	1.7	8
59	Benzimidazolium-based simple host for fluorometric sensing of . <i>Tetrahedron Letters</i> , 2011 , 52, 5098-5103	3.3	33
58	Experimental and theoretical anion binding studies on coumarin linked thiourea and urea molecules. <i>Journal of Molecular Structure</i> , 2011 , 1004, 193-203	3.4	13
57	Triphenylamine-based simple chemosensor for selective fluorometric detection of fluoride, acetate and dihydrogenphosphate ions in different solvents. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2011 , 70, 97-107		9
56	Naphthyridine amide-urea conjugate: a case toward selective fluorometric sensing of N-acetyl proline carboxylate. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2011 , 71, 243-248		3

55	A new ortho-phenylenediamine-based cleft for selective sensing of H ₂ PO ₄ ²⁻ and ATP. <i>New Journal of Chemistry</i> , 2011 , 35, 1397	3.6	26
54	(rac)-1,1'-binaphthyl-based simple receptors designed for fluorometric discrimination of maleic and fumaric acids. <i>Journal of Physical Chemistry B</i> , 2011 , 115, 8597-608	3.4	14
53	Benzimidazolium-based flexible tripodal fluorescent chemosensor for selective sensing of dihydrogenphosphate and ATP. <i>Supramolecular Chemistry</i> , 2011 , 23, 518-526	1.8	12
52	Anthracene-based hetero bisamide chemosensor in fluorescence sensing of monocarboxylates over monocarboxylic acids. <i>Supramolecular Chemistry</i> , 2011 , 23, 539-549	1.8	3
51	Anthracene coupled adenine for the selective sensing of copper ions. <i>Beilstein Journal of Organic Chemistry</i> , 2010 , 6, 44	2.5	11
50	Anthracene appended pyridinium amide-urea conjugate in selective fluorometric sensing of L-N-acetylvaline salt. <i>Beilstein Journal of Organic Chemistry</i> , 2010 , 6, 1211-8	2.5	4
49	Naphthyridine-based symmetrical and unsymmetrical pyridinium amides in sensing of biotin salt. <i>Supramolecular Chemistry</i> , 2010 , 22, 81-94	1.8	6
48	trans-Pyridyl and naphthyridyl cinnamides as alternatives for urea in complexation of carboxylic acid and formation of water-templated assemblies in the solid state. <i>Journal of Physical Chemistry B</i> , 2010 , 114, 321-9	3.4	5
47	One-pot synthesis of linearly fused N-heterocycles from their angular analogues and studies of their redox and electrochromic properties. <i>Journal of Organic Chemistry</i> , 2010 , 75, 2065-8	4.2	6
46	Binding induced destruction of an excimer in anthracene-linked benzimidazole diamide: a case toward the selective detection of organic sulfonic acids and metal ions. <i>New Journal of Chemistry</i> , 2010 , 34, 1387	3.6	20
45	A new benzimidazolium receptor for fluorescence sensing of iodide. <i>Supramolecular Chemistry</i> , 2010 , 22, 311-317	1.8	15
44	Naphthyridine-based receptors for fluorometric detection of urea and biotin. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2010 , 67, 271-280		7
43	Hydrogen bonded assemblies of 1,8-naphthyridine derivatives: discrete or polymeric structures in the solid state. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2010 , 68, 193-199		3
42	Naphthalene appended 2,5-diketopiperazine towards fluorometric response of dihydrogenphosphate. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2010 , 68, 447-452		3
41	Triphenylamine-based receptor for selective recognition of dicarboxylates. <i>Tetrahedron Letters</i> , 2010 , 51, 343-347	2	42
40	Selective sensing of Zn(II) ion by a simple anthracene-based tripodal chemosensor. <i>Tetrahedron Letters</i> , 2010 , 51, 4995-4999	2	37
39	Triphenylamine-Based Pyridine N-Oxide and Pyridinium Salts for Size-Selective Recognition of Dicarboxylates. <i>European Journal of Organic Chemistry</i> , 2009 , 2009, 4515-4524	3.2	44
38	Anthracene-based macrocyclic fluorescent chemosensor for selective sensing of dicarboxylate. <i>Tetrahedron Letters</i> , 2009 , 50, 85-88	2	54

37	Design and synthesis of an ortho-phenylenediamine-based open cleft: a selective fluorescent chemosensor for dihydrogen phosphate. <i>Tetrahedron Letters</i> , 2009 , 50, 2392-2397	2	36
36	A benzthiazole-based simple receptor in fluorescence sensing of biotin ester and urea. <i>Tetrahedron Letters</i> , 2009 , 50, 4096-4100	2	10
35	Pyridinium amide-based simple synthetic receptor for selective recognition of dihydrogenphosphate. <i>Tetrahedron Letters</i> , 2009 , 50, 6557-6561	2	34
34	Anthracene labeled pyridine amides: A class of prototype PET sensors towards monocarboxylic acid. <i>Journal of Photochemistry and Photobiology A: Chemistry</i> , 2009 , 203, 40-49	4-7	8
33	Triphenylamine-based receptors in selective recognition of dicarboxylic acids. <i>Journal of Physical Chemistry B</i> , 2009 , 113, 7800-9	3-4	24
32	Anthracene-based open and macrocyclic receptors in the flurometric detection of urea. <i>New Journal of Chemistry</i> , 2009 , 33, 1965	3.6	8
31	Quinoline based receptor in fluorometric discrimination of carboxylic acids. <i>Beilstein Journal of Organic Chemistry</i> , 2008 , 4, 52	2.5	11
30	Anthracene coupled trans-pyridylcinnamide: a new fluororeceptor for selective sensing of dicarboxylates. <i>Tetrahedron Letters</i> , 2008 , 49, 2592-2597	2	22
29	A quinoline-based tripodal fluororeceptor for citric acid. <i>Tetrahedron Letters</i> , 2008 , 49, 658-663	2	15
28	Anthracene-based ortho-phenylenediamine clefts for sensing carboxylates. <i>Tetrahedron Letters</i> , 2008 , 49, 4591-4595	2	21
27	A pyridine-based macrocyclic host for urea and acetone. <i>Tetrahedron Letters</i> , 2008 , 49, 5063-5066	2	12
26	Adenine-based urea receptors in fluorescent recognition of iodide. <i>Tetrahedron Letters</i> , 2008 , 49, 7204-7208		29
25	A naphthyridine-based receptor for sensing citric acid. <i>Tetrahedron Letters</i> , 2007 , 48, 2935-2938	2	45
24	Anthracene-based ureidopyridyl fluororeceptor for dicarboxylates. <i>Tetrahedron Letters</i> , 2007 , 48, 6129-6132		28
23	N-(6-Methyl-2-pyridyl)acrylamide: a case of amide hydrolysis without the assistance of acid or base in the synthesis of water-driven H-bonded polymeric chains. <i>Tetrahedron Letters</i> , 2007 , 48, 6308-6311	2	1
22	Adenine-based receptor for dicarboxylic acids. <i>Tetrahedron Letters</i> , 2007 , 48, 7022-7026	2	22
21	An anthracene based bispyridinium amide receptor for selective sensing of anions. <i>Tetrahedron Letters</i> , 2007 , 48, 8725-8729	2	51
20	Anthracene-coupled Pyridine Amines: A New Off-On Switch for Molecular Recognition Studies on Dicarboxylic Acids. <i>Chemistry Letters</i> , 2006 , 35, 414-415	1.7	26

19	Water templated hydrogen-bonded network of pyridine amide appended carbamate in solid state. <i>Journal of Molecular Structure</i> , 2006 , 785, 63-67	3-4	9
18	Triphenylamine-based novel PET sensors in selective recognition of dicarboxylic acids. <i>Tetrahedron Letters</i> , 2006 , 47, 2365-2369	2	42
17	Fluorescence sensing of tartaric acid: a case of excimer emission caused by hydrogen bond-mediated complexation. <i>Tetrahedron Letters</i> , 2006 , 47, 3577-3581	2	45
16	Colorimetric and fluorescence sensing of anions using thiourea based coumarin receptors. <i>Tetrahedron Letters</i> , 2006 , 47, 8165-8169	2	85
15	Effect of a hydroxyl group in an anthracene-labelled pyridine amide receptor in molecular recognition of keto and hydroxy monocarboxylic acids. <i>Tetrahedron Letters</i> , 2006 , 47, 9233-9237	2	17
14	Non-bonded O \cdots S contacts and O \cdots H \cdots S hydrogen bonds in isomeric hydroxyphenyl-1,3-dithianes. <i>CrystEngComm</i> , 2005 , 7, 210-215	3-3	7
13	Anthracene-appended Pyridine Amide: A Simple Sensor for Monocarboxylic Acids. <i>Supramolecular Chemistry</i> , 2005 , 17, 331-334	1.8	21
12	Urotropine: a unique scaffold in molecular recognition for phenolic substrates. <i>Journal of Molecular Structure</i> , 2005 , 737, 201-206	3-4	20
11	2-(2-Hydroxyphenyl)-1,3-dithiane. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2003 , 59, o773-o775		5
10	Recognition of insoluble tartaric acid in chloroform. <i>Tetrahedron</i> , 2001 , 57, 4987-4993	2.4	37
9	N-bromosuccinimide reactions of some heterocycles in the presence or absence of water: An overview of ring versus side chain bromination for the synthesis of important brominated heterocyclic synthons. <i>Journal of Heterocyclic Chemistry</i> , 2001 , 38, 173-178	1.9	18
8	1:1 hetero-assembly of 2-aminopyrimidine and (+)-camphoric acid. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2000 , 56 (Pt 4), 477-8		6
7	Self-assembly of 2-pivaloyl-6-chloropterin. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2000 , 56 (Pt 6), 716-7		2
6	Troger's base molecular scaffolds in dicarboxylic acid recognition. <i>Journal of Organic Chemistry</i> , 2000 , 65, 1907-14	4.2	141
5	Novel Motif of Hydrogen Bonds in the Water-assisted Supramolecular Self-assembly of 2-acetylamino-6-methylpyridine-N-oxide and Hetero-assembly of 1:1 Co-crystal of o-Phenylenediamine with Catechol. <i>Supramolecular Chemistry</i> , 2000 , 11, 191-199	1.8	5
4	2-Aminopyrimidine \cdots terephthalic acid (1:1) complex. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999 , 55, 87-89		11
3	Molecular recognition: Hydrogen bonding induced configurational locking of a new photoresponsive receptor by dicarboxylic acids. <i>Tetrahedron Letters</i> , 1999 , 40, 1735-1738	2	33
2	Molecular recognition: Chain length selectivity studies of dicarboxylic acids by the cavity of a new Troger's base receptor. <i>Tetrahedron Letters</i> , 1997 , 38, 4503-4506	2	76

- 1 Molecular recognition: Connection and disconnection of hydrogen bonds, a case study with dimeric and highly associated monocarboxylic acids with simple receptors. *Tetrahedron*, **1996**, 52, 12223-12232 ^{2.4} 24