

Kristina Schoonjans

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7778028/kristina-schoonjans-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

158
papers

23,800
citations

76
h-index

154
g-index

163
ext. papers

26,893
ext. citations

12.6
avg, IF

6.51
L-index

#	Paper	IF	Citations
158	Identification of a Crosstalk among TGR5, GLIS2, and TP53 Signaling Pathways in the Control of Undifferentiated Germ Cell Homeostasis and Chemoresistance.. <i>Advanced Science</i> , 2022 , e2200626	13.6	0
157	Nuclear receptors FXR and SHP regulate protein N-glycan modifications in the liver. <i>Science Advances</i> , 2021 , 7,	14.3	1
156	Central anorexigenic actions of bile acids are mediated by TGR5. <i>Nature Metabolism</i> , 2021 , 3, 595-603	14.6	17
155	Muricholic Acids Promote Resistance to Hypercholesterolemia in Cholesterol-Fed Mice. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	2
154	Hypothalamic bile acid-TGR5 signaling protects from obesity. <i>Cell Metabolism</i> , 2021 , 33, 1483-1492.e10	24.6	22
153	Molecular Physiology of Bile Acid Signaling in Health, Disease, and Aging. <i>Physiological Reviews</i> , 2021 , 101, 683-731	47.9	31
152	TGR5/Cathepsin E signaling regulates macrophage innate immune activation in liver ischemia and reperfusion injury. <i>American Journal of Transplantation</i> , 2021 , 21, 1453-1464	8.7	6
151	Pancreatic Sirtuin 3 Deficiency Promotes Hepatic Steatosis by Enhancing 5-Hydroxytryptamine Synthesis in Mice With Diet-Induced Obesity. <i>Diabetes</i> , 2021 , 70, 119-131	0.9	4
150	The transcriptional coactivator CBP/p300 is an evolutionarily conserved node that promotes longevity in response to mitochondrial stress. <i>Nature Aging</i> , 2021 , 1, 165-178		12
149	Dietary Fiber Is Essential to Maintain Intestinal Size, L-Cell Secretion, and Intestinal Integrity in Mice. <i>Frontiers in Endocrinology</i> , 2021 , 12, 640602	5.7	3
148	Emerging functions of the nuclear receptor LRH-1 in liver physiology and pathology. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2021 , 1867, 166145	6.9	6
147	Downregulation of TGR5 (GPBAR1) in biliary epithelial cells contributes to the pathogenesis of sclerosing cholangitis. <i>Journal of Hepatology</i> , 2021 , 75, 634-646	13.4	11
146	Bile Acids Signal via TGR5 to Activate Intestinal Stem Cells and Epithelial Regeneration. <i>Gastroenterology</i> , 2020 , 159, 956-968.e8	13.3	38
145	L-Cell Differentiation Is Induced by Bile Acids Through GPBAR1 and Paracrine GLP-1 and Serotonin Signaling. <i>Diabetes</i> , 2020 , 69, 614-623	0.9	24
144	Compound 18 Improves Glucose Tolerance in a Hepatocyte TGR5-dependent Manner in Mice. <i>Nutrients</i> , 2020 , 12,	6.7	8
143	Mechano-modulatory synthetic niches for liver organoid derivation. <i>Nature Communications</i> , 2020 , 11, 3416	17.4	49
142	Maternal glucose homeostasis is impaired in mouse models of gestational cholestasis. <i>Scientific Reports</i> , 2020 , 10, 11523	4.9	5

141	Transcriptomic analysis across liver diseases reveals disease-modulating activation of constitutive androstane receptor in cholestasis. <i>JHEP Reports</i> , 2020 , 2, 100140	10.3	2
140	TGR5 Regulates Macrophage Inflammation in Nonalcoholic Steatohepatitis by Modulating NLRP3 Inflammasome Activation. <i>Frontiers in Immunology</i> , 2020 , 11, 609060	8.4	15
139	The RNA-Binding Protein PUM2 Impairs Mitochondrial Dynamics and Mitophagy During Aging. <i>Molecular Cell</i> , 2019 , 73, 775-787.e10	17.6	60
138	A new class of protein biomarkers based on subcellular distribution: application to a mouse liver cancer model. <i>Scientific Reports</i> , 2019 , 9, 6913	4.9	9
137	Bile acids drive colonic secretion of glucagon-like-peptide 1 and peptide-YY in rodents. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 316, G574-G584	5.1	25
136	The orphan nuclear receptor LRH-1/NR5a2 critically regulates T cell functions. <i>Science Advances</i> , 2019 , 5, eaav9732	14.3	9
135	NTCP deficiency in mice protects against obesity and hepatosteatosis. <i>JCI Insight</i> , 2019 , 5,	9.9	11
134	Identifying gene function and module connections by the integration of multispecies expression compendia. <i>Genome Research</i> , 2019 , 29, 2034-2045	9.7	10
133	The G Protein-Coupled Bile Acid Receptor TGR5 (Gpbar1) Modulates Endothelin-1 Signaling in Liver. <i>Cells</i> , 2019 , 8,	7.9	23
132	LRH-1 agonism favours an immune-islet dialogue which protects against diabetes mellitus. <i>Nature Communications</i> , 2018 , 9, 1488	17.4	31
131	Transcriptional regulation by NR5A2 links differentiation and inflammation in the pancreas. <i>Nature</i> , 2018 , 554, 533-537	50.4	57
130	The Orphan Nuclear Receptor Liver Homolog Receptor-1 (Nr5a2) Regulates Ovarian Granulosa Cell Proliferation. <i>Journal of the Endocrine Society</i> , 2018 , 2, 24-41	0.4	22
129	TGR5 signalling promotes mitochondrial fission and beige remodelling of white adipose tissue. <i>Nature Communications</i> , 2018 , 9, 245	17.4	90
128	Bile acids are important direct and indirect regulators of the secretion of appetite- and metabolism-regulating hormones from the gut and pancreas. <i>Molecular Metabolism</i> , 2018 , 11, 84-95	8.8	86
127	An Integrated Systems Genetics and Omics Toolkit to Probe Gene Function. <i>Cell Systems</i> , 2018 , 6, 90-102.e4	11.4	23
126	De novo NAD synthesis enhances mitochondrial function and improves health. <i>Nature</i> , 2018 , 563, 354-359	50.4	163
125	Bile acids deoxycholic acid and ursodeoxycholic acid differentially regulate human Defensin-1 and -2 secretion by colonic epithelial cells. <i>FASEB Journal</i> , 2017 , 31, 3848-3857	0.9	15
124	Ovary-specific depletion of the nuclear receptor Nr5a2 compromises expansion of the cumulus oophorus but not fertilization by intracytoplasmic sperm injection. <i>Biology of Reproduction</i> , 2017 , 96, 1231-1243	3.9	12

123	Small heterodimer partner deletion prevents hepatic steatosis and when combined with farnesoid X receptor loss protects against type 2 diabetes in mice. <i>Hepatology</i> , 2017 , 66, 1854-1865	11.2	22
122	Megatrends in bile acid receptor research. <i>Hepatology Communications</i> , 2017 , 1, 831-835	6	2
121	Plasma membrane-bound G protein-coupled bile acid receptor attenuates liver ischemia/reperfusion injury via the inhibition of toll-like receptor 4 signaling in mice. <i>Liver Transplantation</i> , 2017 , 23, 63-74	4.5	29
120	Inhibiting poly ADP-ribosylation increases fatty acid oxidation and protects against fatty liver disease. <i>Journal of Hepatology</i> , 2017 , 66, 132-141	13.4	80
119	FXR deficiency protects against obesity through a crosstalk between liver, microbiota, and brown adipose tissue. <i>JCI Insight</i> , 2017 , 2,	9.9	31
118	Impaired SUMOylation of nuclear receptor LRH-1 promotes nonalcoholic fatty liver disease. <i>Journal of Clinical Investigation</i> , 2017 , 127, 583-592	15.9	31
117	LRH-1-dependent programming of mitochondrial glutamine processing drives liver cancer. <i>Genes and Development</i> , 2016 , 30, 1255-60	12.6	41
116	Eliciting the mitochondrial unfolded protein response by nicotinamide adenine dinucleotide repletion reverses fatty liver disease in mice. <i>Hepatology</i> , 2016 , 63, 1190-204	11.2	223
115	Bile acid-FXR pathways regulate male sexual maturation in mice. <i>Oncotarget</i> , 2016 , 7, 19468-82	3.3	18
114	NAD ⁺ repletion improves mitochondrial and stem cell function and enhances life span in mice. <i>Science</i> , 2016 , 352, 1436-43	33.3	645
113	Farnesoid X receptor inhibits glucagon-like peptide-1 production by enteroendocrine L cells. <i>Nature Communications</i> , 2015 , 6, 7629	17.4	202
112	Bile Acids Trigger GLP-1 Release Predominantly by Accessing Basolaterally Located G Protein-Coupled Bile Acid Receptors. <i>Endocrinology</i> , 2015 , 156, 3961-70	4.8	199
111	TGR5 and Immunometabolism: Insights from Physiology and Pharmacology. <i>Trends in Pharmacological Sciences</i> , 2015 , 36, 847-857	13.2	91
110	LRH-1 mediates anti-inflammatory and antifungal phenotype of IL-13-activated macrophages through the PPAR α ligand synthesis. <i>Nature Communications</i> , 2015 , 6, 6801	17.4	35
109	Molecular basis for the regulation of the nuclear receptor LRH-1. <i>Current Opinion in Cell Biology</i> , 2015 , 33, 26-34	9	39
108	Phosphorylation of the nuclear receptor corepressor 1 by protein kinase B switches its corepressor targets in the liver in mice. <i>Hepatology</i> , 2015 , 62, 1606-18	11.2	37
107	The Sirt1 activator SRT3025 provides atheroprotection in Apoe ^{-/-} mice by reducing hepatic Pcsk9 secretion and enhancing Ldlr expression. <i>European Heart Journal</i> , 2015 , 36, 51-9	9.5	92
106	Intestinal FXR agonism promotes adipose tissue browning and reduces obesity and insulin resistance. <i>Nature Medicine</i> , 2015 , 21, 159-65	50.5	420

105	Nr5a2 heterozygosity sensitises to, and cooperates with, inflammation in KRas(G12V)-driven pancreatic tumourigenesis. <i>Gut</i> , 2014 , 63, 647-55	19.2	71
104	Pharmacological Inhibition of poly(ADP-ribose) polymerases improves fitness and mitochondrial function in skeletal muscle. <i>Cell Metabolism</i> , 2014 , 19, 1034-41	24.6	175
103	Hepatic glucose sensing and integrative pathways in the liver. <i>Cellular and Molecular Life Sciences</i> , 2014 , 71, 1453-67	10.3	65
102	Vitamin D and energy homeostasis: of mice and men. <i>Nature Reviews Endocrinology</i> , 2014 , 10, 79-87	15.2	83
101	A SIRT7-dependent acetylation switch of GABP β controls mitochondrial function. <i>Cell Metabolism</i> , 2014 , 20, 856-869	24.6	171
100	SUMOylation-dependent LRH-1/PROX1 interaction promotes atherosclerosis by decreasing hepatic reverse cholesterol transport. <i>Cell Metabolism</i> , 2014 , 20, 603-13	24.6	60
99	Another Shp on the horizon for bile acids. <i>Cell Metabolism</i> , 2014 , 20, 203-5	24.6	3
98	Loss of Sirt1 function improves intestinal anti-bacterial defense and protects from colitis-induced colorectal cancer. <i>PLoS ONE</i> , 2014 , 9, e102495	3.7	30
97	SIRT2 deficiency modulates macrophage polarization and susceptibility to experimental colitis. <i>PLoS ONE</i> , 2014 , 9, e103573	3.7	72
96	Bile acids alter male fertility through G-protein-coupled bile acid receptor 1 signaling pathways in mice. <i>Hepatology</i> , 2014 , 60, 1054-65	11.2	37
95	TGR5 reduces macrophage migration through mTOR-induced C/EBP β differential translation. <i>Journal of Clinical Investigation</i> , 2014 , 124, 5424-36	15.9	121
94	Liver receptor homolog-1 is essential for pregnancy. <i>Nature Medicine</i> , 2013 , 19, 1061-6	50.5	70
93	The NAD(+)/Sirtuin Pathway Modulates Longevity through Activation of Mitochondrial UPR and FOXO Signaling. <i>Cell</i> , 2013 , 154, 430-41	56.2	747
92	Transcriptional regulation of adipocyte formation by the liver receptor homologue 1 (Lrh1)-Small hetero-dimerization partner (Shp) network. <i>Molecular Metabolism</i> , 2013 , 2, 314-23	8.8	7
91	The receptor TGR5 mediates the prokinetic actions of intestinal bile acids and is required for normal defecation in mice. <i>Gastroenterology</i> , 2013 , 144, 145-54	13.3	198
90	The receptor TGR5 protects the liver from bile acid overload during liver regeneration in mice. <i>Hepatology</i> , 2013 , 58, 1451-60	11.2	132
89	Probing the Binding Site of Bile Acids in TGR5. <i>ACS Medicinal Chemistry Letters</i> , 2013 , 4, 1158-62	4.3	31
88	NR5A2 regulates Lhb and Fshb transcription in gonadotrope-like cells in vitro, but is dispensable for gonadotropin synthesis and fertility in vivo. <i>PLoS ONE</i> , 2013 , 8, e59058	3.7	17

87	The TGR5 receptor mediates bile acid-induced itch and analgesia. <i>Journal of Clinical Investigation</i> , 2013 , 123, 1513-30	15.9	229
86	TGR5, bile acids and the metabolic syndrome.. <i>FASEB Journal</i> , 2013 , 27, 453.1	0.9	
85	TGR5 potentiates GLP-1 secretion in response to anionic exchange resins. <i>Scientific Reports</i> , 2012 , 2, 430	4.9	126
84	Systems genetics of metabolism: the use of the BXD murine reference panel for multiscalar integration of traits. <i>Cell</i> , 2012 , 150, 1287-99	56.2	150
83	The NAD(+) precursor nicotinamide riboside enhances oxidative metabolism and protects against high-fat diet-induced obesity. <i>Cell Metabolism</i> , 2012 , 15, 838-47	24.6	732
82	Bile acid binding resin improves metabolic control through the induction of energy expenditure. <i>PLoS ONE</i> , 2012 , 7, e38286	3.7	81
81	Local glucocorticoid production in the mouse lung is induced by immune cell stimulation. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2012 , 67, 227-34	9.3	42
80	LRH-1-dependent glucose sensing determines intermediary metabolism in liver. <i>Journal of Clinical Investigation</i> , 2012 , 122, 2817-26	15.9	77
79	NCOR1 is a conserved physiological modulator of muscle mass and oxidative function. <i>Cell</i> , 2011 , 147, 827-39	56.2	170
78	PARP-1 inhibition increases mitochondrial metabolism through SIRT1 activation. <i>Cell Metabolism</i> , 2011 , 13, 461-468	24.6	555
77	Mitochondrial matrix calcium is an activating signal for hormone secretion. <i>Cell Metabolism</i> , 2011 , 13, 601-11	24.6	119
76	TGR5 activation inhibits atherosclerosis by reducing macrophage inflammation and lipid loading. <i>Cell Metabolism</i> , 2011 , 14, 747-57	24.6	364
75	Emerging actions of the nuclear receptor LRH-1 in the gut. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2011 , 1812, 947-55	6.9	55
74	The bile acid membrane receptor TGR5 as an emerging target in metabolism and inflammation. <i>Journal of Hepatology</i> , 2011 , 54, 1263-72	13.4	262
73	Lack of IL-2 in PPAR- δ deficient mice triggers allergic contact dermatitis by affecting regulatory T cells. <i>European Journal of Immunology</i> , 2011 , 41, 1980-91	6.1	15
72	Dual farnesoid X receptor/TGR5 agonist INT-767 reduces liver injury in the Mdr2 ^{-/-} (Abcb4 ^{-/-}) mouse cholangiopathy model by promoting biliary HCO ₃ ⁻ output. <i>Hepatology</i> , 2011 , 54, 1303-12	11.2	167
71	Lowering bile acid pool size with a synthetic farnesoid X receptor (FXR) agonist induces obesity and diabetes through reduced energy expenditure. <i>Journal of Biological Chemistry</i> , 2011 , 286, 26913-20	5.4	185
70	The bile acid membrane receptor TGR5: a valuable metabolic target. <i>Digestive Diseases</i> , 2011 , 29, 37-44	3.2	92

69	Reversible acetylation of PGC-1: connecting energy sensors and effectors to guarantee metabolic flexibility. <i>Oncogene</i> , 2010 , 29, 4617-24	9.2	126
68	Lipopolysaccharide induces intestinal glucocorticoid synthesis in a TNFalpha-dependent manner. <i>FASEB Journal</i> , 2010 , 24, 1340-6	0.9	30
67	Mitochondrial matrix pH controls oxidative phosphorylation and metabolism-secretion coupling in INS-1E clonal beta cells. <i>FASEB Journal</i> , 2010 , 24, 4613-26	0.9	45
66	The intestinal nuclear receptor signature with epithelial localization patterns and expression modulation in tumors. <i>Gastroenterology</i> , 2010 , 138, 636-48, 648.e1-12	13.3	65
65	Histone methyl transferases and demethylases; can they link metabolism and transcription?. <i>Cell Metabolism</i> , 2010 , 12, 321-327	24.6	204
64	Structure-activity relationship study of betulinic acid, a novel and selective TGR5 agonist, and its synthetic derivatives: potential impact in diabetes. <i>Journal of Medicinal Chemistry</i> , 2010 , 53, 178-90	8.3	159
63	Targeting the TGR5-GLP-1 pathway to combat type 2 diabetes and non-alcoholic fatty liver disease. <i>Gastroenterologie Clinique Et Biologique</i> , 2010 , 34, 270-3		12
62	Redefining the TGR5 triterpenoid binding pocket at the C-3 position. <i>ChemMedChem</i> , 2010 , 5, 1983-8	3.7	20
61	Raised hepatic bile acid concentrations during pregnancy in mice are associated with reduced farnesoid X receptor function. <i>Hepatology</i> , 2010 , 52, 1341-9	11.2	68
60	Discovery of 6alpha-ethyl-23(S)-methylcholic acid (S-EMCA, INT-777) as a potent and selective agonist for the TGR5 receptor, a novel target for diabetes. <i>Journal of Medicinal Chemistry</i> , 2009 , 52, 7958-61	8.3	194
59	TGR5-mediated bile acid sensing controls glucose homeostasis. <i>Cell Metabolism</i> , 2009 , 10, 167-77	24.6	1184
58	The orphan nuclear receptor small heterodimer partner mediates male infertility induced by diethylstilbestrol in mice. <i>Journal of Clinical Investigation</i> , 2009 , 119, 3752-64	15.9	47
57	Targeting bile-acid signalling for metabolic diseases. <i>Nature Reviews Drug Discovery</i> , 2008 , 7, 678-93	64.1	864
56	Bile acids and the membrane bile acid receptor TGR5--connecting nutrition and metabolism. <i>Thyroid</i> , 2008 , 18, 167-74	6.2	123
55	Molecular field analysis and 3D-quantitative structure-activity relationship study (MFA 3D-QSAR) unveil novel features of bile acid recognition at TGR5. <i>Journal of Chemical Information and Modeling</i> , 2008 , 48, 1792-801	6.1	20
54	Cholesterol supply and SREBPs modulate transcription of the Niemann-Pick C-1 gene in steroidogenic tissues. <i>Journal of Lipid Research</i> , 2008 , 49, 1024-33	6.3	22
53	Liver receptor homolog 1 is essential for ovulation. <i>Genes and Development</i> , 2008 , 22, 1871-6	12.6	150
52	The genetic ablation of SRC-3 protects against obesity and improves insulin sensitivity by reducing the acetylation of PGC-1{alpha}. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 17187-92	11.5	158

51	Cell cycle-dependent regulation of extra-adrenal glucocorticoid synthesis in murine intestinal epithelial cells. <i>FASEB Journal</i> , 2008 , 22, 4117-25	0.9	29
50	Structure-based design of a superagonist ligand for the vitamin D nuclear receptor. <i>Chemistry and Biology</i> , 2008 , 15, 383-92		42
49	Novel potent and selective bile acid derivatives as TGR5 agonists: biological screening, structure-activity relationships, and molecular modeling studies. <i>Journal of Medicinal Chemistry</i> , 2008 , 51, 1831-41	8.3	218
48	Sirtuin functions in health and disease. <i>Molecular Endocrinology</i> , 2007 , 21, 1745-55		343
47	Nongenomic actions of bile acids. Synthesis and preliminary characterization of 23- and 6,23-alkyl-substituted bile acid derivatives as selective modulators for the G-protein coupled receptor TGR5. <i>Journal of Medicinal Chemistry</i> , 2007 , 50, 4265-8	8.3	89
46	Peroxisome proliferator-activated receptor-alpha activation inhibits Langerhans cell function. <i>Journal of Immunology</i> , 2007 , 178, 4362-72	5.3	32
45	Compromised intestinal lipid absorption in mice with a liver-specific deficiency of liver receptor homolog 1. <i>Molecular and Cellular Biology</i> , 2007 , 27, 8330-9	4.8	127
44	Adipose tissue-specific inactivation of the retinoblastoma protein protects against diabetes because of increased energy expenditure. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 10703-8	11.5	85
43	LRH-1-mediated glucocorticoid synthesis in enterocytes protects against inflammatory bowel disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 13098-103	11.5	111
42	The small heterodimer partner is a gonadal gatekeeper of sexual maturation in male mice. <i>Genes and Development</i> , 2007 , 21, 303-15	12.6	70
41	Sirtuins: the magnificent seven function, metabolism and longevity. <i>Annals of Medicine</i> , 2007 , 39, 335-45	4.5	353
40	The nuclear receptor LRH-1 critically regulates extra-adrenal glucocorticoid synthesis in the intestine. <i>Journal of Experimental Medicine</i> , 2006 , 203, 2057-62	16.6	92
39	Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. <i>Nature</i> , 2006 , 439, 484-9	50.4	1508
38	Liver receptor homolog 1 contributes to intestinal tumor formation through effects on cell cycle and inflammation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 2058-62	11.5	116
37	Adipose tissue expression of the lipid droplet-associating proteins S3-12 and perilipin is controlled by peroxisome proliferator-activated receptor-gamma. <i>Diabetes</i> , 2004 , 53, 1243-52	0.9	164
36	Peroxisome proliferator-activated receptor (PPAR)-beta/delta stimulates differentiation and lipid accumulation in keratinocytes. <i>Journal of Investigative Dermatology</i> , 2004 , 122, 971-83	4.3	177
35	LRH-1: an orphan nuclear receptor involved in development, metabolism and steroidogenesis. <i>Trends in Cell Biology</i> , 2004 , 14, 250-60	18.3	340
34	Synergy between LRH-1 and beta-catenin induces G1 cyclin-mediated cell proliferation. <i>Molecular Cell</i> , 2004 , 15, 499-509	17.6	238

33	Pancreatic-duodenal homeobox 1 regulates expression of liver receptor homolog 1 during pancreas development. <i>Molecular and Cellular Biology</i> , 2003 , 23, 6713-24	4.8	80
32	Liver receptor homolog 1 controls the expression of carboxyl ester lipase. <i>Journal of Biological Chemistry</i> , 2003 , 278, 35725-31	5.4	44
31	Topical peroxisome proliferator activated receptor-alpha activators reduce inflammation in irritant and allergic contact dermatitis models. <i>Journal of Investigative Dermatology</i> , 2002 , 118, 94-101	4.3	140
30	Role of peroxisome proliferator-activated receptor alpha in epidermal development in utero. <i>Journal of Investigative Dermatology</i> , 2002 , 119, 1298-303	4.3	39
29	The role of PPAR-gamma/RXR-alpha heterodimers in the regulation of human trophoblast invasion. <i>Annals of the New York Academy of Sciences</i> , 2002 , 973, 26-30	6.5	47
28	Liver receptor homolog 1 controls the expression of the scavenger receptor class B type I. <i>EMBO Reports</i> , 2002 , 3, 1181-7	6.5	117
27	Progesterone receptor knockout mice have an improved glucose homeostasis secondary to beta-cell proliferation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 15644-8	11.5	115
26	The small heterodimer partner interacts with the liver X receptor alpha and represses its transcriptional activity. <i>Molecular Endocrinology</i> , 2002 , 16, 2065-76		178
25	Attenuation of colon inflammation through activators of the retinoid X receptor (RXR)/peroxisome proliferator-activated receptor gamma (PPARgamma) heterodimer. A basis for new therapeutic strategies. <i>Journal of Experimental Medicine</i> , 2001 , 193, 827-38	16.6	371
24	XoL INXS: role of the liver X and the farnesol X receptors. <i>Current Opinion in Lipidology</i> , 2001 , 12, 113-20	4.4	21
23	PPAR/RXR Heterodimers Control Human Trophoblast Invasion. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 5017-5024	5.6	75
22	Expression of peroxisome proliferator-activated receptor gamma (PPARgamma) in normal human pancreatic islet cells. <i>Diabetologia</i> , 2000 , 43, 1165-9	10.3	171
21	Farnesol stimulates differentiation in epidermal keratinocytes via PPARalpha. <i>Journal of Biological Chemistry</i> , 2000 , 275, 11484-91	5.4	88
20	Induction of LPL gene expression by sterols is mediated by a sterol regulatory element and is independent of the presence of multiple E boxes. <i>Journal of Molecular Biology</i> , 2000 , 304, 323-34	6.5	65
19	Molecular basis for feedback regulation of bile acid synthesis by nuclear receptors. <i>Molecular Cell</i> , 2000 , 6, 507-15	17.6	1195
18	Thiazolidinediones: an update. <i>Lancet, The</i> , 2000 , 355, 1008-10	4.0	174
17	3-Hydroxy-3-methylglutaryl CoA reductase inhibitors reduce serum triglyceride levels through modulation of apolipoprotein C-III and lipoprotein lipase. <i>FEBS Letters</i> , 1999 , 452, 160-4	3.8	68
16	Regulation of peroxisome proliferator-activated receptor gamma expression by adipocyte differentiation and determination factor 1/sterol regulatory element binding protein 1: implications for adipocyte differentiation and metabolism. <i>Molecular and Cellular Biology</i> , 1999 , 19, 5495-503	4.8	356

15	Mechanism of action of fibrates on lipid and lipoprotein metabolism. <i>Circulation</i> , 1998 , 98, 2088-93	16.7	1322
14	Transcriptional regulation of apolipoprotein A-I gene expression by the nuclear receptor RORalpha. <i>Journal of Biological Chemistry</i> , 1997 , 272, 22401-4	5.4	100
13	The organization, promoter analysis, and expression of the human PPARgamma gene. <i>Journal of Biological Chemistry</i> , 1997 , 272, 18779-89	5.4	889
12	Coordinate regulation of the expression of the fatty acid transport protein and acyl-CoA synthetase genes by PPARalpha and PPARgamma activators. <i>Journal of Biological Chemistry</i> , 1997 , 272, 28210-7	5.4	404
11	Peroxisome proliferator-activated receptors, orphans with ligands and functions. <i>Current Opinion in Lipidology</i> , 1997 , 8, 159-66	4.4	395
10	The effects of fibrates and thiazolidinediones on plasma triglyceride metabolism are mediated by distinct peroxisome proliferator activated receptors (PPARs). <i>Biochimie</i> , 1997 , 79, 95-9	4.6	65
9	Regulation of triglyceride metabolism by PPARs: fibrates and thiazolidinediones have distinct effects. <i>Journal of Atherosclerosis and Thrombosis</i> , 1996 , 3, 81-9	4	89
8	Retinoids increase human apolipoprotein A-11 expression through activation of the retinoid X receptor but not the retinoic acid receptor. <i>Molecular and Cellular Biology</i> , 1996 , 16, 3350-60	4.8	51
7	Induction of the acyl-coenzyme A synthetase gene by fibrates and fatty acids is mediated by a peroxisome proliferator response element in the C promoter. <i>Journal of Biological Chemistry</i> , 1995 , 270, 19269-76	5.4	303
6	Fibrates increase human apolipoprotein A-II expression through activation of the peroxisome proliferator-activated receptor. <i>Journal of Clinical Investigation</i> , 1995 , 96, 741-50	15.9	302
5	A Fos-Jun element in the first intron of an alpha 2u-globulin gene. <i>Molecular and Cellular Biochemistry</i> , 1993 , 125, 127-36	4.2	5
4	Acyl-CoA synthetase mRNA expression is controlled by fibric-acid derivatives, feeding and liver proliferation. <i>FEBS Journal</i> , 1993 , 216, 615-22		91
3	Developmental extinction of liver lipoprotein lipase mRNA expression might be regulated by an NF-1-like site. <i>FEBS Letters</i> , 1993 , 329, 89-95	3.8	18
2	Lipoprotein lipase: recent contributions from molecular biology. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 1992 , 29, 243-68	9.4	122
1	New insights into apolipoprotein B and low density lipoprotein physiology; implications for atherosclerosis. <i>Acta Clinica Belgica</i> , 1991 , 46, 355-8	1.8	3