Hermes Licea-Perez

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/778206/hermes-licea-perez-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

29	513	13	22
papers	citations	h-index	g-index
29	579 ext. citations	3.3	3.53
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
29	Chemical derivatization in combination with supercritical fluid chromatography to improve resolution of stereoisomers. <i>Bioanalysis</i> , 2021 , 13, 985-999	2.1	1
28	Strategies for effective development of ultra-sensitive LC-MS/MS assays: application to a novel STING agonist. <i>Bioanalysis</i> , 2020 , 12, 467-484	2.1	1
27	Multicenter Validation Study of Quantitative Imaging Mass Spectrometry. <i>Analytical Chemistry</i> , 2019 , 91, 6266-6274	7.8	33
26	The importance of evaluating the chemical structures and strategies to avoid pitfalls in quantitative bioanalysis. <i>Bioanalysis</i> , 2019 , 11, 85-101	2.1	2
25	Development of an ultra-sensitive assay for the determination of an aminoalkyl glucosaminide 4-phosphate, GSK1795091, in plasma to support a first time in human study. <i>Analytical Methods</i> , 2018 , 10, 3074-3080	3.2	3
24	New insights into supercritical fluid chromatography for chiral separations. <i>Analytical Methods</i> , 2017 , 9, 2603-2610	3.2	6
23	Overcoming challenges associated with the bioanalysis of an ester prodrug and its active acid metabolite. <i>Bioanalysis</i> , 2017 , 9, 1589-1601	2.1	3
22	LC-MS/MS quantification of dimethyl fumarate and methyl hydrogen fumarate in rat blood using tiopronin as trapping reagent. <i>Analytical Methods</i> , 2016 , 8, 6420-6427	3.2	3
21	Overcoming bioanalytical challenges associated with the separation and quantitation of GSK1278863, a HIF-prolyl hydroxylase inhibitor, and its 14 stereoisomeric metabolites. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016 , 1009-1010, 7-16	3.2	11
20	Mass spectrometric characterization of a prolyl hydroxylase inhibitor GSK1278863, its bishydroxylated metabolite, and its implementation into routine doping controls. <i>Drug Testing and Analysis</i> , 2016 , 8, 858-63	3.5	8
19	Development of a multi-sugar LC-MS/MS assay using simple chemical derivatization with acetic anhydride. <i>Analytical Methods</i> , 2016 , 8, 3023-3033	3.2	4
18	Chemical derivatization in bioanalysis. <i>Bioanalysis</i> , 2015 , 7, 2435-7	2.1	10
17	Camphanic acid chloride: a powerful derivatization reagent for stereoisomeric separation and its DMPK applications. <i>Bioanalysis</i> , 2015 , 7, 3005-17	2.1	10
16	Development and validation of a quantitative method for determination of retigabine and its N-acetyl metabolite; overcoming challenges associated with circulating labile N-glucuronide metabolites. <i>Analytical Methods</i> , 2015 , 7, 723-735	3.2	7
15	Analytical approaches for quantification of a Nrf2 pathway activator: overcoming bioanalytical challenges to support a toxicity study. <i>Analyst, The</i> , 2014 , 139, 1902-12	5	5
14	Development and validation of a simple and sensitive method for quantification of levodopa and carbidopa in rat and monkey plasma using derivatization and UPLC-MS/MS. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013 , 926, 47-53	3.2	13
13	A novel approach to capillary plasma microsampling for quantitative bioanalysis. <i>Bioanalysis</i> , 2013 , 5, 1131-5	2.1	48

LIST OF PUBLICATIONS

12	Evaluation of glucuronide metabolite stability in dried blood spots. <i>Bioanalysis</i> , 2012 , 4, 2823-32	2.1	24
11	Development of a semi-automated LC/MS/MS method for the simultaneous quantitation of 14,15-epoxyeicosatrienoic acid, 14,15-dihydroxyeicosatrienoic acid, leukotoxin and leukotoxin diol in human plasma as biomarkers of soluble epoxide hydrolase activity in vivo. <i>Journal of</i>	3.2	14
10	Development of a sensitive and selective LC-MS/MS method for the determination of alpha-fluoro-beta-alanine, 5-fluorouracil and capecitabine in human plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009 , 877, 1040-6	3.2	43
9	Development of a sensitive and selective LC-MS/MS method for simultaneous determination of gemcitabine and 2,2-difluoro-2-deoxyuridine in human plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009 , 877, 2123-9	3.2	38
8	A semi-automated 96-well plate method for the simultaneous determination of oral contraceptives concentrations in human plasma using ultra performance liquid chromatography coupled with tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical</i>	3.2	40
7	and Life Sciences, 2007, 852, 69-76 Dosimetry by means of DNA and hemoglobin adducts in propylene oxide-exposed rats. <i>Toxicology and Applied Pharmacology</i> , 2003, 191, 245-54	4.6	20
6	A sensitive gas chromatographic-tandem mass spectrometric method for detection of alkylating agents in water: application to acrylamide in drinking water, coffee and snuff. <i>Analyst, The</i> , 2003 , 128, 1033-6	5	27
5	Determination of styrene and styrene-7,8-oxide in human blood by gas chromatography-mass spectrometry. <i>Biomedical Applications</i> , 2001 , 757, 59-68		14
4	Biotransformation of the double bond in allyl glycidyl ether to an epoxide ring. Evidence from hemoglobin adducts in mice. <i>Chemico-Biological Interactions</i> , 2000 , 125, 17-28	5	1
3	Simultaneous analysis of hemoglobin adducts of acrylamide and glycidamide by gas chromatography-mass spectrometry. <i>Analytical Biochemistry</i> , 1999 , 274, 59-68	3.1	55
2	Adducts of acrylonitrile with hemoglobin in nonsmokers and in participants in a smoking cessation program. <i>Chemical Research in Toxicology</i> , 1999 , 12, 869-73	4	15
1	Haemoglobin adducts of epoxybutanediol from exposure to 1,3-butadiene or butadiene epoxides. <i>Chemico-Biological Interactions</i> , 1997 , 105, 181-98	5	54