

# Takeshi Asami

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7788676/publications.pdf>

Version: 2024-02-01

21  
papers

784  
citations

623188

14  
h-index

752256

20  
g-index

21  
all docs

21  
docs citations

21  
times ranked

1493  
citing authors

#	ARTICLE	IF	CITATIONS
1	Structural brain abnormalities in adolescent patients with anorexia nervosa at both the acute and weight-recovered phase. <i>Brain Imaging and Behavior</i> , 2022, 16, 1372-1380.	1.1	2
2	Smaller volume of right hippocampal CA2/3 in patients with panic disorder. <i>Brain Imaging and Behavior</i> , 2021, 15, 320-326.	1.1	9
3	The psychological effects of COVID-19 on hospital workers at the beginning of the outbreak with a large disease cluster on the Diamond Princess cruise ship. <i>PLoS ONE</i> , 2021, 16, e0245294.	1.1	17
4	Illness management and recovery program induced neuroprotective effects on language network in schizophrenia. <i>Schizophrenia Research</i> , 2021, 230, 101-103.	1.1	1
5	Exploratory investigation on antibodies to GluN1 and cognitive dysfunction in patients with chronic autoimmune psychosis. <i>Neuroscience Letters</i> , 2021, 743, 135588.	1.0	1
6	Structural abnormalities in nucleus accumbens in patients with panic disorder. <i>Journal of Affective Disorders</i> , 2020, 271, 201-206.	2.0	3
7	Clinical and brain structural effects of the Illness Management and Recovery program in middle-aged and older patients with schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2019, 73, 731-737.	1.0	5
8	Progressive symptom-associated prefrontal volume loss occurs in first-episode schizophrenia but not in affective psychosis. <i>Brain Structure and Function</i> , 2018, 223, 2879-2892.	1.2	16
9	Thalamic shape and volume abnormalities in female patients with panic disorder. <i>PLoS ONE</i> , 2018, 13, e0208152.	1.1	23
10	Smaller volumes in the lateral and basal nuclei of the amygdala in patients with panic disorder. <i>PLoS ONE</i> , 2018, 13, e0207163.	1.1	27
11	Cortical thickness reductions in the middle frontal cortex in patients with panic disorder. <i>Journal of Affective Disorders</i> , 2018, 240, 199-202.	2.0	14
12	Multiple White Matter Volume Reductions in Patients with Panic Disorder: Relationships between Orbitofrontal Gyrus Volume and Symptom Severity and Social Dysfunction. <i>PLoS ONE</i> , 2014, 9, e92862.	1.1	15
13	Prefrontal cortex volume deficit in schizophrenia: A new look using 3T MRI with manual parcellation. <i>Schizophrenia Research</i> , 2014, 152, 184-190.	1.1	30
14	Globally and Locally Reduced MRI Gray Matter Volumes in Neuroleptic-Naive Men With Schizotypal Personality Disorder. <i>JAMA Psychiatry</i> , 2013, 70, 361.	6.0	35
15	Abnormalities of middle longitudinal fascicle and disorganization in patients with schizophrenia. <i>Schizophrenia Research</i> , 2013, 143, 253-259.	1.1	36
16	Longitudinal loss of gray matter volume in patients with first-episode schizophrenia: DARTEL automated analysis and ROI validation. <i>NeuroImage</i> , 2012, 59, 986-996.	2.1	129
17	Midbrain volume increase in patients with panic disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2011, 65, 365-373.	1.0	35
18	Posterior orbitofrontal sulcogyral pattern associated with orbitofrontal cortex volume reduction and anxiety trait in panic disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2010, 64, 318-326.	1.0	54

#	ARTICLE	IF	CITATIONS
19	Sexually dimorphic gray matter volume reduction in patients with panic disorder. <i>Psychiatry Research - Neuroimaging</i> , 2009, 173, 128-134.	0.9	95
20	Smaller amygdala is associated with anxiety in patients with panic disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2009, 63, 266-276.	1.0	141
21	Anterior cingulate cortex volume reduction in patients with panic disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2008, 62, 322-330.	1.0	96