

# Timo Mttnen

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/7790502/timo-mottonen-publications-by-citations.pdf>

**Version:** 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

13  
papers

753  
citations

10  
h-index

13  
g-index

13  
ext. papers

857  
ext. citations

3.2  
avg, IF

2.48  
L-index

#	Paper	IF	Citations
13	Delay to institution of therapy and induction of remission using single-drug or combination-disease-modifying antirheumatic drug therapy in early rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , <b>2002</b> , 46, 894-8		274
12	Combination drug therapy retards the development of rheumatoid atlantoaxial subluxations. <i>Arthritis and Rheumatism</i> , <b>2000</b> , 43, 2397-401		73
11	Single values of serum transferrin receptor and transferrin receptor ferritin index can be used to detect true and functional iron deficiency in rheumatoid arthritis patients with anemia. <i>Arthritis and Rheumatism</i> , <b>2000</b> , 43, 1016-20		67
10	Salmonella-triggered reactive arthritis: use of polymerase chain reaction, immunocytochemical staining, and gas chromatography-mass spectrometry in the detection of bacterial components from synovial fluid. <i>Arthritis and Rheumatism</i> , <b>1999</b> , 42, 84-9		67
9	Infliximab for 6 months added on combination therapy in early rheumatoid arthritis: 2-year results from an investigator-initiated, randomised, double-blind, placebo-controlled study (the NEO-RACo Study). <i>Annals of the Rheumatic Diseases</i> , <b>2013</b> , 72, 851-7	2.4	65
8	Outcome in patients with early rheumatoid arthritis treated according to the "sawtooth" strategy. <i>Arthritis and Rheumatism</i> , <b>1996</b> , 39, 996-1005		54
7	Detection of herpesviruses by polymerase chain reaction in lymphocytes from patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , <b>1993</b> , 36, 1080-6		48
6	Rheumatoid atlantoaxial subluxation can be prevented by intensive use of traditional disease modifying antirheumatic drugs. <i>Journal of Rheumatology</i> , <b>2009</b> , 36, 273-8	4.1	37
5	Targeted treatment with a combination of traditional DMARDs produces excellent clinical and radiographic long-term outcomes in early rheumatoid arthritis regardless of initial infliximab. The 5-year follow-up results of a randomised clinical trial, the NEO-RACo trial. <i>Annals of the Rheumatic Diseases</i> , <b>2017</b> , 76, 1071-6	2.4	31
4	Glycoprotein YKL-40: A potential biomarker of disease activity in rheumatoid arthritis during intensive treatment with csDMARDs and infliximab. Evidence from the randomised controlled NEO-RACo trial. <i>PLoS ONE</i> , <b>2017</b> , 12, e0183294	3.7	20
3	Cervical Spine Involvement among Patients with Rheumatoid Arthritis Treated Actively with Treat-to-target Strategy: 10-year Results of the NEO-RACo Study. <i>Journal of Rheumatology</i> , <b>2020</b> , 47, 1160-1164	4.1	9
2	Early treatment intensification induces favourable radiographic outcomes according to predicted versus observed radiographic progression in early rheumatoid arthritis: a subanalysis of the randomised FIN-RACo and NEO-RACo trials. <i>Clinical and Experimental Rheumatology</i> , <b>2016</b> , 34, 1065-1071	2.2	5
1	Early Targeted Combination Treatment With Conventional Synthetic Disease-Modifying Antirheumatic Drugs and Long-Term Outcomes in Rheumatoid Arthritis: Ten-Year Follow-Up Results of a Randomized Clinical Trial. <i>Arthritis Care and Research</i> , <b>2019</b> , 71, 1450-1458	4.7	3