

Lars Rnnstrand

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7791521/lars-ronnstrand-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140
papers

6,715
citations

44
h-index

78
g-index

155
ext. papers

7,367
ext. citations

6.3
avg, IF

5.81
L-index

#	Paper	IF	Citations
140	Metallopeptidase inhibitor 1 (TIMP-1) promotes receptor tyrosine kinase c-Kit signaling in colorectal cancer. <i>Molecular Oncology</i> , 2019 , 13, 2646-2662	7.9	5
139	FMS-like Tyrosine Kinase 3/FLT3: From Basic Science to Clinical Implications. <i>Physiological Reviews</i> , 2019 , 99, 1433-1466	47.9	42
138	The role of SRC family kinases in FLT3 signaling. <i>International Journal of Biochemistry and Cell Biology</i> , 2019 , 107, 32-37	5.6	12
137	The ALK inhibitor AZD3463 effectively inhibits growth of sorafenib-resistant acute myeloid leukemia. <i>Blood Cancer Journal</i> , 2019 , 9, 5	7	4
136	Internal tandem duplication mutations in the tyrosine kinase domain of FLT3 display a higher oncogenic potential than the activation loop D835Y mutation. <i>Annals of Hematology</i> , 2018 , 97, 773-780	3	7
135	SRC-like adaptor protein 2 (SLAP2) is a negative regulator of KIT-D816V-mediated oncogenic transformation. <i>Scientific Reports</i> , 2018 , 8, 6405	4.9	1
134	Bruton's tyrosine kinase potentiates ALK signaling and serves as a potential therapeutic target of neuroblastoma. <i>Oncogene</i> , 2018 , 37, 6180-6194	9.2	15
133	De novo activating mutations drive clonal evolution and enhance clonal fitness in KMT2A-rearranged leukemia. <i>Nature Communications</i> , 2018 , 9, 1770	17.4	26
132	XK-related protein 5 (XKR5) is a novel negative regulator of KIT/D816V-mediated transformation. <i>Oncogenesis</i> , 2018 , 7, 48	6.6	2
131	Tyrosine 842 in the activation loop is required for full transformation by the oncogenic mutant FLT3-ITD. <i>Cellular and Molecular Life Sciences</i> , 2017 , 74, 2679-2688	10.3	11
130	KIT Induces SRC-Mediated Tyrosine Phosphorylation of MITF and Altered Transcription Program in Melanoma. <i>Molecular Cancer Research</i> , 2017 , 15, 1265-1274	6.6	11
129	The Src family kinase LCK cooperates with oncogenic FLT3/ITD in cellular transformation. <i>Scientific Reports</i> , 2017 , 7, 13734	4.9	13
128	ABL2 suppresses FLT3-ITD-induced cell proliferation through negative regulation of AKT signaling. <i>Oncotarget</i> , 2017 , 8, 12194-12202	3.3	12
127	Internal Tandem Duplication (ITD) in the Tyrosine Kinase Domain of FLT3 Displays Higher Oncogenic Potential in Acute Myeloid Leukemia. <i>Blood</i> , 2016 , 128, 5118-5118	2.2	1
126	Src-like adaptor protein 2 (SLAP2) binds to and inhibits FLT3 signaling. <i>Oncotarget</i> , 2016 , 7, 57770-57783	3.3	9
125	FYN expression potentiates FLT3-ITD induced STAT5 signaling in acute myeloid leukemia. <i>Oncotarget</i> , 2016 , 7, 9964-74	3.3	26
124	HIF2 α contributes to antiestrogen resistance via positive bilateral crosstalk with EGFR in breast cancer cells. <i>Oncotarget</i> , 2016 , 7, 11238-50	3.3	12

123	Expression of GADS enhances FLT3-induced mitogenic signaling. <i>Oncotarget</i> , 2016 , 7, 14112-24	3.3	10
122	Tyrosine 842 Residue in the Activation Loop of FLT3-ITD Is Indispensable for Oncogenic Transformation. <i>Blood</i> , 2016 , 128, 1561-1561	2.2	
121	Loss of Src-like Adaptor Protein 2 Expression Increases the Transforming Potential of Oncogenic FLT3-ITD. <i>Blood</i> , 2016 , 128, 5106-5106	2.2	
120	Role of SRC-like adaptor protein (SLAP) in immune and malignant cell signaling. <i>Cellular and Molecular Life Sciences</i> , 2015 , 72, 2535-44	10.3	16
119	Brain-Expressed X-linked (BEX) proteins in human cancers. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 2015 , 1856, 226-33	11.2	17
118	The PDGFR Receptor Family 2015 , 373-538		2
117	The Phosphatases STS1 and STS2 Regulate Hematopoietic Stem and Progenitor Cell Fitness. <i>Stem Cell Reports</i> , 2015 , 5, 633-46	8	7
116	PI3 kinase is indispensable for oncogenic transformation by the V560D mutant of c-Kit in a kinase-independent manner. <i>Cellular and Molecular Life Sciences</i> , 2015 , 72, 4399-407	10.3	7
115	The role of HOXB2 and HOXB3 in acute myeloid leukemia. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 467, 742-7	3.4	27
114	BEX1 acts as a tumor suppressor in acute myeloid leukemia. <i>Oncotarget</i> , 2015 , 6, 21395-405	3.3	18
113	Aberrant Activation of the PI3K/mTOR Pathway Promotes Resistance to Sorafenib in AML. <i>Blood</i> , 2015 , 126, 2472-2472	2.2	
112	SOCS proteins in regulation of receptor tyrosine kinase signaling. <i>Cellular and Molecular Life Sciences</i> , 2014 , 71, 3297-310	10.3	63
111	SYK is a critical regulator of FLT3 in acute myeloid leukemia. <i>Cancer Cell</i> , 2014 , 25, 226-42	24.3	101
110	EPO-independent functional EPO receptor in breast cancer enhances estrogen receptor activity and promotes cell proliferation. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 445, 163-9	3.4	10
109	Src-like-adaptor protein (SLAP) differentially regulates normal and oncogenic c-Kit signaling. <i>Journal of Cell Science</i> , 2014 , 127, 653-62	5.3	24
108	Keratin 19 expression correlates with poor prognosis in breast cancer. <i>Molecular Biology Reports</i> , 2014 , 41, 7729-35	2.8	29
107	SOCS6 is a selective suppressor of receptor tyrosine kinase signaling. <i>Tumor Biology</i> , 2014 , 35, 10581-9	2.9	25
106	The basic helix-loop-helix (bHLH) proteins in breast cancer progression. <i>Medical Oncology</i> , 2013 , 30, 666	3.7	2

105	Enhanced SOX10 and KIT expression in cutaneous melanoma. <i>Medical Oncology</i> , 2013 , 30, 648	3.7	7
104	Deregulation of protein phosphatase expression in acute myeloid leukemia. <i>Medical Oncology</i> , 2013 , 30, 517	3.7	18
103	Protein kinase C (PKC) as a drug target in chronic lymphocytic leukemia. <i>Medical Oncology</i> , 2013 , 30, 7573.7	3.7	24
102	Protein kinase C expression is deregulated in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2013 , 54, 2288-90	1.9	12
101	Differential activity of c-KIT splice forms is controlled by extracellular peptide insert length. <i>Cellular Signalling</i> , 2013 , 25, 2231-8	4.9	5
100	The presence or absence of IL-3 during long-term culture of Flt3-ITD and c-Kit-D816V expressing Ba/F3 cells influences signaling outcome. <i>Experimental Hematology</i> , 2013 , 41, 585-7	3.1	15
99	The tyrosine kinase CSK associates with FLT3 and c-Kit receptors and regulates downstream signaling. <i>Cellular Signalling</i> , 2013 , 25, 1852-60	4.9	25
98	FLT3 mutations in patients with childhood acute lymphoblastic leukemia (ALL). <i>Medical Oncology</i> , 2013 , 30, 462	3.7	18
97	Suppressor of cytokine signaling 2 (SOCS2) associates with FLT3 and negatively regulates downstream signaling. <i>Molecular Oncology</i> , 2013 , 7, 693-703	7.9	49
96	FLT3 signals via the adapter protein Grb10 and overexpression of Grb10 leads to aberrant cell proliferation in acute myeloid leukemia. <i>Molecular Oncology</i> , 2013 , 7, 402-18	7.9	34
95	Phosphorylation of the activation loop tyrosine 823 in c-Kit is crucial for cell survival and proliferation. <i>Journal of Biological Chemistry</i> , 2013 , 288, 22460-8	5.4	27
94	3,4-Diarylmaleimides-a novel class of kinase inhibitors-effectively induce apoptosis in FLT3-ITD-dependent cells. <i>Annals of Hematology</i> , 2012 , 91, 331-44	3	4
93	SRC is a signaling mediator in FLT3-ITD- but not in FLT3-TKD-positive AML. <i>Blood</i> , 2012 , 119, 4026-33	2.2	47
92	Adaptor protein Lnk binds to and inhibits normal and leukemic FLT3. <i>Blood</i> , 2012 , 120, 3310-7	2.2	33
91	Stem cell factor receptor/c-Kit: from basic science to clinical implications. <i>Physiological Reviews</i> , 2012 , 92, 1619-49	47.9	453
90	HIF-2 α expression is suppressed in SCLC cells, which survive in moderate and severe hypoxia when HIF-1 α s repressed. <i>American Journal of Pathology</i> , 2012 , 180, 494-504	5.8	16
89	Suppressor of cytokine signaling 6 (SOCS6) negatively regulates Flt3 signal transduction through direct binding to phosphorylated tyrosines 591 and 919 of Flt3. <i>Journal of Biological Chemistry</i> , 2012 , 287, 36509-17	5.4	49
88	Src-Like adaptor protein (SLAP) binds to the receptor tyrosine kinase Flt3 and modulates receptor stability and downstream signaling. <i>PLoS ONE</i> , 2012 , 7, e53509	3.7	33

87	Inhibition of MEK5 by BIX02188 induces apoptosis in cells expressing the oncogenic mutant FLT3-ITD. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 412, 307-12	3.4	24
86	Impact of gene dosage, loss of wild-type allele, and FLT3 ligand on Flt3-ITD-induced myeloproliferation. <i>Blood</i> , 2011 , 118, 3613-21	2.2	25
85	Irreversible pan-ERBB inhibitor canertinib elicits anti-leukaemic effects and induces the regression of FLT3-ITD transformed cells in mice. <i>British Journal of Haematology</i> , 2011 , 155, 198-208	4.5	7
84	Protein-tyrosine phosphatase DEP-1 controls receptor tyrosine kinase FLT3 signaling. <i>Journal of Biological Chemistry</i> , 2011 , 286, 10918-29	5.4	51
83	Structural basis for c-KIT inhibition by the suppressor of cytokine signaling 6 (SOCS6) ubiquitin ligase. <i>Journal of Biological Chemistry</i> , 2011 , 286, 480-90	5.4	46
82	C-KIT signaling depends on microphthalmia-associated transcription factor for effects on cell proliferation. <i>PLoS ONE</i> , 2011 , 6, e24064	3.7	32
81	Signaling by the Platelet-Derived Growth Factor Receptor Family 2010 , 427-434		4
80	The D816V mutation of c-Kit circumvents a requirement for Src family kinases in c-Kit signal transduction. <i>Journal of Biological Chemistry</i> , 2009 , 284, 11039-47	5.4	54
79	A novel molecular mechanism of primary resistance to FLT3-kinase inhibitors in AML. <i>Blood</i> , 2009 , 113, 4063-73	2.2	85
78	Oncogenic Flt3 receptors display different specificity and kinetics of autophosphorylation. <i>Experimental Hematology</i> , 2009 , 37, 979-89	3.1	34
77	The c-Kit/D816V mutation eliminates the differences in signal transduction and biological responses between two isoforms of c-Kit. <i>Cellular Signalling</i> , 2009 , 21, 413-8	4.9	22
76	Oncogenic signaling from the hematopoietic growth factor receptors c-Kit and Flt3. <i>Cellular Signalling</i> , 2009 , 21, 1717-26	4.9	95
75	A role of Gab2 association in Flt3 ITD mediated Stat5 phosphorylation and cell survival. <i>British Journal of Haematology</i> , 2009 , 146, 193-202	4.5	26
74	The characterization of epithelial and stromal subsets of candidate stem/progenitor cells in the human adult prostate. <i>European Urology</i> , 2008 , 53, 524-31	10.2	22
73	Stem cell factor induces HIF-1alpha at normoxia in hematopoietic cells. <i>Biochemical and Biophysical Research Communications</i> , 2008 , 377, 98-103	3.4	59
72	Gab2 is involved in differential phosphoinositide 3-kinase signaling by two splice forms of c-Kit. <i>Journal of Biological Chemistry</i> , 2008 , 283, 27444-27451	5.4	42
71	Haematopoietic progenitor cells utilise conventional PKC to suppress PKB/Akt activity in response to c-Kit stimulation. <i>British Journal of Haematology</i> , 2007 , 136, 260-8	4.5	13
70	Grb2 mediates negative regulation of stem cell factor receptor/c-Kit signaling by recruitment of Cbl. <i>Experimental Cell Research</i> , 2007 , 313, 3935-42	4.2	28

69	Identification of Tyrosine Residues of Importance for Survival Signaling through the Scaffolding Protein Gab2 in Both Wild-Type FLT3 and the FLT3-ITD.. <i>Blood</i> , 2007 , 110, 1622-1622	2.2	
68	Interaction and functional cooperation between the serine/threonine kinase bone morphogenetic protein type II receptor with the tyrosine kinase stem cell factor receptor. <i>Journal of Cellular Physiology</i> , 2006 , 206, 457-67	7	21
67	The stem cell factor receptor/c-Kit as a drug target in cancer. <i>Current Cancer Drug Targets</i> , 2006 , 6, 65-75	2.8	69
66	Identification of Y589 and Y599 in the juxtamembrane domain of Flt3 as ligand-induced autophosphorylation sites involved in binding of Src family kinases and the protein tyrosine phosphatase SHP2. <i>Blood</i> , 2006 , 108, 1542-50	2.2	54
65	Direct binding of Cbl to Tyr568 and Tyr936 of the stem cell factor receptor/c-Kit is required for ligand-induced ubiquitination, internalization and degradation. <i>Biochemical Journal</i> , 2006 , 399, 59-67	3.8	65
64	Receptor association and tyrosine phosphorylation of S6 kinases. <i>FEBS Journal</i> , 2006 , 273, 2023-36	5.7	22
63	Co expression of SCF and KIT in gastrointestinal stromal tumours (GISTs) suggests an autocrine/paracrine mechanism. <i>British Journal of Cancer</i> , 2006 , 94, 1180-5	8.7	34
62	Negative Regulation of c-Kit Is Dependent on Direct Binding of Cbl to Tyrosines 568 and 936.. <i>Blood</i> , 2005 , 106, 2288-2288	2.2	
61	Identification of Two Src Recruitment Sites in the Juxtamembrane Region of Flt3 with Opposing Effects on Flt3-Ligand-Induced Signaling.. <i>Blood</i> , 2005 , 106, 2289-2289	2.2	
60	Splice Form Specific Signaling of the Hematopoietic Growth Factor Receptor c-Kit.. <i>Blood</i> , 2005 , 106, 4284-4284	2.2	
59	Platelet-derived growth factor stimulates membrane lipid synthesis through activation of phosphatidylinositol 3-kinase and sterol regulatory element-binding proteins. <i>Journal of Biological Chemistry</i> , 2004 , 279, 35392-402	5.4	93
58	Site-selective regulation of platelet-derived growth factor beta receptor tyrosine phosphorylation by T-cell protein tyrosine phosphatase. <i>Molecular and Cellular Biology</i> , 2004 , 24, 2190-201	4.8	84
57	Gab1 contributes to cytoskeletal reorganization and chemotaxis in response to platelet-derived growth factor. <i>Journal of Biological Chemistry</i> , 2004 , 279, 17897-904	5.4	34
56	p38-MAPK signals survival by phosphorylation of caspase-8 and caspase-3 in human neutrophils. <i>Journal of Experimental Medicine</i> , 2004 , 199, 449-58	16.6	161
55	Signal transduction via the stem cell factor receptor/c-Kit. <i>Cellular and Molecular Life Sciences</i> , 2004 , 61, 2535-48	10.3	322
54	Identification of a Ser/Thr cluster in the C-terminal domain of the human prostaglandin receptor EP4 that is essential for agonist-induced beta-arrestin1 recruitment but differs from the apparent principal phosphorylation site. <i>Biochemical Journal</i> , 2004 , 379, 573-85	3.8	20
53	Src family kinases are involved in the differential signaling from two splice forms of c-Kit. <i>Journal of Biological Chemistry</i> , 2003 , 278, 9159-66	5.4	67
52	Dysfunctionality of a tobacco mosaic virus movement protein mutant mimicking threonine 104 phosphorylation. <i>Journal of General Virology</i> , 2003 , 84, 727-732	4.9	31

51	The adapter protein APS associates with the multifunctional docking sites Tyr-568 and Tyr-936 in c-Kit. <i>Biochemical Journal</i> , 2003 , 370, 1033-8	3.8	45
50	Ligand-induced recruitment of Na ⁺ /H ⁺ -exchanger regulatory factor to the PDGF (platelet-derived growth factor) receptor regulates actin cytoskeleton reorganization by PDGF. <i>Biochemical Journal</i> , 2003 , 376, 505-10	3.8	41
49	Chk1 regulates the S phase checkpoint by coupling the physiological turnover and ionizing radiation-induced accelerated proteolysis of Cdc25A. <i>Cancer Cell</i> , 2003 , 3, 247-58	24.3	460
48	Identification of phosphorylation sites within the SH3 domains of Tec family tyrosine kinases. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2003 , 1645, 123-32	4	29
47	Identification of protein tyrosine phosphatases associating with the PDGF receptor. <i>Biochemistry</i> , 2003 , 42, 2691-9	3.2	44
46	Ezrin is a substrate for Lck in T cells. <i>FEBS Letters</i> , 2003 , 535, 82-6	3.8	33
45	Differential tyrosine phosphorylation of fibroblast growth factor (FGF) receptor-1 and receptor proximal signal transduction in response to FGF-2 and heparin. <i>Experimental Cell Research</i> , 2003 , 287, 190-8	4.2	30
44	Identification of Tyr900 in the kinase domain of c-Kit as a Src-dependent phosphorylation site mediating interaction with c-Crk. <i>Experimental Cell Research</i> , 2003 , 288, 110-8	4.2	37
43	Phosphorylation of the potyvirus capsid protein by protein kinase CK2 and its relevance for virus infection. <i>Plant Cell</i> , 2003 , 15, 2124-39	11.6	97
42	Ser-474 is the major target of insulin-mediated phosphorylation of protein kinase B beta in primary rat adipocytes. <i>Cellular Signalling</i> , 2002 , 14, 175-82	4.9	7
41	SHP-2 is involved in heterodimer specific loss of phosphorylation of Tyr771 in the PDGF beta-receptor. <i>Oncogene</i> , 2002 , 21, 1870-5	9.2	36
40	Different effects of high and low shear stress on platelet-derived growth factor isoform release by endothelial cells: consequences for smooth muscle cell migration. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2002 , 22, 405-11	9.4	54
39	Phosphatidylinositol 3 kinase contributes to the transformation of hematopoietic cells by the D816V c-Kit mutant. <i>Blood</i> , 2001 , 98, 1365-73	2.2	110
38	Mechanisms of platelet-derived growth factor-induced chemotaxis. <i>International Journal of Cancer</i> , 2001 , 91, 757-62	7.5	130
37	Phosphorylation-dependent and -independent functions of p130 cooperate to evoke a sustained G1 block. <i>EMBO Journal</i> , 2001 , 20, 422-32	13	77
36	Activation of Ras, Raf-1 and protein kinase C in differentiating human neuroblastoma cells after treatment with phorbol ester and NGF. <i>Cellular Signalling</i> , 2001 , 13, 95-104	4.9	16
35	DAPP1 undergoes a PI 3-kinase-dependent cycle of plasma-membrane recruitment and endocytosis upon cell stimulation. <i>Current Biology</i> , 2000 , 10, 1403-12	6.3	41
34	TNF-alpha suppresses the PDGF beta-receptor kinase. <i>Experimental Cell Research</i> , 2000 , 258, 65-71	4.2	5

33	Distinct versus redundant properties among members of the INK4 family of cyclin-dependent kinase inhibitors. <i>FEBS Letters</i> , 2000 , 470, 161-6	3.8	61
32	Overactivation of phospholipase C-gamma1 renders platelet-derived growth factor beta-receptor-expressing cells independent of the phosphatidylinositol 3-kinase pathway for chemotaxis. <i>Journal of Biological Chemistry</i> , 1999 , 274, 22089-94	5.4	33
31	Increased mitogenicity of an alphabeta heterodimeric PDGF receptor complex correlates with lack of RasGAP binding. <i>Oncogene</i> , 1999 , 18, 2481-8	9.2	55
30	SHP-2 binds to Tyr763 and Tyr1009 in the PDGF beta-receptor and mediates PDGF-induced activation of the Ras/MAP kinase pathway and chemotaxis. <i>Oncogene</i> , 1999 , 18, 3696-702	9.2	58
29	Phosphorylation of Shc by Src family kinases is necessary for stem cell factor receptor/c-kit mediated activation of the Ras/MAP kinase pathway and c-fos induction. <i>Oncogene</i> , 1999 , 18, 5546-53	9.2	161
28	Characterization of the chronic myelomonocytic leukemia associated TEL-PDGF beta R fusion protein. <i>Oncogene</i> , 1999 , 18, 7055-62	9.2	34
27	Identification of Tyr-703 and Tyr-936 as the primary association sites for Grb2 and Grb7 in the c-Kit/stem cell factor receptor. <i>Biochemical Journal</i> , 1999 , 341, 211-216	3.8	86
26	Functional co-operation between the subunits in heterodimeric platelet-derived growth factor receptor complexes. <i>Biochemical Journal</i> , 1999 , 341, 523-528	3.8	15
25	Identification of Tyr-703 and Tyr-936 as the primary association sites for Grb2 and Grb7 in the c-Kit/stem cell factor receptor. <i>Biochemical Journal</i> , 1999 , 341, 211	3.8	35
24	Functional co-operation between the subunits in heterodimeric platelet-derived growth factor receptor complexes. <i>Biochemical Journal</i> , 1999 , 341, 523	3.8	4
23	Identification of Tyr-762 in the platelet-derived growth factor alpha-receptor as the binding site for Crk proteins. <i>Oncogene</i> , 1998 , 16, 1229-39	9.2	49
22	Signal transduction via platelet-derived growth factor receptors. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 1998 , 1378, F79-113	11.2	296
21	Molecular basis for the dominant white phenotype in the domestic pig. <i>Genome Research</i> , 1998 , 8, 826-33	3.7	142
20	Identification of novel phosphorylation sites in hormone-sensitive lipase that are phosphorylated in response to isoproterenol and govern activation properties in vitro. <i>Journal of Biological Chemistry</i> , 1998 , 273, 215-21	5.4	344
19	Phosphorylation site-specific inhibition of platelet-derived growth factor beta-receptor autophosphorylation by the receptor blocking tyrphostin AG1296. <i>Biochemistry</i> , 1997 , 36, 6260-9	3.2	123
18	Association of coatmer proteins with the beta-receptor for platelet-derived growth factor. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 235, 455-60	3.4	7
17	PDGF-induced phosphorylation of Tyr28 in the N-terminus of Fyn affects Fyn activation. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 241, 355-62	3.4	22
16	Phosphorylation of a 72-kDa protein in PDGF-stimulated cells which forms complex with c-Crk, c-Fyn and Eps15. <i>FEBS Letters</i> , 1997 , 409, 195-200	3.8	8

15	Involvement of phosphatidylinositol 3Kinase in stem-cell-factor-induced phospholipase D activation and arachidonic acid release. <i>FEBS Journal</i> , 1997 , 248, 149-55		35
14	O-5: Identification of the site in the cGMP-inhibited phosphodiesterase phosphorylated in adipocytes in response to insulin and isoproterenol. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 1996 , 104, 10-11	2.3	7
13	Structural determinants in the platelet-derived growth factor alpha-receptor implicated in modulation of chemotaxis. <i>Journal of Biological Chemistry</i> , 1996 , 271, 5101-11	5.4	42
12	Identification of the site in the cGMP-inhibited phosphodiesterase phosphorylated in adipocytes in response to insulin and isoproterenol. <i>Journal of Biological Chemistry</i> , 1996 , 271, 11575-80	5.4	50
11	Identification of the major phosphorylation sites for protein kinase C in kit/stem cell factor receptor in vitro and in intact cells. <i>Journal of Biological Chemistry</i> , 1995 , 270, 14192-200	5.4	72
10	Demonstration of functionally different interactions between phospholipase C-gamma and the two types of platelet-derived growth factor receptors. <i>Journal of Biological Chemistry</i> , 1995 , 270, 7773-81	5.4	45
9	A unique autophosphorylation site in the platelet-derived growth factor alpha receptor from a heterodimeric receptor complex. <i>FEBS Journal</i> , 1994 , 225, 29-41		45
8	Purification of platelet-derived growth factor beta receptor from porcine uterus. <i>Methods in Enzymology</i> , 1991 , 200, 371-8	1.7	2
7	Characterization of the platelet-derived growth factor beta-receptor kinase activity by use of synthetic peptides. <i>Biochemical and Biophysical Research Communications</i> , 1990 , 167, 1333-40	3.4	5
6	Platelet-Derived Growth Factor B Type Receptor 1990 , 303-314		
5	Platelet-derived growth factor receptors in the kidney--upregulated expression in inflammation. <i>Kidney International</i> , 1989 , 36, 1099-102	9.9	109
4	A glioma-derived PDGF A chain homodimer has different functional activities from a PDGF AB heterodimer purified from human platelets. <i>Cell</i> , 1988 , 52, 791-9	56.2	246
3	Purification of human platelet-derived growth factor. <i>Methods in Enzymology</i> , 1987 , 147, 3-13	1.7	43
2	Stimulation of tyrosine phosphorylation by platelet-derived growth factor. <i>Biochemical Society Transactions</i> , 1984 , 12, 759-62	5.1	4
1	Characterization of the fibroblast receptor for platelet-derived growth factor. <i>Cell Biology International Reports</i> , 1983 , 7, 543-544		2