

Michael J Keenan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7795043/publications.pdf>

Version: 2024-02-01

42
papers

1,911
citations

331670

21
h-index

361022

35
g-index

43
all docs

43
docs citations

43
times ranked

2260
citing authors

#	ARTICLE	IF	CITATIONS
1	Resistant starch type 2 and whole grain maize flours enrich different intestinal bacteria and metatranscriptomes. <i>Journal of Functional Foods</i> , 2022, 90, 104982.	3.4	4
2	Gut Microbiota Composition and Predicted Microbial Metabolic Pathways of Obesity Prone and Obesity Resistant Outbred Sprague-Dawley CD Rats May Account for Differences in Their Phenotype. <i>Frontiers in Nutrition</i> , 2021, 8, 746515.	3.7	14
3	Novel Resistant Starch Type 4 Products of Different Starch Origins, Production Methods, and Amounts Are Not Equally Fermented when Fed to Sprague-Dawley Rats. <i>Molecular Nutrition and Food Research</i> , 2020, 64, 1900901.	3.3	8
4	Differences in Capacity of High-Amylose Resistant Starch, Whole-Grain Flour, and a Combination of Both to Modify Intestinal Responses of Male Sprague Dawley Rats Fed Moderate and High Fat Diets. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 15176-15185.	5.2	3
5	Abundance of the species <i>Clostridium butyricum</i> in the gut microbiota contributes to differences in obesity phenotype in outbred Sprague-Dawley CD rats. <i>Nutrition</i> , 2020, 78, 110893.	2.4	15
6	Conserved and variable responses of the gut microbiome to resistant starch type 2. <i>Nutrition Research</i> , 2020, 77, 12-28.	2.9	57
7	CD Obesity-Prone Rats, but not Obesity-Resistant Rats, Robustly Ferment Resistant Starch Without Increased Weight or Fat Accretion. <i>Obesity</i> , 2018, 26, 570-577.	3.0	26
8	Role of resistant starch on diabetes risk factors in people with prediabetes: Design, conduct, and baseline results of the STARCH trial. <i>Contemporary Clinical Trials</i> , 2018, 65, 99-108.	1.8	24
9	Effect of 12 wk of resistant starch supplementation on cardiometabolic risk factors in adults with prediabetes: a randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2018, 108, 492-501.	4.7	54
10	Obese ZDF rats fermented resistant starch with effects on gut microbiota but no reduction in abdominal fat. <i>Molecular Nutrition and Food Research</i> , 2017, 61, 1501025.	3.3	35
11	Misleading conclusions on effects of resistant starch due to inappropriate formulation of controls, inadequate statistical power, and anomalies in the in vitro methods. <i>American Journal of Clinical Nutrition</i> , 2017, 105, 1248-1249.	4.7	0
12	Simultaneous delivery of antibiotics neomycin and ampicillin in drinking water inhibits fermentation of resistant starch in rats. <i>Molecular Nutrition and Food Research</i> , 2017, 61, 1600609.	3.3	12
13	Mice Fed a High-Fat Diet Supplemented with Resistant Starch Display Marked Shifts in the Liver Metabolome Concurrent with Altered Gut Bacteria. <i>Journal of Nutrition</i> , 2016, 146, 2476-2490.	2.9	44
14	Obese Mice Fed a Diet Supplemented with Enzyme-Treated Wheat Bran Display Marked Shifts in the Liver Metabolome Concurrent with Altered Gut Bacteria. <i>Journal of Nutrition</i> , 2016, 146, 2445-2460.	2.9	16
15	The importance of GLP-1 and PYY in resistant starch's effect on body fat in mice. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 1000-1003.	3.3	41
16	Oat consumption reduced intestinal fat deposition and improved health span in <i>Caenorhabditis elegans</i> model. <i>Nutrition Research</i> , 2015, 35, 834-843.	2.9	23
17	Role of Resistant Starch in Improving Gut Health, Adiposity, and Insulin Resistance. <i>Advances in Nutrition</i> , 2015, 6, 198-205.	6.4	194
18	Improving healthspan via changes in gut microbiota and fermentation. <i>Age</i> , 2015, 37, 98.	3.0	33

#	ARTICLE	IF	CITATIONS
19	Prowashonupana barley dietary fibre reduces body fat and increases insulin sensitivity in <i>Caenorhabditis elegans</i> model. <i>Journal of Functional Foods</i> , 2015, 18, 564-574.	3.4	14
20	Induction of Energy Expenditure by Sitagliptin Is Dependent on GLP-1 Receptor. <i>PLoS ONE</i> , 2015, 10, e0126177.	2.5	15
21	Resistant starch from high amylose maize (HAM-RS2) and Dietary butyrate reduce abdominal fat by a different apparent mechanism. <i>Obesity</i> , 2014, 22, 344-348.	3.0	51
22	Resistant starch from high amylose maize (HAM-RS2) reduces body fat and increases gut bacteria in ovariectomized (OVX) rats. <i>Obesity</i> , 2013, 21, 981-984.	3.0	46
23	Dietary resistant starch improves selected brain and behavioral functions in adult and aged rodents. <i>Molecular Nutrition and Food Research</i> , 2013, 57, 2071-2074.	3.3	8
24	High fat diet partially attenuates fermentation responses in rats fed resistant starch from high-amylose maize. <i>Obesity</i> , 2013, 21, 2350-2355.	3.0	49
25	High-Amylose Resistant Starch Increases Hormones and Improves Structure and Function of the Gastrointestinal Tract: A Microarray Study. <i>Journal of Nutrigenetics and Nutrigenomics</i> , 2012, 5, 26-44.	1.3	44
26	Tolerance, fermentation, and cytokine expression in healthy aged male C57BL/6J mice fed resistant starch. <i>Molecular Nutrition and Food Research</i> , 2012, 56, 515-518.	3.3	17
27	Two prebiotics are effective in promoting fermentation in rats fed a high fat diet. <i>FASEB Journal</i> , 2012, 26, 830.9.	0.5	0
28	Dietary-resistant starch improves maternal glycemic control in Goto-Kakizaki rat. <i>Molecular Nutrition and Food Research</i> , 2011, 55, 1499-1508.	3.3	74
29	Dietary Resistant Starch Increases Hypothalamic POMC Expression in Rats. <i>Obesity</i> , 2009, 17, 40-45.	3.0	68
30	Comparative Methodologies for Measuring Metabolizable Energy of Various Types of Resistant High Amylose Corn Starch. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 8474-8479.	5.2	22
31	Failure to Ferment Dietary Resistant Starch in Specific Mouse Models of Obesity Results in No Body Fat Loss. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 8844-8851.	5.2	47
32	Molecular Characterization of Gut Microflora of Mice Fed Dietary Resistant Starch. <i>FASEB Journal</i> , 2009, 23, 919.7.	0.5	1
33	Dietary resistant starch upregulates total GLP-1 and PYY in a sustained day-long manner through fermentation in rodents. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2008, 295, E1160-E1166.	3.5	359
34	Feeding resistant starch maintains elevated plasma levels of GLP-1 and PYY throughout the day and is associated with decreased body fat in rats. <i>FASEB Journal</i> , 2007, 21, A158.	0.5	1
35	Feeding resistant starch to rats alters expression of the cecal cell genome compared to control groups. <i>FASEB Journal</i> , 2007, 21, A364.	0.5	0
36	Resistant starch in a high fat diet produces signaling from the gut, but not reduced body fat. <i>FASEB Journal</i> , 2007, 21, A364.	0.5	0

#	ARTICLE	IF	CITATIONS
37	Effects of Resistant Starch, A Non-digestible Fermentable Fiber, on Reducing Body Fat. Obesity, 2006, 14, 1523-1534.	3.0	255
38	Peptide YY and Proglucagon mRNA Expression Patterns and Regulation in the Gut. Obesity, 2006, 14, 683-689.	3.0	154
39	Resistant starch and fructooligosaccharide improve gut histology and alter gut signaling in rats. FASEB Journal, 2006, 20, A182.	0.5	0
40	Resistant starch reduces abdominal fat more than energy dilution with nonfermentable fiber. FASEB Journal, 2006, 20, A182.	0.5	0
41	Adult Female Rats Defend "Appropriate" Energy Intake after Adaptation to Dietary Energy. Obesity, 2003, 11, 1214-1222.	4.0	14
42	Comparison of Bone Density Measurement Techniques: DXA and Archimedes' Principle. Journal of Bone and Mineral Research, 1997, 12, 1903-1907.	2.8	69