
## Yoshitaka Kaneita

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7797459/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Awareness, Attitudes, and Concerns Regarding Heated Tobacco Products Among Physicians in Japan. Journal of Epidemiology, 2023, 33, 441-449.	2.4	4
2	A cross-sectional epidemiological study of the relationship between sleep duration, quality, and rhythm and presenteeism in workers. Sleep and Biological Rhythms, 2022, 20, 53-63.	1.0	7
3	Association between nighttime urinary frequency and sleep problems among Japanese adolescents. International Journal of Urology, 2022, 29, 152-157.	1.0	2
4	Associations Between the 2011 Great East Japan Earthquake and Tsunami and the Sleep and Mental Health of Japanese People: A 3-Wave Repeated Survey. Nature and Science of Sleep, 2022, Volume 14, 61-73.	2.7	6
5	The validity and reliability of the Japanese version of RU-SATED. Sleep Medicine, 2022, 91, 109-114.	1.6	6
6	Associations between Coping Profile and Work Performance in a Cohort of Japanese Employees. International Journal of Environmental Research and Public Health, 2022, 19, 4806.	2.6	3
7	Psychometric Properties of a New Simplified Scale to Assess the Quality of Workers' Rest and Recreation on Their Days Off. Yonago Acta Medica, 2021, 64, 67-79.	0.7	0
8	Trends in sleep problems and patterns among Japanese adolescents: 2004 to 2017. The Lancet Regional Health - Western Pacific, 2021, 9, 100107.	2.9	12
9	Clarifying the factors affecting the implementation of the "early to bed, early to rise, and don't forget your breakfast―campaign aimed at adolescents in Japan. Sleep and Biological Rhythms, 2021, 19, 325-336.	1.0	3
10	A nationwide cross-sectional study of difficulty waking up for school among adolescents. Sleep, 2021, 44, .	1.1	1
11	The association between Internet usage and sleep problems among Japanese adolescents: three repeated cross-sectional studies. Sleep, 2021, 44, .	1.1	16
12	A Nationwide Cohort Study of Parasomnias Among Adolescents. Journal of Clinical Psychiatry, 2021, 82, .	2.2	1
13	Trends in the co-use of alcohol and tobacco among Japanese adolescents: periodical nationwide cross-sectional surveys 1996–2017. BMJ Open, 2021, 11, e045063.	1.9	8
14	Associations between coping strategies and insomnia: a longitudinal study of Japanese workers. Sleep, 2021, , .	1.1	3
15	Effectiveness of Screening and Brief Alcohol Intervention at the Workplace: A Study Protocol for a Randomized Controlled Trial at Five Japan-Based Companies. Yonago Acta Medica, 2021, 64, 330-338.	0.7	1
16	A school-based sleep hygiene education program for adolescents in Japan: a large-scale comparative intervention study. Sleep and Biological Rhythms, 2020, 18, 27-36.	1.0	10
17	Association between reduced serum BDNF levels and insomnia with short sleep duration among female hospital nurses. Sleep Medicine, 2020, 68, 167-172.	1.6	26
18	A composite measure of sleep health is associated with symptoms of depression among Japanese female hospital nurses. Comprehensive Psychiatry, 2020, 97, 152151.	3.1	27

ΥΟSΗΙΤΑΚΑ ΚΑΝΕΙΤΑ

#	Article	IF	CITATIONS
19	Association between Typus melancholicus and depressive symptoms: A Japanese general population survey. Psychiatry and Clinical Neurosciences, 2020, 74, 672-673.	1.8	2
20	Skipping breakfast, poor sleep quality, and Internet usage and their relation with unhappiness in Japanese adolescents. PLoS ONE, 2020, 15, e0235252.	2.5	6
21	Relationship between internet addiction and sleep disturbance in high school students: a cross-sectional study. BMC Pediatrics, 2020, 20, 379.	1.7	17
22	Gender differences in dietary behaviors among Japanese adolescents. Preventive Medicine Reports, 2020, 20, 101203.	1.8	8
23	The relationship between subjective happiness and sleep problems in Japanese adolescents. Sleep Medicine, 2020, 69, 120-126.	1.6	27
24	Factors Associated with the Development of Childhood Asthma in Japan: A Nationwide Longitudinal Study. Maternal and Child Health Journal, 2020, 24, 911-922.	1.5	8
25	Heat-not-burn tobacco, electronic cigarettes, and combustible cigarette use among Japanese adolescents: a nationwide population survey 2017. BMC Public Health, 2020, 20, 741.	2.9	18
26	Stress-Coping Strategies and Factors Related Distress among Japanese Physicians. Iranian Journal of Public Health, 2020, 49, 1387-1389.	0.5	0
27	Relationship between Internet Addiction and Poor Mental Health among Japanese Adolescents. Iranian Journal of Public Health, 2020, 49, 2069-2077.	0.5	1
28	Sleep disturbance is associated with not only shorter sleep duration, but also longer time in bed: a Japanese general population survey. Sleep and Biological Rhythms, 2019, 17, 407-415.	1.0	6
29	Association between unhealthy dietary behaviors and sleep disturbances among Japanese adolescents: a nationwide representative survey. Sleep and Biological Rhythms, 2019, 17, 93-102.	1.0	27
30	Factors Related to Alcohol Consumption Among Japanese Physicians. Asia-Pacific Journal of Public Health, 2018, 30, 296-306.	1.0	8
31	Sleep-related factors associated with industrial accidents among factory workers and sleep hygiene education intervention. Sleep and Biological Rhythms, 2018, 16, 239-251.	1.0	13
32	Unhealthy lifestyle factors and depressive symptoms: A Japanese general adult population survey. Journal of Affective Disorders, 2018, 234, 156-161.	4.1	26
33	Long sleep duration and health outcomes: A systematic review, meta-analysis and meta-regression. Sleep Medicine Reviews, 2018, 39, 25-36.	8.5	464
34	The association between sleep disturbance and second-hand smoke exposure: a large-scale, nationwide, cross-sectional study of adolescents in Japan. Sleep Medicine, 2018, 50, 29-35.	1.6	29
35	Stressful events and coping strategies associated with symptoms of depression: A Japanese general population survey. Journal of Affective Disorders, 2018, 238, 482-488.	4.1	33
36	Longitudinal Epidemiologic Study of Poor Mental Health Status in Japanese Adolescents. Journal of Clinical Psychiatry, 2018, 79, .	2.2	4

ΥΟSΗΙΤΑΚΑ ΚΑΝΕΙΤΑ

#	Article	IF	CITATIONS
37	The association between alcohol use and problematic internet use: A large-scale nationwide cross-sectional study of adolescents in Japan. Journal of Epidemiology, 2017, 27, 107-111.	2.4	48
38	Short sleep duration, shift work, and actual days taken off work are predictive life-style risk factors for new-onset metabolic syndrome: a seven-year cohort study of 40,000 male workers. Sleep Medicine, 2017, 39, 87-94.	1.6	60
39	Relationship between stress coping and sleep disorders among the general Japanese population: a nationwide representative survey. Sleep Medicine, 2017, 37, 38-45.	1.6	43
40	Short sleep duration and health outcomes: a systematic review, meta-analysis, and meta-regression. Sleep Medicine, 2017, 32, 246-256.	1.6	710
41	Nightmares and sleep paralysis among the general Japanese population: a nationwide representative survey. Sleep and Biological Rhythms, 2017, 16, 187.	1.0	1
42	An Aggregate Measure of Sleep Health Is Associated With Prevalent and Incident Clinically Significant Depression Symptoms Among Community-Dwelling Older Women. Sleep, 2017, 40, .	1.1	78
43	Predictors of insomnia onset in adolescents in Japan. Sleep Medicine, 2017, 38, 37-43.	1.6	16
44	An Epidemiological Study on Psychosocial Factors Associated with Depression among Japanese General Adult Population. Journal of the Nihon University Medical Association, 2016, 75, 81-87.	0.0	0
45	The Association Between Alcohol-Flavoured Non-Alcoholic Beverages and Alcohol Use in Japanese Adolescents. Alcohol and Alcoholism, 2016, 52, 351-357.	1.6	3
46	The association between shift work and health: a review. Sleep and Biological Rhythms, 2016, 14, 231-239.	1.0	13
47	Nationwide epidemiological study of insomnia in Japan. Sleep Medicine, 2016, 25, 130-138.	1.6	57
48	Association Between Smoking and Problematic Internet Use Among Japanese Adolescents: Large-Scale Nationwide Epidemiological Study. Cyberpsychology, Behavior, and Social Networking, 2016, 19, 557-561.	3.9	25
49	Factors influencing adherence to nasal continuous positive airway pressure in obstructive sleep apnea patients in Japan. Sleep and Biological Rhythms, 2016, 14, 339-349.	1.0	3
50	Internet use and problematic Internet use among adolescents in Japan: A nationwide representative survey. Addictive Behaviors Reports, 2016, 4, 58-64.	1.9	70
51	Efficacy of sleep education program based on principles of cognitive behavioral therapy to alleviate workers' distress. Sleep and Biological Rhythms, 2016, 14, 211-219.	1.0	10
52	Napping and associated factors: a Japanese nationwide general population survey. Sleep Medicine, 2016, 20, 72-79.	1.6	38
53	Anger and Impulsivity Among Japanese Adolescents. Journal of Clinical Psychiatry, 2016, 77, e860-e866.	2.2	11
54	Irritable bowel syndrome among <scp>J</scp> apanese adolescents: A nationally representative survey. Journal of Gastroenterology and Hepatology (Australia), 2015, 30, 1354-1360.	2.8	19

Υοςηιτακά Κανειτά

#	Article	IF	CITATIONS
55	Association of short sleep duration and short time in bed with depression: A Japanese general population survey. Sleep and Biological Rhythms, 2015, 13, 136-145.	1.0	38
56	Prevalence of fatigue symptoms and correlations in the general adult population. Sleep and Biological Rhythms, 2015, 13, 146-154.	1.0	17
57	Epidemiological study of the associations between sleep complaints and metabolic syndrome in Japan. Sleep and Biological Rhythms, 2014, 12, 269-278.	1.0	27
58	Excessive daytime sleepiness among pregnant women: An epidemiological study. Sleep and Biological Rhythms, 2014, 12, 12-21.	1.0	4
59	Sleep and Health Promotion. Sleep and Biological Rhythms, 2014, 12, 73-73.	1.0	2
60	Sleep environments of Japanese adolescents: An epidemiological study. Sleep and Biological Rhythms, 2014, 12, 116-126.	1.0	1
61	Factors Affecting Unhappiness at School among Japanese Adolescents: An Epidemiological Study. PLoS ONE, 2014, 9, e111844.	2.5	15
62	Disorders of arousal and sleep-related bruxism among Japanese adolescents: a nationwide representative survey. Sleep Medicine, 2013, 14, 532-541.	1.6	31
63	Associations between sleep disturbance and alcohol drinking: A large-scale epidemiological study of adolescents in Japan. Alcohol, 2013, 47, 619-628.	1.7	29
64	Associations of Work Hours and Actual Availability of Weekly Rest Days with Cardiovascular Risk Factors. Journal of Occupational Health, 2013, 55, 11-20.	2.1	19
65	Prescription hypnotics and associated background factors in a large-scale Japanese database. Sleep and Biological Rhythms, 2012, 10, 319-327.	1.0	5
66	The association between sleep problems and perceived health status: A Japanese nationwide general population survey. Sleep Medicine, 2012, 13, 831-837.	1.6	72
67	Clarification of the factor structure of the 12-item General Health Questionnaire among Japanese adolescents and associated sleep status. Psychiatry Research, 2011, 188, 138-146.	3.3	38
68	Nightmare and sleep paralysis among Japanese adolescents: A nationwide representative survey. Sleep Medicine, 2011, 12, 56-64.	1.6	82
69	Association of onset of obesity with sleep duration and shift work among Japanese adults. Sleep Medicine, 2011, 12, 341-345.	1.6	82
70	The Association between Use of Mobile Phones after Lights Out and Sleep Disturbances among Japanese Adolescents: A Nationwide Cross-Sectional Survey. Sleep, 2011, 34, 1013-1020.	1.1	254
71	Stress coping behaviors and sleep hygiene practices in a sample of Japanese adults with insomnia. Sleep and Biological Rhythms, 2011, 9, 35-45.	1.0	11
72	New onset and natural remission of excessive daytime sleepiness and its correlates among high-school students. Sleep and Biological Rhythms, 2011, 9, 117-126.	1.0	1

Υοςηιτακά Κανειτά

#	Article	IF	CITATIONS
73	Association of Current Work and Sleep Situations with Excessive Daytime Sleepiness and Medical Incidents among Japanese Physicians. Journal of Clinical Sleep Medicine, 2011, 07, 512-522.	2.6	23
74	Insomnia symptoms associated with hyperglycemia. Sleep and Biological Rhythms, 2010, 8, 203-211.	1.0	5
75	Excessive daytime sleepiness and sleep behavior among Japanese adolescents: A nation-wide representative survey. Sleep and Biological Rhythms, 2010, 8, 282-294.	1.0	10
76	Association between Depression and Insomnia Subtypes: A Longitudinal Study on the Elderly in Japan. Sleep, 2010, 33, 1693-1702.	1.1	149
77	Epidemiological study of smoking among Japanese physicians. Preventive Medicine, 2010, 51, 164-167.	3.4	11
78	Decrease in the Prevalence of Adolescent Alcohol Use and its Possible Causes in Japan: Periodical Nationwide Crossâ€Sectional Surveys. Alcoholism: Clinical and Experimental Research, 2009, 33, 247-254.	2.4	31
79	Epidemiological study of nightmare and sleep paralysis among Japanese adolescents. Sleep and Biological Rhythms, 2009, 7, 201-210.	1.0	17
80	Coping strategies and their correlates with depression in the Japanese general population. Psychiatry Research, 2009, 168, 57-66.	3.3	63
81	Associations between sleep disturbance and mental health status: A longitudinal study of Japanese junior high school students. Sleep Medicine, 2009, 10, 780-786.	1.6	118
82	Decrease in the prevalence of smoking among Japanese adolescents and its possible causes: periodic nationwide cross-sectional surveys. Environmental Health and Preventive Medicine, 2008, 13, 219-226.	3.4	21
83	Prevalence of restless legs syndrome with pregnancy and the relationship with sleep disorders in the Japanese large population. Sleep and Biological Rhythms, 2008, 6, 102-109.	1.0	22
84	Associations of Usual Sleep Duration with Serum Lipid and Lipoprotein Levels. Sleep, 2008, 31, 645-652.	1.1	198
85	Is Passive Smoking Associated With Sleep Disturbance Among Pregnant Women?. Sleep, 2007, , .	1.1	Ο
86	Use of alcohol and hypnotic medication as aids to sleep among the Japanese general population. Sleep Medicine, 2007, 8, 723-732.	1.6	69
87	Prevalence of smoking and associated factors among pregnant women in Japan. Preventive Medicine, 2007, 45, 15-20.	3.4	39
88	Association Between Mental Health Status and Sleep Status Among Adolescents in Japan. Journal of Clinical Psychiatry, 2007, 68, 1426-1435.	2.2	118
89	Epidemiological study on passive smoking among Japanese infants and smoking behavior of their respective parents: A nationwide cross-sectional survey. Preventive Medicine, 2006, 42, 210-217.	3.4	19
90	Insomnia Among Japanese Adolescents: A Nationwide Representative Survey. Sleep, 2006, 29, 1543-1550.	1.1	114

Yoshitaka Kaneita

#	Article	IF	CITATIONS
91	Epidemiological study of the relationship between sleep disturbances and somatic and psychological complaints among the Japanese general population. Sleep and Biological Rhythms, 2006, 4, 55-62.	1.0	22
92	The Relationship Between Depression and Sleep Disturbances. Journal of Clinical Psychiatry, 2006, 67, 196-203.	2.2	184
93	Excessive Daytime Sleepiness among the Japanese General Population. Journal of Epidemiology, 2005, 15, 1-8.	2.4	118
94	Relation of smoking and drinking to sleep disturbance among Japanese pregnant women. Preventive Medicine, 2005, 41, 877-882.	3.4	35
95	An Epidemiologic Study of Self-Reported Sleep Problems among Japanese Adolescents. Sleep, 2004, 27, 978-985.	1.1	139
96	Hepatitis C virus is frequently coinfected with serum marker-negative hepatitis B virus: Probable replication promotion of the former by the latter as demonstrated by in vitro cotransfection. Journal of Medical Virology, 1997, 52, 399-405.	5.0	63
97	Longitudinal epidemiological study of subjective sleep quality in Japanese adolescents to investigate predictors of poor sleep quality. Sleep and Biological Rhythms, 0, , 1.	1.0	3
98	The Japanese Youth Version of the Smartphone Addiction Scale Among the Youth in Japan: Reliability and Validity Assessment. International Journal of Mental Health and Addiction, 0, , 1.	7.4	3
99	Association of insomnia symptoms and non-restorative sleep with Typus melancholicus: a Japanese general population survey. Sleep and Biological Rhythms, 0, , 1.	1.0	0