

Firas Abdollah

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7802686/firas-abdollah-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

362
papers

9,367
citations

54
h-index

79
g-index

537
ext. papers

11,131
ext. citations

3.8
avg, IF

5.7
L-index

| # | Paper | IF | Citations |
|-----|--|-----|-----------|
| 362 | Robotic vs laparoscopic nephroureterectomy for upper tract urothelial carcinoma: a multicenter propensity-score matched pair "tetrafecta" analysis (ROBUUST collaborative group).. <i>Journal of Endourology</i> , 2022 , | 2.7 | 2 |
| 361 | Outcomes of Lymph Node Dissection in Nephroureterectomy in the Treatment of Upper Tract Urothelial Carcinoma: Analysis of the ROBUUST Registry.. <i>Journal of Urology</i> , 2022 , 101097JU00000000000002690 | 2.5 | 1 |
| 360 | Regression Discontinuity Analysis of Salvage Radiotherapy in Prostate Cancer. <i>European Urology Oncology</i> , 2021 , 4, 817-820 | 6.7 | 1 |
| 359 | Single stage Xi robotic radical nephroureterectomy for upper tract urothelial carcinoma: surgical technique and outcomes. <i>Minerva Urology and Nephrology</i> , 2021 , | 2.3 | 6 |
| 358 | Improving the stratification of intermediate risk prostate cancer. <i>Minerva Urology and Nephrology</i> , 2021 , | 2.3 | 3 |
| 357 | Editorial Comment. <i>Journal of Urology</i> , 2021 , 101097JU0000000000000225002 | 2.5 | |
| 356 | COVID-19 Infection in Men on Testosterone Replacement Therapy. <i>Journal of Sexual Medicine</i> , 2021 , 18, 215-218 | 1.1 | 13 |
| 355 | Impact of treatment modality on overall survival in localized ductal prostate adenocarcinoma: A national cancer database analysis. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021 , 39, 366.e11-366.e18 | 2.8 | 1 |
| 354 | Generalizability of Prostate-Specific Antigen (PSA) Screening Trials in a "Real World" Setting: A Nationwide Survey Analysis. <i>Urology</i> , 2021 , 148, 1-3 | 1.6 | 1 |
| 353 | Robotic Kidney Transplantation with Regional Hypothermia versus Open Kidney Transplantation for Patients with End Stage Renal Disease: An Ideal Stage 2B Study. <i>Journal of Urology</i> , 2021 , 205, 595-602 | 2.5 | 10 |
| 352 | A Preoperative Nomogram to Predict Renal Function Insufficiency for Cisplatin-based Adjuvant Chemotherapy Following Minimally Invasive Radical Nephroureterectomy (ROBUUST Collaborative Group). <i>European Urology Focus</i> , 2021 , | 5.1 | 2 |
| 351 | Subtotal surgical therapy for localized prostate cancer: a single-center precision prostatectomy experience in 25 patients, and SEER-registry data analysis. <i>Translational Andrology and Urology</i> , 2021 , 10, 3155-3166 | 2.3 | 1 |
| 350 | Author Reply. <i>Urology</i> , 2021 , 153, 361-362 | 1.6 | |
| 349 | Evaluation of lymphovascular invasion as a prognostic predictor of overall survival after radical prostatectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021 , 39, 495.e1-495.e6 | 2.8 | 1 |
| 348 | High-intensity local treatment of clinical node-positive urothelial carcinoma of the bladder alongside systemic chemotherapy improves overall survival. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021 , 40, 62.e1-62.e1 | 2.8 | |
| 347 | Risk Factors for Intravesical Recurrence after Minimally Invasive Nephroureterectomy for Upper Tract Urothelial Cancer (ROBUUST Collaboration). <i>Journal of Urology</i> , 2021 , 206, 568-576 | 2.5 | 6 |
| 346 | Impact of Lymphovascular Invasion on Overall Survival in Patients With Prostate Cancer Following Radical Prostatectomy: Stage-per-Stage Analysis. <i>Clinical Genitourinary Cancer</i> , 2021 , 19, e319-e325 | 3.3 | 0 |

| | | | |
|-----|--|------|----|
| 345 | External validation of genomic classifier-based risk-stratification tool to identify candidates for adjuvant radiation therapy in patients with prostate cancer. <i>World Journal of Urology</i> , 2021 , 39, 3217-3222 | 4.2 | 0 |
| 344 | AUTHOR REPLY.. <i>Urology</i> , 2021 , 158, 115-116 | 1.6 | |
| 343 | The Precision Prostatectomy: "Waiting for Godot". <i>European Urology Focus</i> , 2020 , 6, 227-230 | 5.1 | 4 |
| 342 | Ten-year disease progression and mortality rates in men who experience biochemical recurrence versus persistence after radical prostatectomy and undergo salvage radiation therapy: A post-hoc analysis of RTOG 9601 trial data. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020 , 38, 599.e1-599.e8 | 2.8 | 6 |
| 341 | 462. Prevalence and Outcome of Asymptomatic Procedural Patients with COVID-19 Infection. <i>Open Forum Infectious Diseases</i> , 2020 , 7, S298-S298 | 1 | 2 |
| 340 | Rare Histological Variants of Prostate Adenocarcinoma: A National Cancer Database Analysis. <i>Journal of Urology</i> , 2020 , 204, 260-266 | 2.5 | 6 |
| 339 | Long-term Risk of Recurrence in Surgically Treated Renal Cell Carcinoma: A Post Hoc Analysis of the Eastern Cooperative Oncology Group-American College of Radiology Imaging Network E2805 Trial Cohort. <i>European Urology</i> , 2020 , 77, 277-281 | 10.2 | 5 |
| 338 | Impact of timing on salvage radiation therapy adverse events following radical prostatectomy: A secondary analysis of the RTOG 9601 cohort. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020 , 38, 38.e17-38.e22 | 2.8 | 3 |
| 337 | Pelvic lymph node dissection at robot-assisted radical prostatectomy: Assessing utilization and nodal metastases within a statewide quality improvement consortium. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020 , 38, 198-203 | 2.8 | 2 |
| 336 | Extended pelvic lymph-node dissection is independently associated with improved overall survival in patients with prostate cancer at high-risk of lymph-node invasion. <i>BJU International</i> , 2020 , 125, 756-758 | 5.6 | 5 |
| 335 | A Nationwide Persistent Underutilization of Adjuvant Radiotherapy in North American Prostate Cancer Patients. <i>Clinical Genitourinary Cancer</i> , 2020 , 18, 489-499.e6 | 3.3 | 2 |
| 334 | Re: Timothy J. Wilt, Tien N. Vo, Lisa Langsetmo, et al. Radical Prostatectomy or Observation for Clinically Localized Prostate Cancer: Extended Follow-up of the Prostate Cancer Intervention Versus Observation Trial (PIVOT). <i>Eur Urol</i> 2020;77:713-724: External Validity of the Updated Prostate Cancer Intervention Versus Observation Trial (PIVOT). <i>European Urology Oncology</i> , 2020 , 3, 10-13 | 6.7 | 0 |
| 333 | Omission of Cortical Renorrhaphy During Robotic Partial Nephrectomy: A Vattikuti Collective Quality Initiative Database Analysis. <i>Urology</i> , 2020 , 146, 125-132 | 1.6 | 1 |
| 332 | Nephroureterectomy with or without Bladder Cuff Excision for Localized Urothelial Carcinoma of the Renal Pelvis. <i>European Urology Focus</i> , 2020 , 6, 298-304 | 5.1 | 10 |
| 331 | Variation in Positive Surgical Margin Status After Radical Prostatectomy for pT2 Prostate Cancer. <i>Clinical Genitourinary Cancer</i> , 2019 , 17, e1060-e1068 | 3.3 | 6 |
| 330 | Facility Level Variation in Rates of Definitive Therapy for Low Risk Prostate Cancer in Men with Limited Life Expectancy: An Opportunity for Value Based Care Redesign. <i>Journal of Urology</i> , 2019 , 201, 728-734 | 2.5 | 4 |
| 329 | Minimally Invasive or Abdominal Radical Hysterectomy for Cervical Cancer. <i>New England Journal of Medicine</i> , 2019 , 380, 793 | 59.2 | 7 |
| 328 | A Systematic Review of the Role of Definitive Local Treatment in Patients with Clinically Lymph Node-positive Prostate Cancer. <i>European Urology Oncology</i> , 2019 , 2, 294-301 | 6.7 | 13 |

| | | | |
|-----|--|------|----|
| 327 | Cytoreductive Nephrectomy: Assessing the Generalizability of the CARMENA Trial to Real-world National Cancer Data Base Cases. <i>European Urology</i> , 2019 , 75, 352-353 | 10.2 | 22 |
| 326 | Increasing Rate of Noninterventional Treatment Management in Localized Prostate Cancer Candidates for Active Surveillance: A North American Population-Based Study. <i>Clinical Genitourinary Cancer</i> , 2019 , 17, 72-78.e4 | 3.3 | 5 |
| 325 | Contemporary Trends and Survival Outcomes After Aborted Radical Prostatectomy in Lymph Node Metastatic Prostate Cancer Patients. <i>European Urology Focus</i> , 2019 , 5, 381-388 | 5.1 | 9 |
| 324 | The Impact of Lymph Node Metastases Burden at Radical Prostatectomy. <i>European Urology Focus</i> , 2019 , 5, 399-406 | 5.1 | 13 |
| 323 | Comparison of Partial Versus Radical Nephrectomy Effect on Other-cause Mortality, Cancer-specific Mortality, and 30-day Mortality in Patients Older Than 75 Years. <i>European Urology Focus</i> , 2019 , 5, 467-473 | 5.1 | 10 |
| 322 | Contemporary Trends in the Incidence of Metastatic Prostate Cancer Among US Men: Results from Nationwide Analyses. <i>European Urology Focus</i> , 2019 , 5, 77-80 | 5.1 | 32 |
| 321 | The effect of race on survival after local therapy in metastatic prostate cancer patients. <i>Canadian Urological Association Journal</i> , 2019 , 13, 175-181 | 1.2 | 1 |
| 320 | A contemporary analysis of radiotherapy effect in surgically treated retroperitoneal sarcoma. <i>Radiotherapy and Oncology</i> , 2018 , 127, 318-325 | 5.3 | 11 |
| 319 | Comparative effectiveness of robot-assisted vs. open radical cystectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 88.e1-88.e9 | 2.8 | 35 |
| 318 | Functional Recovery, Oncologic Outcomes and Postoperative Complications after Robot-Assisted Radical Prostatectomy: An Evidence-Based Analysis Comparing the Retzius Sparing and Standard Approaches. <i>Journal of Urology</i> , 2018 , 199, 1210-1217 | 2.5 | 73 |
| 317 | Reply. <i>Urology</i> , 2018 , 112, 64-65 | 1.6 | |
| 316 | Re: Stenting prior to Cystectomy is an Independent Risk Factor for Upper Urinary Tract Recurrence: B. Kiss, M. A. Furrer, P. Y. Wuethrich, F. C. Burkhard, G. N. Thalmann and B. Roth <i>J Urol</i> 2017;198:1263-1268. <i>Journal of Urology</i> , 2018 , 199, 1069-1070 | 2.5 | |
| 315 | Impact of Adjuvant Radiotherapy in Node-positive Prostate Cancer Patients: The Importance of Patient Selection. <i>European Urology</i> , 2018 , 74, 253-256 | 10.2 | 32 |
| 314 | Reply by the Authors. <i>Urology</i> , 2018 , 116, 232-233 | 1.6 | |
| 313 | Contemporary Management of Prostate Cancer Patients Suitable for Active Surveillance: A North American Population-based Study. <i>European Urology Focus</i> , 2018 , 4, 68-74 | 5.1 | 12 |
| 312 | Impact of Baseline Characteristics on the Survival Benefit of High-Intensity Local Treatment in Metastatic Urothelial Carcinoma of the Bladder. <i>European Urology Focus</i> , 2018 , 4, 568-571 | 5.1 | 5 |
| 311 | Trends in Prostate-Specific Antigen Screening Since the Implementation of the 2012 US Preventive Services Task Force Recommendations. <i>European Urology Focus</i> , 2018 , 4, 1002-1004 | 5.1 | 2 |
| 310 | Rate and Extent of Pelvic Lymph Node Dissection in the US Prostate Cancer Patients Treated With Radical Prostatectomy. <i>Clinical Genitourinary Cancer</i> , 2018 , 16, e451-e467 | 3.3 | 11 |

| | | | |
|-----|--|------|-----|
| 309 | Testing the external validity of the EORTC randomized trial 30904 comparing overall survival after radical nephrectomy vs nephron-sparing surgery in contemporary North American patients with renal cell cancer. <i>BJU International</i> , 2018 , 121, 345-347 | 5.6 | 7 |
| 308 | Efficacy of Local Treatment in Prostate Cancer Patients with Clinically Pelvic Lymph Node-positive Disease at Initial Diagnosis. <i>European Urology</i> , 2018 , 73, 452-461 | 10.2 | 29 |
| 307 | Variation in the use of active surveillance for low-risk prostate cancer. <i>Cancer</i> , 2018 , 124, 55-64 | 6.4 | 26 |
| 306 | The Association between Mortality and Distance to Treatment Facility in Patients with Muscle Invasive Bladder Cancer. <i>Journal of Urology</i> , 2018 , 199, 424-429 | 2.5 | 22 |
| 305 | Re: Massimiliano Spaliviero, Nicholas E. Power, Katie S. Murray, et al. Intravenous Mannitol Versus Placebo During Partial Nephrectomy in Patients with Normal Kidney Function: A Double-blind, Clinically-integrated, Randomized Trial. <i>Eur Urol</i> 2018;73:53-9. <i>European Urology</i> , 2018 , 74, e48-e49 | 10.2 | 1 |
| 304 | Racial disparity in quality of care and overall survival among black vs. white patients with muscle-invasive bladder cancer treated with radical cystectomy: A national cancer database analysis. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 469.e1-469.e11 | 2.8 | 16 |
| 303 | Effectiveness of Neoadjuvant Chemotherapy for Muscle-invasive Bladder Cancer in the Current Real World Setting in the USA. <i>European Urology Oncology</i> , 2018 , 1, 83-90 | 6.7 | 34 |
| 302 | Testing the impact of adjuvant radiotherapy (aRT) after radical prostatectomy (RP) on overall mortality (OM) in prostate cancer patients with pathologically node positive disease: A nationwide analysis.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 5035-5035 | 2.2 | |
| 301 | Re: Follow-up of Prostatectomy Versus Observation for Early Prostate Cancer. <i>European Urology</i> , 2018 , 73, 302-303 | 10.2 | 2 |
| 300 | Performance of a Prostate Cancer Genomic Classifier in Predicting Metastasis in Men with Prostate-specific Antigen Persistence Postprostatectomy. <i>European Urology</i> , 2018 , 74, 107-114 | 10.2 | 36 |
| 299 | State-by-state Variation in Prostate-specific Antigen Screening Trends Following the 2011 United States Preventive Services Task Force Panel Update. <i>Urology</i> , 2018 , 112, 56-65 | 1.6 | 4 |
| 298 | First North American validation and head-to-head comparison of four preoperative nomograms for prediction of lymph node invasion before radical prostatectomy. <i>BJU International</i> , 2018 , 121, 592-599 | 5.6 | 24 |
| 297 | Prostate Cancer Screening in Early Medicaid Expansion States. <i>Journal of Urology</i> , 2018 , 199, 81-88 | 2.5 | 17 |
| 296 | Racial Disparity in Delivering Definitive Therapy for Intermediate/High-risk Localized Prostate Cancer: The Impact of Facility Features and Socioeconomic Characteristics. <i>European Urology</i> , 2018 , 73, 445-451 | 10.2 | 27 |
| 295 | Development and Validation of a Novel Integrated Clinical-Genomic Risk Group Classification for Localized Prostate Cancer. <i>Journal of Clinical Oncology</i> , 2018 , 36, 581-590 | 2.2 | 107 |
| 294 | Comparing Adjuvant vs Early-Salvage Radiotherapy After Radical Prostatectomy. <i>JAMA Oncology</i> , 2018 , 4, 1618-1619 | 13.4 | 1 |
| 293 | Oncological and functional efficacy of nephron-sparing surgery versus radical nephrectomy in renal cell carcinoma stages T1b: a single institution, matched analysis. <i>Central European Journal of Urology</i> , 2018 , 71, 48-57 | 0.9 | 1 |
| 292 | Development of the Vattikuti Institute Prostatectomy: Historical Perspective and Technical Nuances 2018 , 255-273 | | |

| | | | |
|-----|--|------|----|
| 291 | Functional and Oncological Outcomes of Robotic Radical Prostatectomy 2018 , 409-425 | | |
| 290 | Robotic Urologic Surgery: How to Make an Effective Robotic Program 2018 , 77-82 | | 1 |
| 289 | Robot-Assisted Laparoscopic Radical Prostatectomy in Patients with Clinically High-Risk Prostate Cancer 2018 , 363-373 | | |
| 288 | Development and Validation of a Prostate Cancer Genomic Signature that Predicts Early ADT Treatment Response Following Radical Prostatectomy. <i>Clinical Cancer Research</i> , 2018 , 24, 3908-3916 | 12.9 | 10 |
| 287 | Surgically Treated Retroperitoneal Sarcoma: A Population-based Competing Risks Analysis. <i>European Urology Oncology</i> , 2018 , 1, 346-351 | 6.7 | 4 |
| 286 | Endocrine, Sexual Function, and Infertility Side Effects of Immune Checkpoint Inhibitor Therapy for Genitourinary Cancers. <i>Current Urology Reports</i> , 2018 , 19, 68 | 2.9 | 6 |
| 285 | Tumor volume improves the long-term prediction of biochemical recurrence-free survival after radical prostatectomy for localized prostate cancer with positive surgical margins. <i>World Journal of Urology</i> , 2017 , 35, 199-206 | 4 | 14 |
| 284 | The Impact of Local Treatment on Overall Survival in Patients with Metastatic Prostate Cancer on Diagnosis: A National Cancer Data Base Analysis. <i>European Urology</i> , 2017 , 72, 14-19 | 10.2 | 97 |
| 283 | North American Population-Based Validation of the National Comprehensive Cancer Network Practice Guideline Recommendation of Pelvic Lymphadenectomy in Contemporary Prostate Cancer. <i>Prostate</i> , 2017 , 77, 542-548 | 4.2 | 13 |
| 282 | 10-Year Outcomes in Localized Prostate Cancer. <i>New England Journal of Medicine</i> , 2017 , 376, 180 | 59.2 | 12 |
| 281 | An Evaluation of the Timing of Surgical Complications Following Radical Cystectomy: Data From the American College of Surgeons National Surgical Quality Improvement Program. <i>Urology</i> , 2017 , 103, 91-98 | 1.6 | 22 |
| 280 | Calculating life expectancy to inform prostate cancer screening and treatment decisions. <i>BJU International</i> , 2017 , 120, 9-11 | 5.6 | 8 |
| 279 | Salvage Radiation Therapy for Biochemical Recurrence After Radical Prostatectomy: Is Earlier Always Better?. <i>Journal of Clinical Oncology</i> , 2017 , 35, 1489-1490 | 2.2 | 4 |
| 278 | Improved Survival With Local Treatment of Prostate Cancer in Men With Metastatic Disease: Look Before You Leap. <i>Journal of Clinical Oncology</i> , 2017 , 35, 914-915 | 2.2 | 1 |
| 277 | Could lead-time bias explain the apparent benefits of early salvage radiotherapy?. <i>Nature Reviews Urology</i> , 2017 , 14, 193-194 | 5.5 | 9 |
| 276 | Racial differences in prostate-specific antigen-based prostate cancer screening: State-by-state and region-by-region analyses. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017 , 35, 460.e9-460.e20 | 2.8 | 17 |
| 275 | Re: Diagnostic Accuracy of Multi-parametric Magnetic Resonance Imaging and Transrectal Ultrasound Biopsy in Prostate Cancer (PROMIS): A Paired Validating Confirmatory Study. <i>European Urology</i> , 2017 , 72, 315-316 | 10.2 | |
| 274 | Comparative Effectiveness of Trimodal Therapy Versus Radical Cystectomy for Localized Muscle-invasive Urothelial Carcinoma of the Bladder. <i>European Urology</i> , 2017 , 72, 483-487 | 10.2 | 65 |

| | | | |
|-----|---|------|-----|
| 273 | A Pragmatic Randomized Controlled Trial Examining the Impact of the Retzius-sparing Approach on Early Urinary Continence Recovery After Robot-assisted Radical Prostatectomy. <i>European Urology</i> , 2017 , 72, 677-685 | 10.2 | 99 |
| 272 | Impact of travel distance to the treatment facility on overall mortality in US patients with prostate cancer. <i>Cancer</i> , 2017 , 123, 3241-3252 | 6.4 | 54 |
| 271 | Efficacy of Systemic Chemotherapy Plus Radical Nephroureterectomy for Metastatic Upper Tract Urothelial Carcinoma. <i>European Urology</i> , 2017 , 71, 714-718 | 10.2 | 25 |
| 270 | Androgen Deprivation Therapy and Dose-Escalated Radiotherapy for Intermediate- and High-Risk Prostate Cancer: Sign of Changing Times?. <i>JAMA Oncology</i> , 2017 , 3, 280-281 | 13.4 | 1 |
| 269 | Therapeutic Value of Standard Versus Extended Pelvic Lymph Node Dissection During Radical Prostatectomy for High-Risk Prostate Cancer. <i>Current Urology Reports</i> , 2017 , 18, 51 | 2.9 | 10 |
| 268 | Local Therapy Improves Survival in Metastatic Prostate Cancer. <i>European Urology</i> , 2017 , 72, 118-124 | 10.2 | 77 |
| 267 | Genomic Classifier Augments the Role of Pathological Features in Identifying Optimal Candidates for Adjuvant Radiation Therapy in Patients With Prostate Cancer: Development and Internal Validation of a Multivariable Prognostic Model. <i>Journal of Clinical Oncology</i> , 2017 , 35, 1982-1990 | 2.2 | 56 |
| 266 | Individual Patient-Level Meta-Analysis of the Performance of the Decipher Genomic Classifier in High-Risk Men After Prostatectomy to Predict Development of Metastatic Disease. <i>Journal of Clinical Oncology</i> , 2017 , 35, 1991-1998 | 2.2 | 127 |
| 265 | Robot-Assisted Laparoscopic Repair of Extraperitoneal Ureteral Inguinal Hernia with Mesh Placement. <i>Journal of Endourology Case Reports</i> , 2017 , 3, 97-100 | 0.3 | 5 |
| 264 | Variation in Locoregional Prostate Cancer Care and Treatment Trends at Commission on Cancer Designated Facilities: A National Cancer Data Base Analysis 2004 to 2013. <i>Clinical Genitourinary Cancer</i> , 2017 , 15, e955-e968 | 3.3 | 12 |
| 263 | PD58-12 PSA SCREENING AT THE INTERSECTION OF POLITICS AND POLICY. <i>Journal of Urology</i> , 2017 , 197, | 2.5 | 1 |
| 262 | Editorial Comment. <i>Journal of Urology</i> , 2017 , 198, 361 | 2.5 | |
| 261 | PD32-05 PROSTATE CANCER SCREENING: EFFECT OF EARLY MEDICAID EXPANSION. <i>Journal of Urology</i> , 2017 , 197, | 2.5 | 1 |
| 260 | Emergency Department Utilization in Patients With Neurogenic Bladder: Contemporary Burden and National Trends in Prevalence, Inpatient Admission, and Associated Charges, 2006-2011. <i>Urology</i> , 2017 , 109, 74-81 | 1.6 | |
| 259 | Classification of Partial Nephrectomy as an Outpatient Surgery under CMS'Part B Reimbursement Program Does the Evidence Justify the Recommendation?. <i>Urology Practice</i> , 2017 , 4, 444-447 | 0.8 | 2 |
| 258 | Editorial Comment. <i>Journal of Urology</i> , 2017 , 198, 1067 | 2.5 | |
| 257 | Population-Based External Validation of the Updated 2012 Partin Tables in Contemporary North American Prostate Cancer Patients. <i>Prostate</i> , 2017 , 77, 105-113 | 4.2 | 20 |
| 256 | Postoperative sepsis prediction in patients undergoing major cancer surgery. <i>Journal of Surgical Research</i> , 2017 , 209, 60-69 | 2.5 | 9 |

| | | | |
|-----|--|------|----|
| 255 | Contemporary rates of pathological features and mortality for adenocarcinoma of the urinary bladder in the USA. <i>International Journal of Urology</i> , 2017 , 24, 117-123 | 2.3 | 12 |
| 254 | Generalizability of the Prostate Cancer Intervention Versus Observation Trial (PIVOT) Results to Contemporary North American Men with Prostate Cancer. <i>European Urology</i> , 2017 , 71, 511-514 | 10.2 | 14 |
| 253 | Efficacy of local treatment in patients with prostate cancer with clinically pelvic lymph node-positive disease at initial diagnosis.. <i>Journal of Clinical Oncology</i> , 2017 , 35, 164-164 | 2.2 | 1 |
| 252 | Genomic classifier to augment the role of pathological features in identifying optimal candidates for adjuvant radiation therapy in patients with prostate cancer: Development and internal validation of a multivariable prognostic model.. <i>Journal of Clinical Oncology</i> , 2017 , 35, 142-142 | 2.2 | |
| 251 | Is neoadjuvant chemotherapy beneficial before radical cystectomy? Examining the external validity of the SWOG-8710 trial.. <i>Journal of Clinical Oncology</i> , 2017 , 35, 331-331 | 2.2 | |
| 250 | Decipher correlation patterns post prostatectomy: initial experience from 2 342 prospective patients. <i>Prostate Cancer and Prostatic Diseases</i> , 2016 , 19, 374-379 | 6.2 | 28 |
| 249 | MP21-11 AGE DEPENDENT VARIATION IN THE EFFECT OF PHYSICIAN RECOMMENDATIONS TO UNDERGO PROSTATE SPECIFIC ANTIGEN (PSA) SCREENING FOLLOWING THE UNITED STATES PREVENTIVE SERVICES TASK FORCE 2012 STATEMENT AGAINST PSA SCREENING.. <i>Journal of Urology</i> , 2016 , 195, | 2.5 | 1 |
| 248 | MP39-14 IMPACT OF THE 2012 UNITED STATES PREVENTIVE SERVICES TASK FORCE RECOMMENDATION AGAINST PROSTATE SPECIFIC ANTIGEN SCREENING ON PROSTATE CANCER RISK GROUP STRATIFICATION. <i>Journal of Urology</i> , 2016 , 195, | 2.5 | 1 |
| 247 | PD15-03 DIFFERENCES IN PROSTATE SPECIFIC ANTIGEN TESTING AMONG UROLOGISTS AND PRIMARY CARE PROVIDERS IN THE UNITED STATES FOLLOWING THE 2011 USPSTF RECOMMENDATIONS. <i>Journal of Urology</i> , 2016 , 195, | 2.5 | 1 |
| 246 | MP02-06 EVALUATION OF A GENOMIC CLASSIFIER IN RADICAL PROSTATECTOMY PATIENTS WITH LYMPH NODE METASTASIS. <i>Journal of Urology</i> , 2016 , 195, | 2.5 | 2 |
| 245 | Utilization of a Genomic Classifier for Prediction of Metastasis Following Salvage Radiation Therapy after Radical Prostatectomy. <i>European Urology</i> , 2016 , 70, 588-596 | 10.2 | 56 |
| 244 | Prevalence of Nonrecommended Screening for Prostate Cancer and Breast Cancer in the United States: A Nationwide Survey Analysis. <i>JAMA Oncology</i> , 2016 , 2, 543-5 | 13.4 | 5 |
| 243 | Validation of a Genomic Classifier for Predicting Post-Prostatectomy Recurrence in a Community Based Health Care Setting. <i>Journal of Urology</i> , 2016 , 195, 1748-53 | 2.5 | 28 |
| 242 | Determinants of Prostate Specific Antigen Screening among Black Men in the United States in the Contemporary Era. <i>Journal of Urology</i> , 2016 , 195, 913-8 | 2.5 | 20 |
| 241 | Adverse Event Rates, Timing of Complications, and the Impact of Specialty on Outcomes Following Adrenal Surgery: An Analysis of 30-Day Outcome Data From the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP). <i>Urology</i> , 2016 , 90, 62-8 | 1.6 | 10 |
| 240 | Racial Differences in the Surgical Care of Medicare Beneficiaries With Localized Prostate Cancer. <i>JAMA Oncology</i> , 2016 , 2, 85-93 | 13.4 | 61 |
| 239 | Development and validation of genomic signature to predict ADT treatment failure.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 5018-5018 | 2.2 | 1 |
| 238 | Contemporary Role of the Decipher [®] Test in Prostate Cancer Management: Current Practice and Future Perspectives. <i>Reviews in Urology</i> , 2016 , 18, 1-9 | 1 | 19 |

| | | | |
|-----|---|------|----|
| 237 | Development and validation of an ADT resistance signature to predict adjuvant hormone treatment failure.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 106-106 | 2.2 | |
| 236 | Efficacy of early and delayed radiation in a prostatectomy cohort adjusted for genomic and clinical risk.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 12-12 | 2.2 | |
| 235 | The Technique of Robotic Nerve-Sparing Prostatectomy 2016 , 315-326 | | 1 |
| 234 | Evaluation of a genomic classifier in radical prostatectomy patients with lymph node metastasis. <i>Research and Reports in Urology</i> , 2016 , 8, 77-84 | 1.3 | 12 |
| 233 | Robot-assisted partial cystectomy with intraoperative frozen section examination: Evolution and evaluation of a novel technique. <i>Investigative and Clinical Urology</i> , 2016 , 57, 221-8 | 1.9 | 9 |
| 232 | The Authors Respond. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2016 , 14, 117-22 | 7.3 | |
| 231 | Treatment patterns, testicular loss and disparities in inpatient surgical management of testicular torsion in boys: a population-based study 1998-2010. <i>BJU International</i> , 2016 , 118, 969-979 | 5.6 | 4 |
| 230 | The importance of frailty: Know thy patient. <i>BJU International</i> , 2016 , 117, 716-7 | 5.6 | |
| 229 | Suicide and accidental deaths among patients with non-metastatic prostate cancer. <i>BJU International</i> , 2016 , 118, 286-97 | 5.6 | 30 |
| 228 | Dose-dependent effect of androgen deprivation therapy for localized prostate cancer on adverse cardiac events. <i>BJU International</i> , 2016 , 118, 221-9 | 5.6 | 16 |
| 227 | When Should a Positive Surgical Margin Ring a Bell? An Analysis of a Multi-Institutional Robot-Assisted Laparoscopic Radical Prostatectomy Database. <i>Journal of Endourology</i> , 2016 , 30, 201-7 | 2.7 | 10 |
| 226 | MP14-12 EFFICACY OF EARLY AND DELAYED RADIATION IN A PROSTATECTOMY COHORT ADJUSTED FOR GENOMIC AND CLINICAL RISK. <i>Journal of Urology</i> , 2016 , 195, | 2.5 | 1 |
| 225 | Reply to Michael Froehner, Rainer Koch, Manfred P. Wirth Letter to the Editor re: Jesse D. Sammon, Firas Abdollah, Anthony D'Amico, et al. Predicting Life Expectancy in Men Diagnosed with Prostate Cancer. <i>Eur Urol</i> 2015;68:756-65. <i>European Urology</i> , 2016 , 69, e129 | 10.2 | |
| 224 | PD15-01 THE IMPACT OF 2011 UNITED STATES PREVENTIVE SERVICES TASK FORCE PANEL UPDATE ON PSA SCREENING PRACTICE: A NATIONWIDE, AND STATE-BY-STATE LEVEL ANALYSES.. <i>Journal of Urology</i> , 2016 , 195, | 2.5 | 2 |
| 223 | PD43-12 URINARY CONTINENCE OUTCOMES AFTER RETZIUS-SPARING ROBOT-ASSISTED RADICAL PROSTATECTOMY: A PROSPECTIVE, NON-RANDOMIZED, IDEAL STAGE 2B (EXPLORATION) STUDY.. <i>Journal of Urology</i> , 2016 , 195, | 2.5 | 3 |
| 222 | Determinants of long-term survival of patients with locally advanced prostate cancer: the role of extensive pelvic lymph node dissection. <i>Prostate Cancer and Prostatic Diseases</i> , 2016 , 19, 63-7 | 6.2 | 36 |
| 221 | Intermediate-term cancer control outcomes in prostate cancer patients treated with robotic-assisted laparoscopic radical prostatectomy: a multi-institutional analysis. <i>World Journal of Urology</i> , 2016 , 34, 1357-66 | 4 | 7 |
| 220 | Understanding the Use of Prostate Biopsy Among Men with Limited Life Expectancy in a Statewide Quality Improvement Collaborative. <i>European Urology</i> , 2016 , 70, 854-861 | 10.2 | 9 |

| | | | |
|-----|---|------|----|
| 219 | Efficacy of post-operative radiation in a prostatectomy cohort adjusted for clinical and genomic risk. <i>Prostate Cancer and Prostatic Diseases</i> , 2016 , 19, 277-82 | 6.2 | 17 |
| 218 | The importance of adjuvant therapy in patients with node-positive prostate cancer: A nationwide validation study. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016 , 34, 577-578 | 2.8 | |
| 217 | The diminishing returns of robotic diffusion: complications after robot-assisted radical prostatectomy. <i>BJU International</i> , 2016 , 117, 211-2 | 5.6 | 4 |
| 216 | Predicting lymph node invasion in patients treated with robot-assisted radical prostatectomy. <i>Canadian Journal of Urology</i> , 2016 , 23, 8141-50 | 0.8 | 2 |
| 215 | Reply to Jovo Bogdanović and Vuk Sekulić Letter to the Editor re: Firas Abdollah, Giorgio Gandaglia, Nazareno Suardi, et al. More Extensive Pelvic Lymph Node Dissection Improves Survival in Patients with Node-positive Prostate Cancer. <i>Eur Urol</i> 2015;67:212-9. <i>European Urology</i> , 2015 , 68, e37-8 | 10.2 | 1 |
| 214 | Re: AR-V7 and Resistance to Enzalutamide and Abiraterone in Prostate Cancer. <i>European Urology</i> , 2015 , 68, 162-3 | 10.2 | 3 |
| 213 | A Comparison of 30-Day Perioperative Outcomes in Open Versus Minimally Invasive Nephroureterectomy for Upper Tract Urothelial Carcinoma: Analysis of 896 Patients from the American College of Surgeons-National Surgical Quality Improvement Program Database. <i>Journal of Endourology</i> , 2015 , 29, 1052-8 | 2.7 | 18 |
| 212 | Preventable mortality after common urological surgery: failing to rescue?. <i>BJU International</i> , 2015 , 115, 666-74 | 5.6 | 10 |
| 211 | Long-term cancer control outcomes in patients with clinically high-risk prostate cancer treated with robot-assisted radical prostatectomy: results from a multi-institutional study of 1100 patients. <i>European Urology</i> , 2015 , 68, 497-505 | 10.2 | 63 |
| 210 | Effect of preoperative angina pectoris on cardiac outcomes in patients with previous myocardial infarction undergoing major noncardiac surgery (data from ACS-NSQIP). <i>American Journal of Cardiology</i> , 2015 , 115, 1080-4 | 3 | 11 |
| 209 | Comparison of 30-day perioperative outcomes in adults undergoing open versus minimally invasive pyeloplasty for ureteropelvic junction obstruction: analysis of 593 patients in a prospective national database. <i>World Journal of Urology</i> , 2015 , 33, 2107-13 | 4 | 9 |
| 208 | An evaluation of the timing of surgical complications following nephrectomy: data from the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP). <i>World Journal of Urology</i> , 2015 , 33, 2031-8 | 4 | 20 |
| 207 | Predicting Life Expectancy in Men Diagnosed with Prostate Cancer. <i>European Urology</i> , 2015 , 68, 756-65 | 10.2 | 43 |
| 206 | The Effect of Body Mass Index on Perioperative Outcomes After Major Surgery: Results from the National Surgical Quality Improvement Program (ACS-NSQIP) 2005-2011. <i>World Journal of Surgery</i> , 2015 , 39, 2376-85 | 3.3 | 47 |
| 205 | Early radiotherapy after radical prostatectomy improves cancer-specific survival only in patients with highly aggressive prostate cancer: validation of recently released criteria. <i>International Journal of Urology</i> , 2015 , 22, 89-95 | 2.3 | 20 |
| 204 | Reply to C.G. Rusthoven et al. <i>Journal of Clinical Oncology</i> , 2015 , 33, 1989 | 2.2 | 1 |
| 203 | Rates of Kidney Transplantation From Living and Deceased Donors for Blacks and Whites in the United States, 1998 to 2011. <i>JAMA Internal Medicine</i> , 2015 , 175, 1716-8 | 11.5 | 7 |
| 202 | Contemporary nationwide patterns of self-reported prostate-specific antigen screening in US veterans. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 503.e7-15 | 2.8 | 7 |

| | | | |
|-----|---|------|-----|
| 201 | Medical androgen deprivation therapy and increased non-cancer mortality in non-metastatic prostate cancer patients aged ≥ 66 years. <i>European Journal of Surgical Oncology</i> , 2015 , 41, 1529-39 | 3.6 | 14 |
| 200 | The influence of physician recommendation on prostate-specific antigen screening. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 424.e1-7 | 2.8 | 21 |
| 199 | Adjuvant Radiotherapy in Prostate Cancer Patients Treated with Surgery: The Impact of Age and Tumor Characteristics. <i>European Urology Focus</i> , 2015 , 1, 191-199 | 5.1 | 5 |
| 198 | Prostate-Specific Antigen Screening After 2012 US Preventive Services Task Force Recommendations. <i>JAMA - Journal of the American Medical Association</i> , 2015 , 314, 2077-9 | 27.4 | 84 |
| 197 | Identifying optimal candidates for local treatment of the primary tumor among patients diagnosed with metastatic prostate cancer: a SEER-based study. <i>European Urology</i> , 2015 , 67, 3-6 | 10.2 | 95 |
| 196 | Contemporary role of salvage lymphadenectomy in patients with recurrence following radical prostatectomy. <i>European Urology</i> , 2015 , 67, 839-49 | 10.2 | 63 |
| 195 | More extensive pelvic lymph node dissection improves survival in patients with node-positive prostate cancer. <i>European Urology</i> , 2015 , 67, 212-9 | 10.2 | 134 |
| 194 | MP32-08 READMISSIONS AFTER MAJOR UROLOGIC CANCER SURGERY. <i>Journal of Urology</i> , 2015 , 193, | 2.5 | 1 |
| 193 | The Impact of Insurance Status on Tumor Characteristics and Treatment Selection in Contemporary Patients With Prostate Cancer. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2015 , 13, 1351-8 | 7.3 | 15 |
| 192 | Racial Disparities in End-of-Life Care Among Patients With Prostate Cancer: A Population-Based Study. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2015 , 13, 1131-8 | 7.3 | 30 |
| 191 | Racial/Ethnic Disparities in Perioperative Outcomes of Major Procedures: Results From the National Surgical Quality Improvement Program. <i>Annals of Surgery</i> , 2015 , 262, 955-64 | 7.8 | 69 |
| 190 | PD6-12 DIMINISHING RETURNS OF ROBOTIC DIFFUSION: COMPLICATIONS FOLLOWING ROBOT-ASSISTED RADICAL PROSTATECTOMY. <i>Journal of Urology</i> , 2015 , 193, | 2.5 | 1 |
| 189 | Minimally invasive renal autotransplantation. <i>Journal of Surgical Oncology</i> , 2015 , 112, 717-22 | 2.8 | 10 |
| 188 | An evaluation of the weekend effect on patients admitted with metastatic prostate cancer. <i>BJU International</i> , 2015 , 116, 911-9 | 5.6 | 7 |
| 187 | Robot-assisted hepatic mobilization and control of suprahepatic infradiaphragmatic inferior vena cava for level 3 vena caval thrombectomy: An IDEAL stage 0 study. <i>Journal of Surgical Oncology</i> , 2015 , 112, 741-5 | 2.8 | 8 |
| 186 | Predicting pathological outcomes in patients undergoing robot-assisted radical prostatectomy for high-risk prostate cancer: a preoperative nomogram. <i>BJU International</i> , 2015 , 116, 703-12 | 5.6 | 6 |
| 185 | Patterns of Declining Use and the Adverse Effect of Primary Androgen Deprivation on All-cause Mortality in Elderly Men with Prostate Cancer. <i>European Urology</i> , 2015 , 68, 32-9 | 10.2 | 35 |
| 184 | Genomic classifier identifies men with adverse pathology after radical prostatectomy who benefit from adjuvant radiation therapy. <i>Journal of Clinical Oncology</i> , 2015 , 33, 944-51 | 2.2 | 151 |

| | | | |
|-----|---|------|-----|
| 183 | The Role of Biomarkers and Genetics in the Diagnosis of Prostate Cancer. <i>European Urology Focus</i> , 2015 , 1, 99-108 | 5.1 | 6 |
| 182 | Differential effect on survival of pelvic lymph node dissection at radical cystectomy for muscle invasive bladder cancer. <i>European Journal of Surgical Oncology</i> , 2015 , 41, 353-60 | 3.6 | 28 |
| 181 | A genomic classifier to identify men with adverse pathology post radical prostatectomy who benefit from adjuvant radiation therapy.. <i>Journal of Clinical Oncology</i> , 2015 , 33, 168-168 | 2.2 | |
| 180 | Prevalence of non-recommended screening for prostate cancer and breast cancer in the United States.. <i>Journal of Clinical Oncology</i> , 2015 , 33, e17528-e17528 | 2.2 | |
| 179 | Gonadotropin-releasing hormone agonists and acute kidney injury in patients with prostate cancer. <i>European Urology</i> , 2014 , 66, 1125-32 | 10.2 | 21 |
| 178 | Impact of adjuvant radiation therapy on urinary continence recovery after radical prostatectomy. <i>European Urology</i> , 2014 , 65, 546-51 | 10.2 | 68 |
| 177 | Head-to-head comparison of lymph node density and number of positive lymph nodes in stratifying the outcome of patients with lymph node-positive prostate cancer submitted to radical prostatectomy and extended lymph node dissection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 29.e21-8 | 2.8 | 20 |
| 176 | Distribution of metastatic sites in patients with prostate cancer: A population-based analysis. <i>Prostate</i> , 2014 , 74, 210-6 | 4.2 | 208 |
| 175 | The effect of neoadjuvant chemotherapy on perioperative outcomes in patients who have bladder cancer treated with radical cystectomy: a population-based study. <i>European Urology</i> , 2014 , 66, 561-8 | 10.2 | 52 |
| 174 | Survival benefit of definitive therapy in patients with clinically advanced prostate cancer: estimations of the number needed to treat based on competing-risks analysis. <i>BJU International</i> , 2014 , 114, E62-E69 | 5.6 | 14 |
| 173 | The number of cores at first biopsy may suggest the need for a confirmatory biopsy in patients eligible for active surveillance-implication for clinical decision making in the real-life setting. <i>Urology</i> , 2014 , 84, 634-41 | 1.6 | 6 |
| 172 | Is robot-assisted radical prostatectomy safe in men with high-risk prostate cancer? Assessment of perioperative outcomes, positive surgical margins, and use of additional cancer treatments. <i>Journal of Endourology</i> , 2014 , 28, 784-91 | 2.7 | 36 |
| 171 | 11C-choline PET/CT predicts prostate cancer-specific survival in patients with biochemical failure during androgen-deprivation therapy. <i>Journal of Nuclear Medicine</i> , 2014 , 55, 233-41 | 8.9 | 71 |
| 170 | Pathologic nodal staging scores in patients treated with radical prostatectomy: a postoperative decision tool. <i>European Urology</i> , 2014 , 66, 439-46 | 10.2 | 18 |
| 169 | The impact of robot-assisted radical prostatectomy on the use and extent of pelvic lymph node dissection in the "post-dissemination" period. <i>European Journal of Surgical Oncology</i> , 2014 , 40, 1080-6 | 3.6 | 40 |
| 168 | Management of localized kidney cancer: calculating cancer-specific mortality and competing risks of death for surgery and nonsurgical management. <i>European Urology</i> , 2014 , 65, 235-41 | 10.2 | 83 |
| 167 | The key role of time in predicting progression-free survival in patients with renal cell carcinoma treated with partial or radical nephrectomy: conditional survival analysis. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 43.e9-16 | 2.8 | 10 |
| 166 | Long-term survival in patients with germ cell testicular cancer: a population-based competing-risks regression analysis. <i>European Journal of Surgical Oncology</i> , 2014 , 40, 103-12 | 3.6 | 19 |

| | | | |
|-----|---|------|-----|
| 165 | PD12-12 ADJUVANT RADIOTHERAPY IMPROVES CANCER-SPECIFIC SURVIVAL ONLY IN PATIENTS WITH HIGHLY AGGRESSIVE PROSTATE CANCER. VALIDATION OF RECENTLY RELEASED CRITERIA. <i>Journal of Urology</i> , 2014 , 191, | 2.5 | 1 |
| 164 | Comparative effectiveness of robot-assisted versus open radical prostatectomy cancer control. <i>European Urology</i> , 2014 , 66, 666-72 | 10.2 | 81 |
| 163 | PD12-11 SURVIVAL BENEFIT OF RADICAL PROSTATECTOMY IN PATIENTS WITH CLINICALLY ADVANCED PROSTATE CANCER: ESTIMATIONS OF THE NUMBER NEEDED TO TREAT BASED ON COMPETING-RISKS ANALYSIS. <i>Journal of Urology</i> , 2014 , 191, | 2.5 | 2 |
| 162 | A population-based competing-risks analysis of survival after nephrectomy for renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 46.e1-7 | 2.8 | 18 |
| 161 | The effect of age at diagnosis on prostate cancer mortality: a grade-for-grade and stage-for-stage analysis. <i>European Journal of Surgical Oncology</i> , 2014 , 40, 1706-15 | 3.6 | 15 |
| 160 | Extent of lymphadenectomy does not improve the survival of patients with renal cell carcinoma and nodal metastases: biases associated with the handling of missing data. <i>BJU International</i> , 2014 , 113, 36-42 | 5.6 | 27 |
| 159 | Clinical nodal staging scores for prostate cancer: a proposal for preoperative risk assessment. <i>British Journal of Cancer</i> , 2014 , 111, 213-9 | 8.7 | 21 |
| 158 | External validation of the European association of urology recommendations for pelvic lymph node dissection in patients treated with robot-assisted radical prostatectomy. <i>Journal of Endourology</i> , 2014 , 28, 416-23 | 2.7 | 29 |
| 157 | Contemporary incidence and mortality rates of kidney cancer in the United States. <i>Canadian Urological Association Journal</i> , 2014 , 8, 247-52 | 1.2 | 68 |
| 156 | Preoperative erectile function is the only predictor of the use of a high number of phosphodiesterase type-5 inhibitors after bilateral nerve-sparing radical prostatectomy. <i>International Journal of Impotence Research</i> , 2014 , 26, 201-4 | 2.3 | 8 |
| 155 | Intensity-modulated radiation therapy leads to survival benefit only in patients with high-risk prostate cancer: a population-based study. <i>Annals of Oncology</i> , 2014 , 25, 979-86 | 10.3 | 7 |
| 154 | Radical prostatectomy vs radiotherapy vs observation among older patients with clinically localized prostate cancer: a comparative effectiveness evaluation. <i>BJU International</i> , 2014 , 113, 200-8 | 5.6 | 48 |
| 153 | Extent of lymph node dissection at nephrectomy affects cancer-specific survival and metastatic progression in specific sub-categories of patients with renal cell carcinoma (RCC). <i>BJU International</i> , 2014 , 114, 210-5 | 5.6 | 51 |
| 152 | Survival after nephroureterectomy for upper tract urothelial carcinoma: a population-based competing-risks analysis. <i>International Journal of Urology</i> , 2014 , 21, 249-56 | 2.3 | 17 |
| 151 | Benefit in regionalisation of care for patients treated with radical cystectomy: a nationwide inpatient sample analysis. <i>BJU International</i> , 2014 , 113, 733-40 | 5.6 | 21 |
| 150 | Impact of adjuvant radiotherapy on survival of patients with node-positive prostate cancer. <i>Journal of Clinical Oncology</i> , 2014 , 32, 3939-47 | 2.2 | 184 |
| 149 | Percentage of high-grade tumour volume does not meaningfully improve prediction of early biochemical recurrence after radical prostatectomy compared with Gleason score. <i>BJU International</i> , 2014 , 113, 399-407 | 5.6 | 18 |
| 148 | In reply to the letter to the editor On Reply to Gandaglia et al. By De Bari et al. <i>Annals of Oncology</i> , 2014 , 25, 1862-1863 | 10.3 | |

| | | | |
|-----|--|------|----|
| 147 | The impact of resident involvement in minimally-invasive urologic oncology procedures. <i>Canadian Urological Association Journal</i> , 2014 , 8, 334-40 | 1.2 | 38 |
| 146 | Predicting the risk of bone metastasis in prostate cancer. <i>Cancer Treatment Reviews</i> , 2014 , 40, 3-11 | 14.4 | 43 |
| 145 | MP51-15 TIME FROM SURGERY TO URINARY CONTINENCE SIGNIFICANTLY INFLUENCES THE SUBSEQUENT RECOVERY OF ERECTILE FUNCTION IN PATIENTS TREATED WITH BILATERAL NERVE-SPARING RADICAL PROSTATECTOMY. <i>Journal of Urology</i> , 2014 , 191, | 2.5 | 2 |
| 144 | Utility of [11C]choline PET/CT in guiding lesion-targeted salvage therapies in patients with prostate cancer recurrence localized to a single lymph node at imaging: results from a pathologically validated series. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 38.e9-16 | 2.8 | 54 |
| 143 | Predicting survival of patients with node-positive prostate cancer following multimodal treatment. <i>European Urology</i> , 2014 , 65, 554-62 | 10.2 | 61 |
| 142 | The impact of androgen-deprivation therapy (ADT) on the risk of cardiovascular (CV) events in patients with non-metastatic prostate cancer: a population-based study. <i>BJU International</i> , 2014 , 114, E82-E89 | 5.6 | 66 |
| 141 | Comparative effectiveness of radical cystectomy versus bladder-sparing treatment for muscle-invasive urothelial carcinoma: A population-based report.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 334-334 | 2.2 | 1 |
| 140 | Role of robot-assisted radical prostatectomy in the management of high-risk prostate cancer. <i>Indian Journal of Urology</i> , 2014 , 30, 410-7 | 0.8 | 11 |
| 139 | Using adjuvant radiotherapy to improve cancer-specific survival in patients with highly aggressive prostate cancer: Examining recently released criteria.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 30-30 | 2.2 | |
| 138 | Population-based comparison of surgical margin status for robotic versus open radical prostatectomy.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 51-51 | 2.2 | |
| 137 | Comparison of partial vs radical nephrectomy with regard to other-cause mortality in T1 renal cell carcinoma among patients aged ≥ 5 years with multiple comorbidities. <i>BJU International</i> , 2013 , 111, 67-73 | 5.6 | 46 |
| 136 | Staging lymphadenectomy in renal cell carcinoma must be extended: a sensitivity curve analysis. <i>BJU International</i> , 2013 , 111, 412-8 | 5.6 | 15 |
| 135 | Nerve-sparing approach during radical prostatectomy is strongly associated with the rate of postoperative urinary continence recovery. <i>BJU International</i> , 2013 , 111, 717-22 | 5.6 | 73 |
| 134 | Conditional survival after nephrectomy for renal cell carcinoma (RCC): changes in future survival probability over time. <i>BJU International</i> , 2013 , 111, E283-9 | 5.6 | 26 |
| 133 | When to perform lymph node dissection in patients with renal cell carcinoma: a novel approach to the preoperative assessment of risk of lymph node invasion at surgery and of lymph node progression during follow-up. <i>BJU International</i> , 2013 , 112, E59-66 | 5.6 | 35 |
| 132 | Preoperative sex steroids are significant predictors of early biochemical recurrence after radical prostatectomy. <i>World Journal of Urology</i> , 2013 , 31, 275-80 | 4 | 25 |
| 131 | Pelvic lymph node dissection for prostate cancer: adherence and accuracy of the recent guidelines. <i>International Journal of Urology</i> , 2013 , 20, 405-10 | 2.3 | 18 |
| 130 | Impact of age and comorbidities on long-term survival of patients with high-risk prostate cancer treated with radical prostatectomy: a multi-institutional competing-risks analysis. <i>European Urology</i> , 2013 , 63, 693-701 | 10.2 | 78 |

| | | | |
|-----|---|------|-----|
| 129 | Assessing the most accurate formula to predict the risk of lymph node metastases from prostate cancer in contemporary patients treated with radical prostatectomy and extended pelvic lymph node dissection. <i>Radiotherapy and Oncology</i> , 2013 , 109, 211-6 | 5.3 | 15 |
| 128 | An evidence-based guide to the selection of sequential therapies in metastatic renal cell carcinoma. <i>Therapeutic Advances in Urology</i> , 2013 , 5, 121-8 | 3.2 | 15 |
| 127 | Postoperative phosphodiesterase type 5 inhibitor administration increases the rate of urinary continence recovery after bilateral nerve-sparing radical prostatectomy. <i>International Journal of Urology</i> , 2013 , 20, 413-9 | 2.3 | 17 |
| 126 | A population-based analysis of the effect of marital status on overall and cancer-specific mortality in patients with squamous cell carcinoma of the penis. <i>Cancer Causes and Control</i> , 2013 , 24, 71-9 | 2.8 | 28 |
| 125 | 774 HEAD-TO-HEAD COMPARISON OF LYMPH NODE DENSITY AND NUMBER OF POSITIVE LYMPH NODES IN STRATIFYING THE OUTCOME OF PATIENTS WITH LYMPH NODE POSITIVE PROSTATE CANCER SUBMITTED TO RADICAL PROSTATECTOMY AND EXTENDED PELVIC LYMPH NODE DISSECTION. <i>Journal of Urology</i> , 2013 , 189, | 2.5 | 1 |
| 124 | The number of cores taken in patients diagnosed with a single microfocus at initial biopsy is a major predictor of insignificant prostate cancer. <i>Journal of Urology</i> , 2013 , 189, 854-9 | 2.5 | 11 |
| 123 | Survival following biochemical recurrence after radical prostatectomy and adjuvant radiotherapy in patients with prostate cancer: the impact of competing causes of mortality and patient stratification. <i>European Urology</i> , 2013 , 64, 557-64 | 10.2 | 33 |
| 122 | 1838 WHEN TO PERFORM LYMPH NODE DISSECTION IN RENAL CELL CARCINOMA PATIENTS: A NOVEL APPROACH TO PREOPERATIVELY ASSESS THE RISK OF LYMPH NODE INVASION AT SURGERY AND NODAL PROGRESSION DURING FOLLOW UP. <i>Journal of Urology</i> , 2013 , 189, | 2.5 | 1 |
| 121 | 961 PREDICTORS OF EARLY BIOCHEMICAL RECURRENCE AFTER RADICAL PROSTATECTOMY AND ADJUVANT RADIOTHERAPY IN MEN WITH PT3N0 PROSTATE CANCER. IMPLICATIONS FOR MULTI-MODAL THERAPIES. <i>Journal of Urology</i> , 2013 , 189, | 2.5 | 1 |
| 120 | 1831 THE NUMBER OF LYMPH NODES REMOVED IN RENAL CELL CARCINOMA DOES AFFECT CANCER SPECIFIC SURVIVAL IN SPECIFIC SUBGROUPS OF PATIENTS: RESULTS FROM A SYSTEMATIC ANALYSIS. <i>Journal of Urology</i> , 2013 , 189, | 2.5 | 2 |
| 119 | Head-to-head comparison of prostate health index and urinary PCA3 for predicting cancer at initial or repeat biopsy. <i>Journal of Urology</i> , 2013 , 190, 496-501 | 2.5 | 72 |
| 118 | Presence of positive surgical margin in patients with organ-confined prostate cancer equals to extracapsular extension negative surgical margin. A plea for TNM staging system reclassification. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1497-503 | 2.8 | 11 |
| 117 | 1634 MANAGEMENT OF LOCALIZED KIDNEY CANCER: CALCULATING CANCER-SPECIFIC MORTALITY AND COMPETING-RISKS OF DEATH TRADEOFFS BETWEEN SURGERY AND ACTIVE SURVEILLANCE. <i>Journal of Urology</i> , 2013 , 189, | 2.5 | 2 |
| 116 | Selecting the optimal candidate for adjuvant radiotherapy after radical prostatectomy for prostate cancer: a long-term survival analysis. <i>European Urology</i> , 2013 , 63, 998-1008 | 10.2 | 88 |
| 115 | In-hospital mortality and failure to rescue after cytoreductive nephrectomy. <i>European Urology</i> , 2013 , 63, 1107-14 | 10.2 | 45 |
| 114 | Reply to Chris Parker, Matthew R. Sydes and Howard Kynaston letter to the editor re: Firas Abdollah, Nazareno Suardi, Cesare Cozzarini, et al. Selecting the optimal candidate for adjuvant radiotherapy after radical prostatectomy for prostate cancer: a long-term survival analysis. <i>Eur Urol</i> 2013;63:998-1008. <i>European Urology</i> , 2013 , 64, e103-4 | 10.2 | 1 |
| 113 | Incidence, survival and mortality rates of stage-specific bladder cancer in United States: a trend analysis. <i>Cancer Epidemiology</i> , 2013 , 37, 219-25 | 2.8 | 178 |
| 112 | Circulating sex steroids and prostate cancer: introducing the time-dependency theory. <i>World Journal of Urology</i> , 2013 , 31, 267-73 | 4 | 5 |

| | | | |
|-----|--|------|-----|
| 111 | Nodal involvement at nephrectomy is associated with worse survival: a stage-for-stage and grade-for-grade analysis. <i>International Journal of Urology</i> , 2013 , 20, 372-80 | 2.3 | 16 |
| 110 | An analysis of patients with T2 renal cell carcinoma (RCC) according to tumour size: a population-based analysis. <i>BJU International</i> , 2013 , 111, 1184-90 | 5.6 | 3 |
| 109 | Impact of the introduction of a robotic training programme on prostate cancer stage migration at a single tertiary referral centre. <i>BJU International</i> , 2013 , 111, 1222-30 | 5.6 | 28 |
| 108 | Competing-risks analysis in patients with T1 squamous cell carcinoma of the penis. <i>BJU International</i> , 2013 , 111, E174-9 | 5.6 | 7 |
| 107 | A novel tool to assess the risk of urinary incontinence after nerve-sparing radical prostatectomy. <i>BJU International</i> , 2013 , 111, 905-13 | 5.6 | 13 |
| 106 | External validation of the updated nomogram predicting lymph node invasion in patients with prostate cancer undergoing extended pelvic lymph node dissection. <i>Urologia Internationalis</i> , 2013 , 90, 277-82 | 1.9 | 27 |
| 105 | Extended pelvic lymph node dissection in prostate cancer: a 20-year audit in a single center. <i>Annals of Oncology</i> , 2013 , 24, 1459-66 | 10.3 | 65 |
| 104 | Effect of number and location of distant metastases on renal cell carcinoma mortality in candidates for cytoreductive nephrectomy: implications for multimodal therapy. <i>International Journal of Urology</i> , 2013 , 20, 572-9 | 2.3 | 19 |
| 103 | Response to Re: Lymph node count threshold for optimal pelvic lymph node staging in prostate cancer. <i>International Journal of Urology</i> , 2013 , 20, 845-6 | 2.3 | 1 |
| 102 | Rates of open versus laparoscopic and partial versus radical nephrectomy for T1a renal cell carcinoma: a population-based evaluation. <i>International Journal of Urology</i> , 2013 , 20, 1064-71 | 2.3 | 32 |
| 101 | Response to Re: Comparison of mortality outcomes after radical prostatectomy versus radiotherapy in patients with localized prostate cancer: a population-based analysis. <i>International Journal of Urology</i> , 2013 , 20, 548-9 | 2.3 | |
| 100 | Spatial distribution of positive cores improves the selection of patients with low-risk prostate cancer as candidates for active surveillance. <i>BJU International</i> , 2013 , 112, E234-42 | 5.6 | 1 |
| 99 | Clinical nodal staging scores for bladder cancer: a proposal for preoperative risk assessment. <i>European Urology</i> , 2012 , 61, 237-42 | 10.2 | 54 |
| 98 | Updated nomogram predicting lymph node invasion in patients with prostate cancer undergoing extended pelvic lymph node dissection: the essential importance of percentage of positive cores. <i>European Urology</i> , 2012 , 61, 480-7 | 10.2 | 425 |
| 97 | Reply to Michael Froehner Letter to the Editor re: Firas Abdollah, Maxine Sun, Jan Schmitges, et al. Cancer-Specific and Other-Cause Mortality After Radical Prostatectomy Versus Observation in Patients with Prostate Cancer: Competing-Risks Analysis of a Large North American Population-Based Cohort. <i>Eur Urol</i> 2011;60:920B0. <i>European Urology</i> , 2012 , 61, e12 | 10.2 | |
| 96 | Performance characteristics of computed tomography in detecting lymph node metastases in contemporary patients with prostate cancer treated with extended pelvic lymph node dissection. <i>European Urology</i> , 2012 , 61, 1132-8 | 10.2 | 95 |
| 95 | Identifying the best candidate for radical prostatectomy among patients with high-risk prostate cancer. <i>European Urology</i> , 2012 , 61, 584-92 | 10.2 | 94 |
| 94 | A non-cancer-related survival benefit is associated with partial nephrectomy. <i>European Urology</i> , 2012 , 61, 725-31 | 10.2 | 95 |

| | | | |
|----|--|------|-----|
| 93 | Prediction of functional outcomes after nerve-sparing radical prostatectomy: results of conditional survival analyses. <i>European Urology</i> , 2012 , 62, 42-52 | 10.2 | 54 |
| 92 | Are Caucasian-European men delaying fatherhood? Results of a 7 year observational study of infertile couples with male factor infertility. <i>Journal of Developmental and Physical Disabilities</i> , 2012 , 35, 125-32 | | 13 |
| 91 | Choosing the best candidates for penile rehabilitation after bilateral nerve-sparing radical prostatectomy. <i>Journal of Sexual Medicine</i> , 2012 , 9, 608-17 | 1.1 | 29 |
| 90 | Erectile function outcome after bilateral nerve sparing radical prostatectomy: which patients may be left untreated?. <i>Journal of Sexual Medicine</i> , 2012 , 9, 903-8 | 1.1 | 29 |
| 89 | Is a treatment delay in radical prostatectomy safe in individuals with low-risk prostate cancer?. <i>Journal of Sexual Medicine</i> , 2012 , 9, 2961-9 | 1.1 | 25 |
| 88 | Extended pelvic lymph node dissection does not affect erectile function recovery in patients treated with bilateral nerve-sparing radical prostatectomy. <i>Journal of Sexual Medicine</i> , 2012 , 9, 2187-94 | 1.1 | 17 |
| 87 | Comparison of mortality outcomes after radical prostatectomy versus radiotherapy in patients with localized prostate cancer: a population-based analysis. <i>International Journal of Urology</i> , 2012 , 19, 836-44 | 2.3 | 76 |
| 86 | Survival after radical cystectomy of non-bilharzial squamous cell carcinoma vs urothelial carcinoma: a competing-risks analysis. <i>BJU International</i> , 2012 , 109, 564-9 | 5.6 | 32 |
| 85 | Influence of obesity on tumour volume in patients with prostate cancer. <i>BJU International</i> , 2012 , 109, 678-84 | 5.6 | 22 |
| 84 | Stage-specific impact of pelvic lymph node dissection on survival in patients with non-metastatic bladder cancer treated with radical cystectomy. <i>BJU International</i> , 2012 , 109, 1147-54 | 5.6 | 52 |
| 83 | A population-based assessment of the National Comprehensive Cancer Network practice guideline indications for pelvic lymph node dissection at radical prostatectomy. <i>BJU International</i> , 2012 , 109, 1177-82 | 5.6 | 11 |
| 82 | Diagnosis of isolated high-grade prostatic intra-epithelial neoplasia: proposal of a nomogram for the prediction of cancer detection at saturation re-biopsy. <i>BJU International</i> , 2012 , 109, 1329-34 | 5.6 | 18 |
| 81 | Cytoreductive nephrectomy in the elderly: a population-based cohort from the USA. <i>BJU International</i> , 2012 , 109, 1807-12 | 5.6 | 26 |
| 80 | Unilateral positive biopsies in low risk prostate cancer patients diagnosed with extended transrectal ultrasound-guided biopsy schemes do not predict unilateral prostate cancer at radical prostatectomy. <i>BJU International</i> , 2012 , 110, E64-8 | 5.6 | 27 |
| 79 | Venous thromboembolism after radical prostatectomy: the effect of surgical caseload. <i>BJU International</i> , 2012 , 110, 828-33 | 5.6 | 13 |
| 78 | Treatment management of small renal masses in the 21st century: a paradigm shift. <i>Annals of Surgical Oncology</i> , 2012 , 19, 2380-7 | 3.1 | 74 |
| 77 | Preoperative circulating sex hormones are not predictors of positive surgical margins at open radical prostatectomy. <i>World Journal of Urology</i> , 2012 , 30, 533-9 | 4 | 6 |
| 76 | Chronic kidney disease after nephrectomy in patients with small renal masses: a retrospective observational analysis. <i>European Urology</i> , 2012 , 62, 696-703 | 10.2 | 110 |

| | | | |
|----|---|-----|----|
| 75 | Indications for pelvic nodal treatment in prostate cancer should change. Validation of the Roach formula in a large extended nodal dissection series. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012 , 83, 624-9 | 4 | 20 |
| 74 | Competing-risks mortality after radiotherapy vs. observation for localized prostate cancer: a population-based study. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012 , 84, 95-103 | 4 | 21 |
| 73 | In reply to Yu. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012 , 84, 301-2 | 4 | |
| 72 | Hospital volume is a determinant of postoperative complications, blood transfusion and length of stay after radical or partial nephrectomy. <i>Journal of Urology</i> , 2012 , 187, 405-10 | 2.5 | 50 |
| 71 | Preoperative erectile function represents a significant predictor of postoperative urinary continence recovery in patients treated with bilateral nerve sparing radical prostatectomy. <i>Journal of Urology</i> , 2012 , 187, 569-74 | 2.5 | 30 |
| 70 | 982 STAGING LYMPHADENECTOMY IN RENAL CELL CARCINOMA MUST BE EXTENDED: A SENSITIVITY CURVE ANALYSES. <i>Journal of Urology</i> , 2012 , 187, | 2.5 | 1 |
| 69 | 771 SHOULD AN EXTENDED NODAL TEMPLATE FOR HIGH RISK PROSTATE CANCER ALWAYS INCLUDE REMOVAL OF COMMON ILIAC LYMPH NODES?. <i>Journal of Urology</i> , 2012 , 187, | 2.5 | 1 |
| 68 | Survival benefit of radical prostatectomy in patients with localized prostate cancer: estimations of the number needed to treat according to tumor and patient characteristics. <i>Journal of Urology</i> , 2012 , 188, 73-83 | 2.5 | 35 |
| 67 | Blood transfusions in radical prostatectomy: a contemporary population-based analysis. <i>Urology</i> , 2012 , 79, 332-8 | 1.6 | 16 |
| 66 | Annual prostatectomy volume is related to rectal laceration rate after radical prostatectomy. <i>Urology</i> , 2012 , 79, 796-803 | 1.6 | 10 |
| 65 | Conditional survival of patients with urothelial carcinoma of the urinary bladder treated with radical cystectomy. <i>European Journal of Cancer</i> , 2012 , 48, 1503-11 | 7.5 | 39 |
| 64 | Propensity-score matched comparison of complications, blood transfusions, length of stay, and in-hospital mortality between open and laparoscopic partial nephrectomy: a national series. <i>European Journal of Surgical Oncology</i> , 2012 , 38, 80-7 | 3.6 | 27 |
| 63 | Predicting the risk of lymph node invasion during radical prostatectomy using the European Association of Urology guideline nomogram: a validation study. <i>European Journal of Surgical Oncology</i> , 2012 , 38, 624-9 | 3.6 | 6 |
| 62 | National Comprehensive Cancer Network practice guidelines 2011: Need for more accurate recommendations for pelvic lymph node dissection in prostate cancer. <i>Journal of Urology</i> , 2012 , 188, 423-8 | 2.5 | 19 |
| 61 | The extent of lymphadenectomy does affect cancer specific survival in pathologically confirmed T4 renal cell carcinoma. <i>Urologia</i> , 2012 , 79, 109-15 | 1.2 | 17 |
| 60 | Lymphatic spread of nodal metastases in high-risk prostate cancer: The ascending pathway from the pelvis to the retroperitoneum. <i>Prostate</i> , 2012 , 72, 186-92 | 4.2 | 58 |
| 59 | Assessing the risk of lymph node invasion in patients with intermediate risk prostate cancer treated with extended pelvic lymph node dissection. A novel prediction tool. <i>Prostate</i> , 2012 , 72, 499-506 | 4.2 | 10 |
| 58 | There is no way to identify patients who will harbor small volume, unilateral prostate cancer at final pathology. implications for focal therapies. <i>Prostate</i> , 2012 , 72, 925-30 | 4.2 | 11 |

| | | | |
|----|---|------|----|
| 57 | Development and validation of a reference table for prediction of postoperative mortality rate in patients treated with radical cystectomy: a population-based study. <i>Annals of Surgical Oncology</i> , 2012 , 19, 309-17 | 3.1 | 30 |
| 56 | The effect of annual surgical caseload on the rates of in-hospital pneumonia and other in-hospital outcomes after radical prostatectomy. <i>International Urology and Nephrology</i> , 2012 , 44, 799-806 | 2.3 | 7 |
| 55 | Lymph node count threshold for optimal pelvic lymph node staging in prostate cancer. <i>International Journal of Urology</i> , 2012 , 19, 645-51 | 2.3 | 45 |
| 54 | Optimizing postoperative sexual function after radical prostatectomy. <i>Therapeutic Advances in Urology</i> , 2012 , 4, 347-65 | 3.2 | 12 |
| 53 | Serum sex steroids depict a nonlinear u-shaped association with high-risk prostate cancer at radical prostatectomy. <i>Clinical Cancer Research</i> , 2012 , 18, 3648-57 | 12.9 | 53 |
| 52 | Prostate saturation biopsy following a first negative biopsy: state of the art. <i>Urologia Internationalis</i> , 2012 , 89, 126-35 | 1.9 | 20 |
| 51 | Impact of caseload on total hospital charges: a direct comparison between minimally invasive and open radical prostatectomy—a population based study. <i>Journal of Urology</i> , 2011 , 185, 855-61 | 2.5 | 22 |
| 50 | Tumor grade improves the prognostic ability of American Joint Committee on Cancer stage in patients with penile carcinoma. <i>Journal of Urology</i> , 2011 , 185, 501-7 | 2.5 | 26 |
| 49 | Impact of annual surgical volume on length of stay in patients undergoing minimally invasive prostatectomy: a population-based study. <i>European Journal of Surgical Oncology</i> , 2011 , 37, 429-34 | 3.6 | 7 |
| 48 | Perioperative mortality is significantly greater in septuagenarian and octogenarian patients treated with radical cystectomy for urothelial carcinoma of the bladder. <i>Urology</i> , 2011 , 77, 660-6 | 1.6 | 55 |
| 47 | Trans-rectal versus trans-perineal saturation rebiopsy of the prostate: is there a difference in cancer detection rate?. <i>Urology</i> , 2011 , 77, 921-5 | 1.6 | 73 |
| 46 | Head-to-head comparison of three commonly used preoperative tools for prediction of lymph node invasion at radical prostatectomy. <i>Urology</i> , 2011 , 78, 1363-7 | 1.6 | 15 |
| 45 | The importance of pelvic lymph node dissection in the elderly population: implications for interpreting the 2010 National Comprehensive Cancer Network practice guidelines for bladder cancer treatment. <i>Journal of Urology</i> , 2011 , 185, 2078-84 | 2.5 | 10 |
| 44 | Hospital and surgical caseload are predictors of comprehensive surgical treatment for bladder cancer: a population based study. <i>Journal of Urology</i> , 2011 , 186, 824-8 | 2.5 | 7 |
| 43 | Metabolic syndrome and benign prostatic hyperplasia: evidence of a potential relationship, hypothesized etiology, and prevention. <i>Korean Journal of Urology</i> , 2011 , 52, 507-16 | | 38 |
| 42 | How can we predict lymphorrhoea and clinically significant lymphocoeles after radical prostatectomy and pelvic lymphadenectomy? Clinical implications. <i>BJU International</i> , 2011 , 107, 1095-101 | 5.6 | 42 |
| 41 | The role of transrectal saturation biopsy in tumour localization: pathological correlation after retropubic radical prostatectomy and implication for focal ablative therapy. <i>BJU International</i> , 2011 , 108, 366-71 | 5.6 | 29 |
| 40 | The impact of surgical experience on total hospital charges for minimally invasive prostatectomy: a population-based study. <i>BJU International</i> , 2011 , 108, 888-93 | 5.6 | 7 |

| | | | |
|----|---|------|-----|
| 39 | Critical assessment of the European Association of Urology guideline indications for pelvic lymph node dissection at radical prostatectomy. <i>BJU International</i> , 2011 , 108, 1769-75 | 5.6 | 18 |
| 38 | Does diabetes mellitus increase the risk of high-grade prostate cancer in patients undergoing radical prostatectomy?. <i>Prostate Cancer and Prostatic Diseases</i> , 2011 , 14, 74-8 | 6.2 | 31 |
| 37 | A competing-risks analysis of survival after alternative treatment modalities for prostate cancer patients: 1988-2006. <i>European Urology</i> , 2011 , 59, 88-95 | 10.2 | 136 |
| 36 | Age-adjusted incidence, mortality, and survival rates of stage-specific renal cell carcinoma in North America: a trend analysis. <i>European Urology</i> , 2011 , 59, 135-41 | 10.2 | 214 |
| 35 | Reply to Charles B. Simone II, Charles B. Simone Letter to the Editor re: Firas Abdollah, Maxine Sun, Rodolphe Thuret, et al. A Competing-Risks Analysis of Survival After Alternative Treatment Modalities for Prostate Cancer Patients: 1988-2006. <i>Eur Urol</i> 2011;59:88-95. <i>European Urology</i> , 2011 , 59, e31-e32 | 10.2 | |
| 34 | A population-based analysis of temporal perioperative complication rates after minimally invasive radical prostatectomy. <i>European Urology</i> , 2011 , 60, 564-71 | 10.2 | 17 |
| 33 | Cancer-specific and other-cause mortality after radical prostatectomy versus observation in patients with prostate cancer: competing-risks analysis of a large North American population-based cohort. <i>European Urology</i> , 2011 , 60, 920-30 | 10.2 | 60 |
| 32 | The optimal rebiopsy prostatic scheme depends on patient clinical characteristics: results of a recursive partitioning analysis based on a 24-core systematic scheme. <i>European Urology</i> , 2011 , 60, 834-41 | 10.2 | 27 |
| 31 | Pelvic/retroperitoneal salvage lymph node dissection for patients treated with radical prostatectomy with biochemical recurrence and nodal recurrence detected by [11C]choline positron emission tomography/computed tomography. <i>European Urology</i> , 2011 , 60, 935-43 | 10.2 | 163 |
| 30 | A stage-for-stage and grade-for-grade analysis of cancer-specific mortality rates in renal cell carcinoma according to age: a competing-risks regression analysis. <i>European Urology</i> , 2011 , 60, 1152-9 | 10.2 | 28 |
| 29 | Impact of surgical experience on in-hospital complication rates in patients undergoing minimally invasive prostatectomy: a population-based study. <i>Annals of Surgical Oncology</i> , 2011 , 18, 839-47 | 3.1 | 20 |
| 28 | A contemporary population-based assessment of the rate of lymph node dissection for penile carcinoma. <i>Annals of Surgical Oncology</i> , 2011 , 18, 439-46 | 3.1 | 26 |
| 27 | Surgical caseload is an important determinant of continent urinary diversion rate at radical cystectomy: a population-based study. <i>Annals of Surgical Oncology</i> , 2011 , 18, 2680-7 | 3.1 | 10 |
| 26 | Mortality and morbidity after cytoreductive nephrectomy for metastatic renal cell carcinoma: a population-based study. <i>Annals of Surgical Oncology</i> , 2011 , 18, 2988-96 | 3.1 | 49 |
| 25 | Trends of retroperitoneal lymphadenectomy use in patients with nonseminomatous germ cell tumor of the testis: a population-based study. <i>Annals of Surgical Oncology</i> , 2011 , 18, 2997-3004 | 3.1 | 12 |
| 24 | The effect of marital status on stage and survival of prostate cancer patients treated with radical prostatectomy: a population-based study. <i>Cancer Causes and Control</i> , 2011 , 22, 1085-95 | 2.8 | 37 |
| 23 | A critical assessment of the value of lymph node dissection at radical prostatectomy: A population-based study. <i>Prostate</i> , 2011 , 71, 1587-94 | 4.2 | 18 |
| 22 | Racial disparities and socioeconomic status in men diagnosed with testicular germ cell tumors: a survival analysis. <i>Cancer</i> , 2011 , 117, 4277-85 | 6.4 | 61 |

| | | | |
|----|---|------|----|
| 21 | Conditional survival predictions after surgery for patients with penile carcinoma. <i>Cancer</i> , 2011 , 117, 3723-30 | 6.4 | 29 |
| 20 | Preoperative hypogonadism is not an independent predictor of high-risk disease in patients undergoing radical prostatectomy. <i>Cancer</i> , 2011 , 117, 3953-62 | 6.4 | 44 |
| 19 | Circulating estradiol, but not testosterone, is a significant predictor of high-grade prostate cancer in patients undergoing radical prostatectomy. <i>Cancer</i> , 2011 , 117, 5029-38 | 6.4 | 31 |
| 18 | Gleason 6 prostate cancer in one or two biopsy cores can harbor more aggressive disease. <i>Journal of Endourology</i> , 2011 , 25, 699-703 | 2.7 | 8 |
| 17 | Clinical experience and critical evaluation of the role of everolimus in advanced renal cell carcinoma. <i>Open Access Journal of Urology</i> , 2011 , 3, 43-8 | | |
| 16 | The importance of pelvic lymph node dissection in the elderly population: implications for interpreting the 2010 national comprehensive cancer network practice guidelines for bladder cancer treatment. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2011 , 37, 413-414 | 2 | |
| 15 | Should all men having a radical prostatectomy have a pelvic lymph node dissection? No. <i>Canadian Urological Association Journal</i> , 2010 , 4, 425-6 | 1.2 | 1 |
| 14 | Application of ice cold irrigation during vascular pedicle control of robot-assisted radical prostatectomy: EnSeal instrument cooling to reduce collateral thermal tissue damage. <i>Journal of Endourology</i> , 2010 , 24, 1991-6 | 2.7 | 22 |
| 13 | Annual surgical caseload and open radical prostatectomy outcomes: improving temporal trends. <i>Journal of Urology</i> , 2010 , 184, 2285-90 | 2.5 | 33 |
| 12 | Biopsy schemes with the fewest cores for detecting 95% of the prostate cancers detected by a 24-core biopsy. <i>European Urology</i> , 2010 , 57, 1-8 | 10.2 | 83 |
| 11 | Management of upper urinary tract urothelial carcinoma. <i>Expert Review of Anticancer Therapy</i> , 2010 , 10, 1955-65 | 3.5 | 7 |
| 10 | Decreasing rate and extent of lymph node staging in patients undergoing radical prostatectomy may undermine the rate of diagnosis of lymph node metastases in prostate cancer. <i>European Urology</i> , 2010 , 58, 882-92 | 10.2 | 75 |
| 9 | Surgery and erectile dysfunction. <i>Archivos Espanoles De Urologia</i> , 2010 , 63, 640-8 | 0.4 | 3 |
| 8 | Are infertile men less healthy than fertile men? Results of a prospective case-control survey. <i>European Urology</i> , 2009 , 56, 1025-31 | 10.2 | 94 |
| 7 | Obesity does not increase the risk of lymph node metastases in patients with clinically localized prostate cancer undergoing radical prostatectomy and extended pelvic lymph node dissection. <i>International Journal of Urology</i> , 2009 , 16, 676-81 | 2.3 | 8 |
| 6 | Prevention and Management of Postprostatectomy Erectile Dysfunction. <i>European Urology Supplements</i> , 2009 , 8, 80-87 | 0.9 | 3 |
| 5 | Is sperm banking of interest to patients with nongerm cell urological cancer before potentially fertility damaging treatments?. <i>Journal of Urology</i> , 2009 , 182, 1101-7 | 2.5 | 14 |
| 4 | VALIDATION OF THE CRITERIA SUGGESTED BY CURRENT GUIDELINES TO INDICATE THE NEED FOR BASELINE STAGING BONE SCAN IN PATIENTS WITH NEWLY DIAGNOSED PROSTATE CANCER. <i>Journal of Urology</i> , 2009 , 181, 782-782 | 2.5 | 3 |

| | | | |
|---|---|-----|----|
| 3 | Holmium laser enucleation of the prostate and holmium laser ablation of the prostate: indications and outcome. <i>Current Opinion in Urology</i> , 2009 , 19, 38-43 | 2.8 | 37 |
| 2 | Does educational status affect a patient's behavior toward erectile dysfunction?. <i>Journal of Sexual Medicine</i> , 2008 , 5, 1941-8 | 1.1 | 15 |
| 1 | Re: Orgasm associated incontinence (climacturia) following radical pelvic surgery: rates of occurrence and predictors. J. M. Choi, C. J. Nelson, J. Stasi and J. P. Mulhall. <i>J Urol</i> 2007; 177: 2223-2226. <i>Journal of Urology</i> , 2008 , 180, 1187; author reply 1187-8 | 2.5 | 1 |