

Yingwei Qiu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7802700/publications.pdf>

Version: 2024-02-01

24
papers

584
citations

566801

15
h-index

642321

23
g-index

25
all docs

25
docs citations

25
times ranked

871
citing authors

#	ARTICLE	IF	CITATIONS
1	Progressive Brain Structural Impairment Assessed via Network and Causal Analysis in Patients With Hepatitis B Virus-Related Cirrhosis. <i>Frontiers in Neurology</i> , 2022, 13, .	1.1	4
2	Divergent white matter changes in patients with nasopharyngeal carcinoma post-radiotherapy with different outcomes: a potential biomarker for prediction of radiation necrosis. <i>European Radiology</i> , 2022, 32, 7036-7047.	2.3	1
3	Longitudinal study of irradiation-induced brain functional network alterations in patients with nasopharyngeal carcinoma. <i>Radiotherapy and Oncology</i> , 2022, 173, 277-284.	0.3	5
4	Progressive Disruption of Dynamic Functional Network Connectivity in Patients With Hepatitis B Virus-related cirrhosis. <i>Journal of Magnetic Resonance Imaging</i> , 2021, 54, 1830-1840.	1.9	9
5	Divergent effects of irradiation on brain cortical morphology in patients with nasopharyngeal carcinoma: one-year follow-up study using structural magnetic resonance imaging. <i>Quantitative Imaging in Medicine and Surgery</i> , 2021, 11, 2307-2320.	1.1	1
6	Standard radiotherapy for patients with nasopharyngeal carcinoma results in progressive tract-specific brain white matter alterations: A one-year follow-up via diffusion tensor imaging. <i>Radiotherapy and Oncology</i> , 2021, 159, 255-264.	0.3	8
7	Prediction of Hepatocellular Carcinoma Response to Transcatheter Arterial Chemoembolization: A Real-World Study Based on Non-Contrast Computed Tomography Radiomics and General Image Features. <i>Journal of Hepatocellular Carcinoma</i> , 2021, Volume 8, 773-782.	1.8	20
8	Irradiation-related longitudinal white matter atrophy underlies cognitive impairment in patients with nasopharyngeal carcinoma. <i>Brain Imaging and Behavior</i> , 2021, 15, 2426-2435.	1.1	8
9	Aberrant inter-hemispheric coordination characterizes the progression of minimal hepatic encephalopathy in patients with HBV-related cirrhosis. <i>NeuroImage: Clinical</i> , 2020, 25, 102175.	1.4	13
10	Radiation-induced hippocampal atrophy in patients with nasopharyngeal carcinoma early after radiotherapy: a longitudinal MR-based hippocampal subfield analysis. <i>Brain Imaging and Behavior</i> , 2019, 13, 1160-1171.	1.1	35
11	Aberrant Brain Activity at Early Delay Stage Post-radiotherapy as a Biomarker for Predicting Neurocognitive Dysfunction Late-Delayed in Patients With Nasopharyngeal Carcinoma. <i>Frontiers in Neurology</i> , 2019, 10, 752.	1.1	19
12	Functional segregation loss over time is moderated by <i>APOE</i> genotype in healthy elderly. <i>Human Brain Mapping</i> , 2018, 39, 2742-2752.	1.9	16
13	Network-level dysconnectivity in patients with nasopharyngeal carcinoma (NPC) early post-radiotherapy: longitudinal resting state fMRI study. <i>Brain Imaging and Behavior</i> , 2018, 12, 1279-1289.	1.1	31
14	Longitudinal brain structural alterations in patients with nasopharyngeal carcinoma early after radiotherapy. <i>NeuroImage: Clinical</i> , 2018, 19, 252-259.	1.4	30
15	Aberrant interhemispheric functional and structural connectivity in heroin-dependent individuals. <i>Addiction Biology</i> , 2017, 22, 1057-1067.	1.4	30
16	Abnormal white matter structural networks characterize heroin-dependent individuals: a network analysis. <i>Addiction Biology</i> , 2016, 21, 667-678.	1.4	26
17	Inter-hemispheric functional dysconnectivity mediates the association of corpus callosum degeneration with memory impairment in AD and amnesic MCI. <i>Scientific Reports</i> , 2016, 6, 32573.	1.6	38
18	Aberrant Default-Mode Functional and Structural Connectivity in Heroin-Dependent Individuals. <i>PLoS ONE</i> , 2015, 10, e0120861.	1.1	40

#	ARTICLE	IF	CITATIONS
19	Structural and Functional Brain Alterations in End Stage Renal Disease Patients on Routine Hemodialysis: A Voxel-Based Morphometry and Resting State Functional Connectivity Study. PLoS ONE, 2014, 9, e98346.	1.1	55
20	Reduced ventral medial prefrontal cortex (vmPFC) volume and impaired vmPFC-default mode network integration in codeine-containing cough syrups users. Drug and Alcohol Dependence, 2014, 134, 314-321.	1.6	23
21	The impulsivity behavior is correlated with prefrontal cortex gray matter volume reduction in heroin-dependent individuals. Neuroscience Letters, 2013, 538, 43-48.	1.0	51
22	Progressive White Matter Microstructure Damage in Male Chronic Heroin Dependent Individuals: A DTI and TBSS Study. PLoS ONE, 2013, 8, e63212.	1.1	49
23	Reduced Regional Homogeneity in Bilateral Frontostriatal System Relates to Higher Impulsivity Behavior in Codeine-Containing Cough Syrups Dependent Individuals. PLoS ONE, 2013, 8, e78738.	1.1	25
24	Disrupted Topological Organization in Whole-Brain Functional Networks of Heroin-Dependent Individuals: A Resting-State fMRI Study. PLoS ONE, 2013, 8, e82715.	1.1	47