

Takahiro Shintani

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/780291/takahiro-shintani-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

54
papers

11,891
citations

23
h-index

55
g-index

55
ext. papers

13,054
ext. citations

5.2
avg. IF

5.65
L-index

#	Paper	IF	Citations
54	Improved recombinant protein production in <i>Aspergillus oryzae</i> lacking both β 1,3-glucan and galactosaminogalactan in batch culture with a lab-scale bioreactor. <i>Journal of Bioscience and Bioengineering</i> , 2021 ,	3.3	1
53	Enzymatic degradation of xyloglucans by <i>Aspergillus</i> species: a comparative view of this genus. <i>Applied Microbiology and Biotechnology</i> , 2021 , 105, 2701-2711	5.7	2
52	Crucial role of the intracellular β glucosidase MalT in the activation of the transcription factor AmyR essential for amyolytic gene expression in <i>Aspergillus oryzae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2021 , 85, 2076-2083	2.1	0
51	Expression profiles of amyolytic genes in AmyR and CreA transcription factor deletion mutants of the black koji mold <i>Aspergillus luchuensis</i> . <i>Journal of Bioscience and Bioengineering</i> , 2021 , 132, 321-326	3.3	0
50	Alternative transcription start sites of the enolase-encoding gene <i>enoA</i> are stringently used in glycolytic/gluconeogenic conditions in <i>Aspergillus oryzae</i> . <i>Current Genetics</i> , 2020 , 66, 729-747	2.9	0
49	Chaperone complex formation of the transcription factor MalR involved in maltose utilization and amyolytic enzyme production in. <i>Bioscience, Biotechnology and Biochemistry</i> , 2018 , 82, 827-835	2.1	2
48	Increased production of biomass-degrading enzymes by double deletion of <i>creA</i> and <i>creB</i> genes involved in carbon catabolite repression in <i>Aspergillus oryzae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2018 , 125, 141-147	3.3	12
47	The PDR-type ABC transporters <i>AtrA</i> and <i>AtrG</i> are involved in azole drug resistance in <i>Aspergillus oryzae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2018 , 82, 1840-1848	2.1	8
46	Nuclear export-dependent degradation of the carbon catabolite repressor <i>CreA</i> is regulated by a region located near the C-terminus in <i>Aspergillus oryzae</i> . <i>Molecular Microbiology</i> , 2018 , 110, 176-190	4.1	9
45	Subcellular localization of aphidicolin biosynthetic enzymes heterologously expressed in <i>Aspergillus oryzae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2018 , 82, 139-147	2.1	3
44	The C-terminal region of the yeast monocarboxylate transporter <i>Jen1</i> acts as a glucose signal-responding degron recognized by the β arrestin <i>Rod1</i> . <i>Journal of Biological Chemistry</i> , 2018 , 293, 10926-10936	5.4	13
43	Cellular responses to the expression of unstable secretory proteins in the filamentous fungus <i>Aspergillus oryzae</i> . <i>Applied Microbiology and Biotechnology</i> , 2017 , 101, 2437-2446	5.7	7
42	Improved β Amylase Production by Dephosphorylation Mutation of <i>CreD</i> , an Arrestin-Like Protein Required for Glucose-Induced Endocytosis of Maltose Permease and Carbon Catabolite Derepression in <i>Aspergillus oryzae</i> . <i>Applied and Environmental Microbiology</i> , 2017 , 83,	4.8	9
41	Quantitative regulation of histone variant H2A.Z during cell cycle by ubiquitin proteasome system and SUMO-targeted ubiquitin ligases. <i>Bioscience, Biotechnology and Biochemistry</i> , 2017 , 81, 1557-1560	2.1	5
40	Cell wall β 1,3-glucan prevents β amylase adsorption onto fungal cell in submerged culture of <i>Aspergillus oryzae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2017 , 124, 47-53	3.3	21
39	Self-excising <i>Cre</i> /mutant <i>lox</i> marker recycling system for multiple gene integrations and consecutive gene deletions in <i>Aspergillus oryzae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2017 , 123, 403-411	3.3	28
38	Induction of autophagy by phosphate starvation in an <i>Atg11</i> -dependent manner in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2017 , 483, 522-527	3.4	18

37	A Novel Zn ² -Cys ⁶ Transcription Factor AtrR Plays a Key Role in an Azole Resistance Mechanism of <i>Aspergillus fumigatus</i> by Co-regulating <i>cyp51A</i> and <i>cdr1B</i> Expressions. <i>PLoS Pathogens</i> , 2017 , 13, e1006096	7.6	68
36	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016 , 12, 1-222	10.2	3838
35	The C ₂ H ₂ -type transcription factor, FlbC, is involved in the transcriptional regulation of <i>Aspergillus oryzae</i> glucoamylase and protease genes specifically expressed in solid-state culture. <i>Applied Microbiology and Biotechnology</i> , 2016 , 100, 5859-68	5.7	13
34	Cargo Proteins Facilitate the Formation of Transport Vesicles, but not Autophagosomes 2016 , 143-154		
33	Endocytosis of a maltose permease is induced when amylolytic enzyme production is repressed in <i>Aspergillus oryzae</i> . <i>Fungal Genetics and Biology</i> , 2015 , 82, 136-44	3.9	19
32	Distinct mechanism of activation of two transcription factors, AmyR and MalR, involved in amylolytic enzyme production in <i>Aspergillus oryzae</i> . <i>Applied Microbiology and Biotechnology</i> , 2015 , 99, 1805-15	5.7	28
31	Purification and enzymatic characterization of secretory glycoside hydrolase family 3 (GH3) aryl β -glucosidases screened from <i>Aspergillus oryzae</i> genome. <i>Journal of Bioscience and Bioengineering</i> , 2015 , 120, 614-23	3.3	22
30	Unfolded protein response is required for <i>Aspergillus oryzae</i> growth under conditions inducing secretory hydrolytic enzyme production. <i>Fungal Genetics and Biology</i> , 2015 , 85, 1-6	3.9	13
29	Improved β -amylase production by <i>Aspergillus oryzae</i> after a double deletion of genes involved in carbon catabolite repression. <i>Applied Microbiology and Biotechnology</i> , 2014 , 98, 335-43	5.7	44
28	Evaluation of baker's yeast strains exhibiting significant growth on Japanese beet molasses and compound analysis of the molasses types. <i>Journal of Bioscience and Bioengineering</i> , 2014 , 117, 715-9	3.3	14
27	Fusion of an intact secretory protein permits a misfolded protein to exit from the endoplasmic reticulum in yeast. <i>Bioscience, Biotechnology and Biochemistry</i> , 2014 , 78, 49-59	2.1	1
26	Assays for autophagy I: the Cvt pathway and nonselective autophagy. <i>Methods in Molecular Biology</i> , 2014 , 1163, 153-64	1.4	14
25	Functional analysis of FarA transcription factor in the regulation of the genes encoding lipolytic enzymes and hydrophobic surface binding protein for the degradation of biodegradable plastics in <i>Aspergillus oryzae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2012 , 113, 549-55	3.3	17
24	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445-544.2	4.2	2783
23	Transcripts of a heterologous gene encoding mite allergen Der f 7 are stabilized by codon optimization in <i>Aspergillus oryzae</i> . <i>Applied Microbiology and Biotechnology</i> , 2012 , 96, 1275-82	5.7	20
22	Identification of potential cell wall component that allows Taka-amylase A adsorption in submerged cultures of <i>Aspergillus oryzae</i> . <i>Applied Microbiology and Biotechnology</i> , 2011 , 92, 961-9	5.7	24
21	Heterologous expression of <i>Aspergillus oryzae</i> xylose reductase and xylitol dehydrogenase genes facilitated xylose utilization in the yeast <i>Saccharomyces cerevisiae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2011 , 75, 168-70	2.1	3
20	Aspartyl aminopeptidase is imported from the cytoplasm to the vacuole by selective autophagy in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2011 , 286, 13704-13	5.4	64

19	In silico analysis of 3Tend-processing signals in <i>Aspergillus oryzae</i> using expressed sequence tags and genomic sequencing data. <i>DNA Research</i> , 2011 , 18, 189-200	4.5	9
18	The conserved oligomeric Golgi complex is involved in double-membrane vesicle formation during autophagy. <i>Journal of Cell Biology</i> , 2010 , 188, 101-14	7.3	157
17	Characterization and expression analysis of a maltose-utilizing (MAL) cluster in <i>Aspergillus oryzae</i> . <i>Fungal Genetics and Biology</i> , 2010 , 47, 1-9	3.9	32
16	Codon optimization increases steady-state mRNA levels in <i>Aspergillus oryzae</i> heterologous gene expression. <i>Applied and Environmental Microbiology</i> , 2008 , 74, 6538-46	4.8	49
15	Fluorescence microscopy-based assays for monitoring yeast Atg protein trafficking. <i>Methods in Enzymology</i> , 2008 , 451, 43-56	1.7	10
14	Atg11 directs autophagosome cargoes to the PAS along actin cables. <i>Autophagy</i> , 2006 , 2, 119-21	10.2	21
13	The actin cytoskeleton is required for selective types of autophagy, but not nonspecific autophagy, in the yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2005 , 16, 5843-56	3.5	127
12	Atg9 cycles between mitochondria and the pre-autophagosomal structure in yeasts. <i>Autophagy</i> , 2005 , 1, 101-9	10.2	207
11	Early stages of the secretory pathway, but not endosomes, are required for Cvt vesicle and autophagosome assembly in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2004 , 15, 2189-204	3.5	123
10	Cargo proteins facilitate the formation of transport vesicles in the cytoplasm to vacuole targeting pathway. <i>Journal of Biological Chemistry</i> , 2004 , 279, 29889-94	5.4	268
9	Autophagy in health and disease: a double-edged sword. <i>Science</i> , 2004 , 306, 990-5	33.3	2103
8	Vps51 is part of the yeast Vps fifty-three tethering complex essential for retrograde traffic from the early endosome and Cvt vesicle completion. <i>Journal of Biological Chemistry</i> , 2003 , 278, 5009-20	5.4	79
7	Mechanism of cargo selection in the cytoplasm to vacuole targeting pathway. <i>Developmental Cell</i> , 2002 , 3, 825-37	10.2	296
6	Apg2p functions in autophagosome formation on the perivacuolar structure. <i>Journal of Biological Chemistry</i> , 2001 , 276, 30452-60	5.4	95
5	Tor-mediated induction of autophagy via an Apg1 protein kinase complex. <i>Journal of Cell Biology</i> , 2000 , 150, 1507-13	7.3	909
4	Apg10p, a novel protein-conjugating enzyme essential for autophagy in yeast. <i>EMBO Journal</i> , 1999 , 18, 5234-41	13	216
3	Engineering of porcine pepsin. Alteration of S1 substrate specificity of pepsin to those of fungal aspartic proteinases by site-directed mutagenesis. <i>Journal of Biological Chemistry</i> , 1997 , 272, 18855-61	5.4	27
2	Characterization of the S1 subsite specificity of aspergillopepsin I by site-directed mutagenesis. <i>Journal of Biochemistry</i> , 1996 , 120, 974-81	3.1	14

- 1 Primary structure of aspergillopepsin I deduced from nucleotide sequence of the gene and aspartic acid-76 is an essential active site of the enzyme for trypsinogen activation. *BBA - Proteins and Proteomics*, **1994**, 1204, 257-64 26