

Riki Okita

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7823835/riki-okita-publications-by-year.pdf>

Version: 2024-04-11

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| | | | |
|-------------------|-----------------------|----------------|-----------------|
| 40 papers | 502 citations | 14 h-index | 22 g-index |
| 42 ext. papers | 609 ext. citations | 3.2 avg, IF | 3.55 L-index |

| # | Paper | IF | Citations |
|----|--|-----|-----------|
| 40 | Rapid development and rupture of a pneumatocele caused by pulmonary dissection in the early postoperative period of lung resection: a case report.. <i>AME Case Reports</i> , 2022 , 6, 6 | 0.6 | |
| 39 | Suspected fibrin glue-induced acute eosinophilic pneumonia after pulmonary resection: A case report. <i>Thoracic Cancer</i> , 2021 , 12, 2126-2129 | 3.2 | 3 |
| 38 | Treatment strategy for primary lung cancer in a lung highly compressed by giant emphysematous bullae: A case report. <i>Thoracic Cancer</i> , 2021 , 12, 268-271 | 3.2 | |
| 37 | Tofacitinib overcomes an IFN γ -induced decrease in NK cell-mediated cytotoxicity via the regulation of immune-related molecules in LC-2/ad. <i>Thoracic Cancer</i> , 2021 , 12, 775-782 | 3.2 | 1 |
| 36 | Tegafur-uracil-induced pericardial effusion during adjuvant chemotherapy for resected lung adenocarcinoma: A case report. <i>Thoracic Cancer</i> , 2021 , 12, 707-710 | 3.2 | 3 |
| 35 | Rapid development and rupture of a pulmonary cyst in the early postoperative period after pulmonary resection: A case report. <i>Thoracic Cancer</i> , 2020 , 11, 1712-1715 | 3.2 | 3 |
| 34 | Multilocular thymic cyst in a patient with preclinical rheumatoid arthritis: A case report. <i>International Journal of Surgery Case Reports</i> , 2020 , 74, 136-139 | 0.8 | 0 |
| 33 | Right upper lobectomy in lung cancer with double aortic arch: A case report. <i>Thoracic Cancer</i> , 2020 , 11, 2365-2369 | 3.2 | |
| 32 | Contralateral pulmonary resection using selective bronchial blockade in postpneumonectomy patients. <i>Thoracic Cancer</i> , 2020 , 11, 3528-3535 | 3.2 | |
| 31 | Comparative study of the PD-L1 expression and CD8+ tumor-infiltrating lymphocyte between surgically resected and matched re-biopsy specimens in recurrent non-small cell lung cancer. <i>Therapeutics and Clinical Risk Management</i> , 2019 , 15, 605-612 | 2.9 | 1 |
| 30 | Clinicopathological relevance of tumor expression of NK group 2 member D ligands in resected non-small cell lung cancer. <i>Oncotarget</i> , 2019 , 10, 6805-6815 | 3.3 | 3 |
| 29 | Effect of platinum-based chemotherapy on the expression of natural killer group 2 member D ligands, programmed cell death-1 ligand 1 and HLA class II in non-small cell lung cancer. <i>Oncology Reports</i> , 2019 , 42, 839-848 | 3.5 | 4 |
| 28 | Impact of COX2 Inhibitor for Regulation of PD-L1 Expression in Non-small Cell Lung Cancer. <i>Anticancer Research</i> , 2018 , 38, 4637-4644 | 2.3 | 19 |
| 27 | Deciduoid type malignant pleural mesothelioma: a case report. <i>AME Case Reports</i> , 2018 , 2, 43 | 0.6 | 2 |
| 26 | PD-L1 overexpression is partially regulated by EGFR/HER2 signaling and associated with poor prognosis in patients with non-small-cell lung cancer. <i>Cancer Immunology, Immunotherapy</i> , 2017 , 66, 865-876 | 7.4 | 94 |
| 25 | Urinary levels of prostaglandin E are positively correlated with intratumoral infiltration of Foxp3 regulatory T cells in non-small cell lung cancer. <i>Oncology Letters</i> , 2017 , 14, 1615-1620 | 2.6 | 3 |
| 24 | Prognostic value of Cox-2 and PD-L1 expression and its relationship with tumor-infiltrating lymphocytes in resected lung adenocarcinoma. <i>Cancer Management and Research</i> , 2017 , 9, 741-750 | 3.6 | 14 |

| | | | |
|----|---|-----|----|
| 23 | Clinicopathological and immunohistochemical features of lung invasive mucinous adenocarcinoma based on computed tomography findings. <i>OncoTargets and Therapy</i> , 2017 , 10, 153-163 | 4.4 | 19 |
| 22 | MHC class I chain-related molecule A and B expression is upregulated by cisplatin and associated with good prognosis in patients with non-small cell lung cancer. <i>Cancer Immunology, Immunotherapy</i> , 2016 , 65, 499-509 | 7.4 | 36 |
| 21 | Pneumatosis intestinalis after gefitinib therapy for pulmonary adenocarcinoma: a case report. <i>World Journal of Surgical Oncology</i> , 2016 , 14, 175 | 3.4 | 32 |
| 20 | Cyclooxygenase-2 expression is a prognostic biomarker for non-small cell lung cancer patients treated with adjuvant platinum-based chemotherapy. <i>World Journal of Surgical Oncology</i> , 2015 , 13, 21 | 3.4 | 16 |
| 19 | Clinical significance of dual-time-point 18F-FDG PET imaging in resectable non-small cell lung cancer. <i>Annals of Nuclear Medicine</i> , 2015 , 29, 854-60 | 2.5 | 22 |
| 18 | Preoperative neutrophil/lymphocyte ratio and prognostic nutritional index predict survival in patients with non-small cell lung cancer. <i>World Journal of Surgical Oncology</i> , 2015 , 13, 291 | 3.4 | 51 |
| 17 | Cyclooxygenase-2 genetic variants influence intratumoral infiltration of Foxp3-positive regulatory T cells in non-small cell lung cancer. <i>Oncology Reports</i> , 2015 , 33, 74-80 | 3.5 | 6 |
| 16 | Contrasting Effects of the Cytotoxic Anticancer Drug Gemcitabine and the EGFR Tyrosine Kinase Inhibitor Gefitinib on NK Cell-Mediated Cytotoxicity via Regulation of NKG2D Ligand in Non-Small-Cell Lung Cancer Cells. <i>PLoS ONE</i> , 2015 , 10, e0139809 | 3.7 | 21 |
| 15 | Prognostic nutritional index before adjuvant chemotherapy predicts chemotherapy compliance and survival among patients with non-small-cell lung cancer. <i>Therapeutics and Clinical Risk Management</i> , 2015 , 11, 1555-61 | 2.9 | 18 |
| 14 | Lung abscess presenting as tension pyopneumothorax in a gastrointestinal cancer patient. <i>Annals of Thoracic and Cardiovascular Surgery</i> , 2014 , 20 Suppl, 478-81 | 1.8 | 2 |
| 13 | Lung metastasis of adenoid cystic carcinoma, which mimicked primary lung cancer. <i>Thoracic Cancer</i> , 2013 , 4, 327-329 | 3.2 | 1 |
| 12 | Case of pleural metastasis of prostate cancer. <i>Thoracic Cancer</i> , 2013 , 4, 465-468 | 3.2 | 1 |
| 11 | Clinical significance of the tumor microenvironment in non-small cell lung cancer. <i>Annals of Translational Medicine</i> , 2013 , 1, 20 | 3.2 | 15 |
| 10 | Successful perioperative management for a breast cancer patient with Bernard-Soulier syndrome. <i>Breast Cancer</i> , 2012 , 19, 88-92 | 3.4 | 2 |
| 9 | HER2/HER3 signaling regulates NK cell-mediated cytotoxicity via MHC class I chain-related molecule A and B expression in human breast cancer cell lines. <i>Journal of Immunology</i> , 2012 , 188, 2136-45 | 5.3 | 41 |
| 8 | Opposing consequences of signaling through EGF family members: Escape from CTLs could be a bait for NK cells. <i>Onc Immunology</i> , 2012 , 1, 1200-1201 | 7.2 | 2 |
| 7 | T cell recognition of HLA-A2 restricted tumor antigens is impaired by the oncogene HER2. <i>International Journal of Cancer</i> , 2011 , 128, 390-401 | 7.5 | 42 |
| 6 | Suggestive synchronous triple squamous cell carcinoma of the lung in the same lobe. <i>General Thoracic and Cardiovascular Surgery</i> , 2010 , 58, 427-30 | 1.6 | 2 |

| | | | |
|---|---|-----|---|
| 5 | Use of entecavir to prevent hepatitis B virus reactivation during cytotoxic chemotherapy for solid malignancy. <i>Clinical Journal of Gastroenterology</i> , 2009 , 2, 214-217 | 1.1 | 6 |
| 4 | Targeting of CD4+CD25high cells while preserving CD4+CD25low cells with low-dose chimeric anti-CD25 antibody in adoptive immunotherapy of cancer. <i>International Journal of Oncology</i> , 2009 , 34, 563-72 | 1 | 9 |
| 3 | Functional inactivation of CD4+CD25high regulatory T cells using low dose human/mouse chimeric anti-CD25 monoclonal antibody enhanced lymphokine-activated killer cells activity. <i>Targeted Oncology</i> , 2008 , 3, 227-234 | 5 | |
| 2 | Enhancement of lymphokine-activated killer cell induction using anti-CD25 and anti-CTLA-4 monoclonal antibodies. <i>Oncology Reports</i> , 2007 , 17, 1429-35 | 3.5 | 2 |
| 1 | Intractable gastrointestinal bleeding from angiodysplasia in a patient of Bernard-Soulier syndrome--report of a case--. <i>Hiroshima Journal of Medical Sciences</i> , 2005 , 54, 113-5 | | 4 |