

Mario González-de la Parra

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7827068/publications.pdf>

Version: 2024-02-01

24
papers

171
citations

1162889

8
h-index

1281743

11
g-index

25
all docs

25
docs citations

25
times ranked

212
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Reference Datasets for Studies in a Replicate Design Intended for Average Bioequivalence with Expanding Limits. <i>AAPS Journal</i> , 2020, 22, 44. | 2.2 | 0 |
| 2 | Application of Design of Experiments (DOE) to the Development and Validation of a Swab Sampling Method for Cleaning Validation. <i>Asian Journal of Chemistry and Pharmaceutical Sciences</i> , 2017, 2, 16. | 0.0 | 0 |
| 3 | Application of Sequential Design of Experiments to Develop Ibuprofen (400 mg) Tablets by Direct Compression. <i>Asian Journal of Chemistry and Pharmaceutical Sciences</i> , 2017, 2, 10. | 0.0 | 2 |
| 4 | Quantification of 4-aminopyridine in plasma by capillary electrophoresis with electrokinetic injection. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2010, 878, 290-294. | 1.2 | 9 |
| 5 | Relative bioavailability of two oral formulations of piroxicam 20 mg: A single-dose, randomized-sequence, open-label, two-period crossover comparison in healthy Mexican adult volunteers. <i>Clinical Therapeutics</i> , 2010, 32, 357-364. | 1.1 | 12 |
| 6 | Bioavailability of two oral-tablet and two oral-suspension formulations of naproxen sodium/paracetamol (acetaminophen): Single-dose, randomized, open-label, two-period crossover comparisons in healthy Mexican adult subjects. <i>Clinical Therapeutics</i> , 2009, 31, 399-410. | 1.1 | 5 |
| 7 | Bioavailability of two oral formulations of a single dose of levofloxacin 500 mg: An open-label, randomized, two-period crossover comparison in healthy mexican volunteers. <i>Clinical Therapeutics</i> , 2009, 31, 1796-1803. | 1.1 | 12 |
| 8 | Bioavailability of Two Single-Dose Oral Formulations of Omeprazole 20 mg: An Open-Label, Randomized Sequence, Two-Period Crossover Comparison in Healthy Mexican Adult Volunteers. <i>Clinical Therapeutics</i> , 2008, 30, 693-699. | 1.1 | 9 |
| 9 | Bioavailability of two sublingual formulations of ketorolac tromethamine 30 mg: A randomized, open-label, single-dose, two-period crossover comparison in healthy mexican adult volunteers. <i>Clinical Therapeutics</i> , 2008, 30, 1667-1674. | 1.1 | 15 |
| 10 | Development and validation of a densitometric HPTLC method for quantitative analysis of levofloxacin in human plasma. <i>Journal of Planar Chromatography - Modern TLC</i> , 2008, 21, 209-212. | 0.6 | 6 |
| 11 | Development and Validation of a High-Performance Thin-Layer Chromatographic Method, with Densitometry, for Quantitative Analysis of Ketorolac Tromethamine in Human Plasma. <i>Journal of AOAC INTERNATIONAL</i> , 2008, 91, 1191-1195. | 0.7 | 7 |
| 12 | Bioequivalence of Two Commercial Preparations of trimethoprim/sulfamethoxazole: A randomized, single-dose, single-blind, crossover trial. <i>Clinical Therapeutics</i> , 2007, 29, 326-333. | 1.1 | 6 |
| 13 | Bioavailability of two oral suspension and two oral tablet formulations of acyclovir 400 mg: Two single-dose, open-label, randomized, two-period crossover comparisons in healthy Mexican adult subjects. <i>Clinical Therapeutics</i> , 2007, 29, 1146-1152. | 1.1 | 8 |
| 14 | Development and validation of a high-performance thin-layer chromatographic method, with densitometry, for quantitative analysis of tizoxanide (a Metabolite of Nitazoxanide) in human plasma. <i>Journal of Planar Chromatography - Modern TLC</i> , 2007, 20, 331-334. | 0.6 | 4 |
| 15 | Bioavailability of two oral formulations of loratadine 20 mg with concomitant ketoconazole: An open-label, randomized, two-period crossover comparison in healthy Mexican adult volunteers. <i>Clinical Therapeutics</i> , 2006, 28, 110-115. | 1.1 | 6 |
| 16 | Using Structural Equation Modeling (SEM) for the Study of Impurity Profiles of Drug Substances. <i>Quality Engineering</i> , 2006, 18, 225-235. | 0.7 | 6 |
| 17 | Bioavailability of two oral formulations of azithromycin 500 mg: A randomized, open-label, two-period crossover comparison in healthy Mexican adult subjects. <i>Clinical Therapeutics</i> , 2005, 27, 1607-1611. | 1.1 | 13 |
| 18 | Preliminary study on the synthesis and high-resolution NMR analysis of Naproxen and Ibuprofen esters. <i>Spectroscopy</i> , 2004, 18, 485-500. | 0.8 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Studying Unidentified Impurities in Drug Substances by the Application of Multivariate (Principal) Tj ETQq1 1 0.784314 rgBT 4/Overloc | 0.7 | 7 |
| 20 | Application of Analysis of Means (ANOM) to Nested Designs for Improving the Visualization and Understanding of the Sources of Variation of Chemical and Pharmaceutical Processes. Quality Engineering, 2003, 15, 663-670. | 0.7 | 7 |
| 21 | Application of the Multivariate T2Control Chart and the Masonâ€“Tracyâ€“Young Decomposition Procedure to the Study of the Consistency of Impurity Profiles of Drug Substances. Quality Engineering, 2003, 16, 127-142. | 0.7 | 14 |
| 22 | The Use of a Germination Bioassay as a Toxicological Screening System to Study the Potential Drugâ€“Drug Interactions of (S)-Naproxen and (S)-Ibuprofen and their corresponding Oxidation State Analogues, (S)-Naproxol and (S)-Ibuprofen Alcohol. ATLA Alternatives To Laboratory Animals, 1999, 27, 461-469. | 0.7 | 0 |
| 23 | Acaricidal potential of piquerols A and B againstBoophilus microplus. Pest Management Science, 1991, 33, 73-80. | 0.6 | 10 |
| 24 | Macrolide biosynthesis: Stereochemistry of the hydroxylation of brefelding C.. Journal of Antibiotics, 1987, 40, 1170-1174. | 1.0 | 8 |