

Byeong-Woo Park

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7828665/byeong-woo-park-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

131
papers

3,239
citations

29
h-index

54
g-index

139
ext. papers

3,870
ext. citations

4.7
avg, IF

4.78
L-index

#	Paper	IF	Citations
131	Adjuvant Capecitabine for Breast Cancer after Preoperative Chemotherapy. <i>New England Journal of Medicine</i> , 2017 , 376, 2147-2159	59.2	691
130	Up-regulation of acetyl-CoA carboxylase alpha and fatty acid synthase by human epidermal growth factor receptor 2 at the translational level in breast cancer cells. <i>Journal of Biological Chemistry</i> , 2007 , 282, 26122-31	5.4	158
129	Rationally designed anti-HER2/neu peptide mimetic disables P185HER2/neu tyrosine kinases in vitro and in vivo. <i>Nature Biotechnology</i> , 2000 , 18, 194-8	44.5	158
128	Characteristics and outcomes according to molecular subtypes of breast cancer as classified by a panel of four biomarkers using immunohistochemistry. <i>Breast</i> , 2012 , 21, 50-7	3.6	141
127	A genome-wide association study identifies a breast cancer risk variant in ERBB4 at 2q34: results from the Seoul Breast Cancer Study. <i>Breast Cancer Research</i> , 2012 , 14, R56	8.3	102
126	Basic Facts of Breast Cancer in Korea in 2014: The 10-Year Overall Survival Progress. <i>Journal of Breast Cancer</i> , 2017 , 20, 1-11	3	83
125	Benign papilloma without atypia diagnosed at US-guided 14-gauge core-needle biopsy: clinical and US features predictive of upgrade to malignancy. <i>Radiology</i> , 2011 , 258, 81-8	20.5	76
124	Krüppel-like factor 4 (KLF4) activates the transcription of the gene for the platelet isoform of phosphofructokinase (PFKP) in breast cancer. <i>Journal of Biological Chemistry</i> , 2011 , 286, 23808-16	5.4	68
123	Expression of estrogen receptor-beta in normal mammary and tumor tissues: is it protective in breast carcinogenesis?. <i>Breast Cancer Research and Treatment</i> , 2003 , 80, 79-85	4.4	53
122	Efficacy of everolimus with exemestane versus exemestane alone in Asian patients with HER2-negative, hormone-receptor-positive breast cancer in BOLERO-2. <i>Breast Cancer</i> , 2014 , 21, 703-14	3.4	52
121	Sentinel lymph node biopsy after neoadjuvant chemotherapy in patients with cytologically proven node-positive breast cancer at diagnosis. <i>Annals of Surgical Oncology</i> , 2013 , 20, 2858-65	3.1	51
120	Human leukocyte antigen-G (HLA-G) polymorphism and expression in breast cancer patients. <i>PLoS ONE</i> , 2014 , 9, e98284	3.7	48
119	Clinicopathologic features and outcomes of metaplastic breast carcinoma: comparison with invasive ductal carcinoma of the breast. <i>Yonsei Medical Journal</i> , 2010 , 51, 864-9	3	47
118	Markedly enhanced cytolysis by E1B-19kD-deleted oncolytic adenovirus in combination with cisplatin. <i>Human Gene Therapy</i> , 2006 , 17, 379-90	4.8	46
117	Feasibility of Charcoal Tattooing of Cytology-Proven Metastatic Axillary Lymph Node at Diagnosis and Sentinel Lymph Node Biopsy after Neoadjuvant Chemotherapy in Breast Cancer Patients. <i>Cancer Research and Treatment</i> , 2018 , 50, 801-812	5.2	45
116	Oncologic safety of breast-conserving surgery compared to mastectomy in patients receiving neoadjuvant chemotherapy for locally advanced breast cancer. <i>Journal of Surgical Oncology</i> , 2013 , 108, 531-6	2.8	45
115	How many sentinel lymph nodes are enough for accurate axillary staging in t1-2 breast cancer?. <i>Journal of Breast Cancer</i> , 2011 , 14, 296-300	3	45

114	Risk predictors of underestimation and the need for sentinel node biopsy in patients diagnosed with ductal carcinoma in situ by preoperative needle biopsy. <i>Journal of Surgical Oncology</i> , 2013 , 107, 388-92	2.8	44
113	Does chemotherapy really affect the quality of life of women with breast cancer?. <i>Journal of Breast Cancer</i> , 2013 , 16, 229-35	3	44
112	Nonmalignant papillary lesions of the breast at US-guided directional vacuum-assisted removal: a preliminary report. <i>European Radiology</i> , 2008 , 18, 1774-83	8	42
111	The impact of caveolin protein expression in tumor stroma on prognosis of breast cancer. <i>Tumor Biology</i> , 2011 , 32, 787-99	2.9	40
110	Unmet Needs and Their Relationship with Quality of Life among Women with Recurrent Breast Cancer. <i>Journal of Breast Cancer</i> , 2012 , 15, 454-61	3	38
109	US surveillance of regional lymph node recurrence after breast cancer surgery. <i>Radiology</i> , 2009 , 252, 673-81	20.5	38
108	Breast lesions with imaging-histologic discordance during US-guided 14G automated core biopsy: can the directional vacuum-assisted removal replace the surgical excision? Initial findings. <i>European Radiology</i> , 2007 , 17, 2376-83	8	36
107	Quality of Life Differences between Younger and Older Breast Cancer Patients. <i>Journal of Breast Cancer</i> , 2011 , 14, 112-8	3	35
106	Clinicopathological features of infiltrating lobular carcinomas comparing with infiltrating ductal carcinomas: a case control study. <i>World Journal of Surgical Oncology</i> , 2010 , 8, 34	3.4	34
105	The perceived care needs of breast cancer patients in Korea. <i>Yonsei Medical Journal</i> , 2006 , 47, 524-33	3	31
104	Elevated levels of serum tumor markers CA 15-3 and CEA are prognostic factors for diagnosis of metastatic breast cancers. <i>Breast Cancer Research and Treatment</i> , 2013 , 141, 477-84	4.4	30
103	Higher expression of androgen receptor is a significant predictor for better endocrine-responsiveness in estrogen receptor-positive breast cancers. <i>Breast Cancer Research and Treatment</i> , 2012 , 133, 311-20	4.4	30
102	Association between genetic polymorphisms of CYP2D6 and outcomes in breast cancer patients with tamoxifen treatment. <i>Journal of Korean Medical Science</i> , 2011 , 26, 1007-13	4.7	28
101	Adding Ovarian Suppression to Tamoxifen for Premenopausal Breast Cancer: A Randomized Phase III Trial. <i>Journal of Clinical Oncology</i> , 2020 , 38, 434-443	2.2	28
100	A nomogram for predicting underestimation of invasiveness in ductal carcinoma in situ diagnosed by preoperative needle biopsy. <i>Breast</i> , 2013 , 22, 869-73	3.6	27
99	Clinicopathological characteristics of mucinous carcinoma of the breast in Korea: comparison with invasive ductal carcinoma-not otherwise specified. <i>Journal of Korean Medical Science</i> , 2010 , 25, 361-8	4.7	27
98	Bilateral synchronous breast cancer in an Asian population: mammographic and sonographic characteristics, detection methods, and staging. <i>American Journal of Roentgenology</i> , 2008 , 190, 208-13	5.4	27
97	Serum 25-hydroxyvitamin D deficiency and increased risk of breast cancer among Korean women: a case-control study. <i>Breast Cancer Research and Treatment</i> , 2015 , 152, 147-154	4.4	26

96	Lack of either estrogen or progesterone receptor expression is associated with poor survival outcome among luminal A breast cancer subtype. <i>Annals of Surgical Oncology</i> , 2013 , 20, 1505-13	3.1	26
95	Long-term follow-up results for ultrasound-guided vacuum-assisted removal of benign palpable breast mass. <i>American Journal of Surgery</i> , 2010 , 199, 1-7	2.7	26
94	Clinicopathological characteristics of male breast cancer. <i>Yonsei Medical Journal</i> , 2008 , 49, 978-86	3	25
93	Clinical significance of progesterone receptor and HER2 status in estrogen receptor-positive, operable breast cancer with adjuvant tamoxifen. <i>Journal of Cancer Research and Clinical Oncology</i> , 2011 , 137, 1123-30	4.9	24
92	Characteristics and Survival of Breast Cancer Patients with Multiple Synchronous or Metachronous Primary Cancers. <i>Yonsei Medical Journal</i> , 2015 , 56, 1213-20	3	23
91	Cyclooxygenase-2 expression in proliferative Ki-67-positive breast cancers is associated with poor outcomes. <i>Breast Cancer Research and Treatment</i> , 2012 , 133, 741-51	4.4	23
90	The feasibility of endoscopy-assisted breast conservation surgery for patients with early breast cancer. <i>Journal of Breast Cancer</i> , 2011 , 14, 52-7	3	22
89	The changes of estrogen receptor-beta variants expression in breast carcinogenesis: Decrease of estrogen receptor-beta2 expression is the key event in breast cancer development. <i>Journal of Surgical Oncology</i> , 2006 , 93, 504-10	2.8	22
88	Region-specific changes in the immunoreactivity of SIRT1 expression in the central nervous system of SOD1(G93A) transgenic mice as an in vivo model of amyotrophic lateral sclerosis. <i>Brain Research</i> , 2012 , 1433, 20-8	3.7	21
87	Axillary lymph node metastasis: CA-15-3 and carcinoembryonic antigen concentrations in fine-needle aspirates for preoperative diagnosis in patients with breast cancer. <i>Radiology</i> , 2010 , 254, 691-7	20.5	20
86	Subcellular localization of Mdm2 expression and prognosis of breast cancer. <i>International Journal of Clinical Oncology</i> , 2014 , 19, 842-51	4.2	19
85	The impact of a focally positive resection margin on the local control in patients treated with breast-conserving therapy. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 600-8	2.8	19
84	Columnar cell lesions of the breast: mammographic and US features. <i>European Journal of Radiology</i> , 2006 , 60, 264-9	4.7	19
83	Increased sclerostin levels after further ablation of remnant estrogen by aromatase inhibitors. <i>Endocrinology and Metabolism</i> , 2015 , 30, 58-64	3.5	15
82	Role of sonography in the detection of contralateral metachronous breast cancer in an Asian population. <i>American Journal of Roentgenology</i> , 2008 , 190, 476-80	5.4	15
81	Do Recent Advances in Diagnostic and Therapeutic Procedures Negate the Benefit of Postmastectomy Radiotherapy in N1 Patients With a Low Risk of Locoregional Recurrence?. <i>Medicine (United States)</i> , 2015 , 94, e1259	1.8	14
80	Sonographic surveillance for the detection of contralateral metachronous breast cancer in an Asian population. <i>American Journal of Roentgenology</i> , 2009 , 192, 221-8	5.4	13
79	Overexpression of class III beta tubulin and amplified HER2 gene predict good response to paclitaxel and trastuzumab therapy. <i>PLoS ONE</i> , 2012 , 7, e45127	3.7	13

78	Breast cancers presenting luminal B subtype features show higher discordant human epidermal growth factor receptor 2 results between immunohistochemistry and fluorescence in situ hybridization. <i>Cancer</i> , 2012 , 118, 914-23	6.4	12
77	Anaplastic lymphoma kinase gene copy number gain in inflammatory breast cancer (IBC): prevalence, clinicopathologic features and prognostic implication. <i>PLoS ONE</i> , 2015 , 10, e0120320	3.7	12
76	Value of ultrasound for postoperative surveillance of asian patients with history of breast cancer surgery: a single-center study. <i>Annals of Surgical Oncology</i> , 2013 , 20, 3461-8	3.1	12
75	The impact of local and regional recurrence on distant metastasis and survival in patients treated with breast conservation therapy. <i>Journal of Breast Cancer</i> , 2011 , 14, 191-7	3	12
74	Magnetic Resonance Imaging after Completion of Neoadjuvant Chemotherapy Can Accurately Discriminate between No Residual Carcinoma and Residual Ductal Carcinoma In Situ in Patients with Triple-Negative Breast Cancer. <i>PLoS ONE</i> , 2016 , 11, e0149347	3.7	12
73	Can additional immunohistochemistry staining replace the surgical excision for the diagnosis of papillary breast lesions classified as benign on 14-gage core needle biopsy?. <i>Breast Cancer Research and Treatment</i> , 2013 , 137, 797-806	4.4	11
72	Evaluation with 3.0-T MR imaging: predicting the pathological response of triple-negative breast cancer treated with anthracycline and taxane neoadjuvant chemotherapy. <i>Acta Radiologica</i> , 2015 , 56, 1069-77	2	11
71	Factors influencing the outcome of breast cancer patients with 10 or more metastasized axillary lymph nodes. <i>International Journal of Clinical Oncology</i> , 2011 , 16, 473-81	4.2	11
70	Comparison of treatment outcome between breast-conservation surgery with radiation and total mastectomy without radiation in patients with one to three positive axillary lymph nodes. <i>International Journal of Radiation Oncology Biology Physics</i> , 2011 , 80, 1446-52	4	11
69	Depression and Coping in Breast Cancer Patients. <i>Journal of Breast Cancer</i> , 2009 , 12, 199	3	11
68	Is the frozen section examination for sentinel lymph node necessary in early breast cancer patients?. <i>Annals of Surgical Treatment and Research</i> , 2019 , 97, 49-57	2	11
67	Fine-needle aspirates CYFRA 21-1 is a useful tumor marker for detecting axillary lymph node metastasis in breast cancer patients. <i>PLoS ONE</i> , 2013 , 8, e57248	3.7	10
66	Imaging-histologic discordance at sonographically guided percutaneous biopsy of breast lesions. <i>European Journal of Radiology</i> , 2008 , 65, 163-9	4.7	10
65	Ductal eccrine carcinoma presenting as a Paget@ disease-like lesion of the breast. <i>Breast Journal</i> , 2001 , 7, 358-62	1.2	10
64	The Understanding of Terminal Cancer and Its Relationship with Attitudes toward End-of-Life Care Issues. <i>Medical Decision Making</i> , 2014 , 34, 720-30	2.5	9
63	The significance of c-erbB-2 overexpression and p53 expression in patients with axillary lymph node--negative breast cancer: a tissue microarray study. <i>International Journal of Surgical Pathology</i> , 2007 , 15, 98-109	1.2	9
62	Invasive papillary carcinoma of the breast presenting as post-traumatic recurrent hemorrhagic cysts. <i>Yonsei Medical Journal</i> , 2006 , 47, 575-7	3	9
61	Adding Ultrasound to the Evaluation of Patients with Pathologic Nipple Discharge to Diagnose Additional Breast Cancers: Preliminary Data. <i>Ultrasound in Medicine and Biology</i> , 2015 , 41, 2099-107	3.5	8

60	Adjuvant chemotherapy and survival among patients 70 years of age and younger with node-negative breast cancer and the 21-gene recurrence score of 26-30. <i>Breast Cancer Research</i> , 2019 , 21, 110	8.3	8
59	Clinical Significance of Age at the Time of Diagnosis among Young Breast Cancer Patients. <i>Journal of Breast Cancer</i> , 2011 , 14, 314-21	3	8
58	Risk Factors Associated with Discordant Ki-67 Levels between Preoperative Biopsy and Postoperative Surgical Specimens in Breast Cancers. <i>PLoS ONE</i> , 2016 , 11, e0151054	3.7	8
57	Infiltrating syringomatous adenoma presenting as microcalcification in the nipple on screening mammogram: case report and review of the literature of radiologic features. <i>Clinical Imaging</i> , 2010 , 34, 462-5	2.7	7
56	Prognostic value of ABO blood types in young patients with breast cancer; a nationwide study in Korean Breast Cancer Society. <i>Medical Oncology</i> , 2017 , 34, 118	3.7	6
55	S-1 combined with docetaxel following doxorubicin plus cyclophosphamide as neoadjuvant therapy in breast cancer: phase II trial. <i>BMC Cancer</i> , 2013 , 13, 583	4.8	6
54	Factors predictive of occult nipple-areolar complex involvement in patients with carcinoma in situ of the breast. <i>Journal of Surgical Oncology</i> , 2017 , 116, 1046-1055	2.8	6
53	Low-grade adenosquamous carcinoma of the breast with diverse expression patterns of myoepithelial cell markers on immunohistochemistry: a case study. <i>Korean Journal of Pathology</i> , 2014 , 48, 229-33		6
52	Comparison of immunohistochemical staining in breast papillary neoplasms of cytokeratin 5/6 and p63 in core needle biopsies and surgical excisions. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2012 , 20, 108-15	1.9	6
51	Treatment-planning CT scan for breast and chest-wall irradiation: how many unexpected abnormalities could we detect?. <i>Clinical Imaging</i> , 2008 , 32, 443-6	2.7	6
50	Impact of patient age on the outcome of primary breast carcinoma. <i>Journal of Surgical Oncology</i> , 2002 , 80, 12-8	2.8	6
49	Lack of both androgen receptor and forkhead box A1 (FOXA1) expression is a poor prognostic factor in estrogen receptor-positive breast cancers. <i>Oncotarget</i> , 2017 , 8, 82940-82955	3.3	6
48	Assessment of the Prognostic Staging System of American Joint Committee on Cancer 8th Edition for Breast Cancer: Comparisons with the Conventional Anatomic Staging System. <i>Journal of Breast Cancer</i> , 2020 , 23, 59-68	3	6
47	Validation of Clinical Treatment Score post-5 years (CTS5) risk stratification in premenopausal breast cancer patients and Ki-67 labelling index. <i>Scientific Reports</i> , 2020 , 10, 16850	4.9	6
46	Survival of Breast-Conserving Surgery Plus Radiotherapy versus Total Mastectomy in Early Breast Cancer. <i>Annals of Surgical Oncology</i> , 2021 , 28, 5039-5047	3.1	6
45	Association between Changes in Serum 25-Hydroxyvitamin D Levels and Survival in Patients with Breast Cancer Receiving Neoadjuvant Chemotherapy. <i>Journal of Breast Cancer</i> , 2018 , 21, 134-141	3	5
44	Prolonged clinical benefit from the maintenance hormone therapy in patients with metastatic breast cancer. <i>Breast</i> , 2013 , 22, 1205-9	3.6	5
43	Comparison of characteristics in patients with both thyroid and breast cancer: Based on order of incidence. <i>Korean Journal of Clinical Oncology</i> , 2017 , 13, 1-9	0.1	5

42	Optimal treatment of pseudoangiomatous stromal hyperplasia of the breast. <i>Asian Journal of Surgery</i> , 2020 , 43, 735-741	1.6	5
41	Breast ultrasonography for detection of metachronous ipsilateral breast tumor recurrence. <i>Acta Radiologica</i> , 2016 , 57, 1171-7	2	4
40	Impact of Micrometastatic Axillary Nodes on Survival of Breast Cancer Patients with Tumors ≥ 1 cm. <i>World Journal of Surgery</i> , 2018 , 42, 3969-3978	3.3	4
39	Ki-67 and breast cancer prognosis: does it matter if Ki-67 level is examined using preoperative biopsy or postoperative specimen?. <i>Breast Cancer Research and Treatment</i> , 2022 , 192, 343	4.4	4
38	Prevalence of breast cancer-related risk factors in underweight premenopausal women: the Korea National Health and Nutrition Examination Survey IV-VI. <i>Breast Cancer Research and Treatment</i> , 2019 , 174, 515-524	4.4	4
37	Homeodomain-interacting protein kinase 1 (HIPK1) expression in breast cancer tissues. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 1138-45	2.8	3
36	Positive predictive value of additional synchronous breast lesions in whole-breast ultrasonography at the diagnosis of breast cancer: clinical and imaging factors. <i>Ultrasonography</i> , 2014 , 33, 170-7	4.3	3
35	Neoadjuvant Chemotherapy and Prognosis of Pregnancy-Associated Breast Cancer: A Time-Trends Study of the Korean Breast Cancer Registry Database. <i>Journal of Breast Cancer</i> , 2018 , 21, 425-432	3	3
34	Can Biannual Ultrasound Surveillance Detect Smaller Second Cancers or Detect Cancers Earlier in Patients with Breast Cancer History?. <i>Ultrasound in Medicine and Biology</i> , 2018 , 44, 1355-1363	3.5	2
33	Dermatofibrosarcoma protuberans arising on the skin of the breast. <i>Breast Journal</i> , 2011 , 17, 93-5	1.2	2
32	Adjuvant Hormonal Therapy: Current Standard and Practical Issues. <i>Journal of Breast Cancer</i> , 2010 , 13, 242	3	2
31	Breast cancer from the excisional scar of a benign mass. <i>Korean Journal of Radiology</i> , 2007 , 8, 254-7	6.9	2
30	The role of sonography in patients with breast cancer presenting as an axillary mass. <i>Korean Journal of Radiology</i> , 2002 , 3, 189-93	6.9	2
29	Clinical Outcome of Immediate Breast Reconstruction after Mastectomy for Breast Cancer. <i>Journal of Breast Disease</i> , 2014 , 2, 44-50	0.3	2
28	The Result of Evaluation According to Radioactivity of Sequential Sentinel Nodes Biopsy in Breast cancer. <i>Journal of Breast Cancer</i> , 2006 , 9, 235	3	2
27	The Prognostic Significance of Survivin Expression in Breast Cancer. <i>Journal of Breast Cancer</i> , 2009 , 12, 285	3	2
26	An Updated Nomogram for Predicting Invasiveness in Preoperative Ductal Carcinoma In Situ of the Breast. <i>Yonsei Medical Journal</i> , 2019 , 60, 1028-1035	3	2
25	Conserving the lymphatics from the arm using fluorescence imaging in patients with breast cancer at high risk of postoperative lymphedema: a pilot study. <i>Gland Surgery</i> , 2020 , 9, 629-636	2.2	2

24	Time-Sequencing of the Neutrophil-to-Lymphocyte Ratio to Predict Prognosis of Triple-Negative Breast Cancer. <i>Cancers</i> , 2021 , 13,	6.6	2
23	Expression of T-Lymphocyte Markers in Human Epidermal Growth Factor Receptor 2-Positive Breast Cancer. <i>Journal of Breast Cancer</i> , 2016 , 19, 385-393	3	2
22	Survival Rates of Breast Cancer Patients Aged 40 to 49 Years according to Detection Modality in Korea: Screening Ultrasound versus Mammography. <i>Korean Journal of Radiology</i> , 2021 , 22, 159-167	6.9	2
21	Axillary response according to neoadjuvant single or dual human epidermal growth factor receptor 2 (HER2) blockade in clinically node-positive, HER2-positive breast cancer. <i>International Journal of Cancer</i> , 2021 , 149, 1585-1592	7.5	2
20	Assessment of Quality of Life and Safety in Postmenopausal Breast Cancer Patients Receiving Letrozole as an Early Adjuvant Treatment. <i>Journal of Breast Cancer</i> , 2018 , 21, 182-189	3	1
19	Can We Predict Phyllodes Tumor among Fibroepithelial Lesions with Cellular Stroma Diagnosed at Breast Core Needle Biopsy?. <i>Journal of the Korean Society of Radiology</i> , 2011 , 64, 603	0.2	1
18	Omission of axillary lymph node dissection in patients who underwent total mastectomy with 1 or 2 metastatic lymph nodes. <i>Annals of Surgical Treatment and Research</i> , 2020 , 98, 283-290	2	1
17	Expression of growth factor receptor family before and after targeted therapy in human epidermal growth factor receptor-2 positive breast cancer tissues. <i>Korean Journal of Clinical Oncology</i> , 2015 , 11, 12-19	0.1	1
16	Five-year changes in ovarian function restoration in premenopausal patients with breast cancer taking tamoxifen after chemotherapy: An ASTRRA study report. <i>European Journal of Cancer</i> , 2021 , 151, 190-200	7.5	1
15	Safety and efficacy of everolimus (EVE) plus exemestane (EXE) in postmenopausal women with locally advanced or metastatic breast cancer: final results from EVEREXES. <i>Breast Cancer Research and Treatment</i> , 2021 , 188, 77-89	4.4	0
14	Solitary drain-site recurrence after lumpectomy for breast cancer. <i>Yonsei Medical Journal</i> , 2010 , 51, 469-31		
13	Paclitaxel combined with ifosfamide in anthracycline- and docetaxel-pretreated metastatic breast cancer: activity independence of prior docetaxel resistance. <i>Cancer Chemotherapy and Pharmacology</i> , 2010 , 66, 425-31	3.5	
12	Changes of Coregulators, MAP Kinase Activity and p27/kip1 with Estrogen or Antiestrogen Treatment in Breast Cancer Cell Line. <i>Journal of Breast Cancer</i> , 2008 , 11, 56	3	
11	Clinicopathological Characteristics and Prognostic Factors of Papillary Carcinoma of the Breast. <i>Journal of Breast Cancer</i> , 2008 , 11, 77	3	
10	Molecular Targets for Treatment of Breast Cancer. <i>Journal of Breast Cancer</i> , 2009 , 12, 229	3	
9	KLF5 enhances SREBP-1 action in androgen-dependent induction of FASN in prostate cancer cell line. <i>FASEB Journal</i> , 2008 , 22, 794.12	0.9	
8	A multicenter phase II neoadjuvant trial of bevacizumab combined with docetaxel plus carboplatin in the treatment of triple-negative breast cancer: Korean Cancer Study Group (KCSG-BR 0905, NCT 01208480).. <i>Journal of Clinical Oncology</i> , 2012 , 30, 2542-2542	2.2	
7	On-treatment derived neutrophil-to-lymphocyte ratio and response to palbociclib and letrozole: Analysis of a multicenter retrospective cohort and the PALOMA-2 study.. <i>Journal of Clinical Oncology</i> , 2021 , 39, 1066-1066	2.2	

6	Clinicopathological Features and Prognosis of Lobular Carcinoma In Situ. <i>Journal of Breast Disease</i> , 2021 , 9, 10-15	0.3
5	Abstract P2-11-13: Distinctive clinicopathological features of premenopausal women with intermediate 21-gene recurrence score. <i>Cancer Research</i> , 2022 , 82, P2-11-13-P2-11-13	10.1
4	Abstract P2-08-13: The prognostic significance of Ki67 and progesterone receptor; could they really have prognostic importance in early breast cancer with estrogen receptor positive, HER-2 negative?. <i>Cancer Research</i> , 2022 , 82, P2-08-13-P2-08-13	10.1
3	Abstract P4-07-14: Long-term prognosis of minimal residual disease in breast cancer patients with breast or nodal pathologic complete response (pCR) after neoadjuvant chemotherapy (NAC). <i>Cancer Research</i> , 2022 , 82, P4-07-14-P4-07-14	10.1
2	Abstract P1-07-03: Clinical treatment score post-5 years (CTS5) and the prediction of survival benefit from extended endocrine therapy for breast cancer patients under and over 50 years of age. <i>Cancer Research</i> , 2022 , 82, P1-07-03-P1-07-03	10.1
1	Abstract P2-10-07: Prediction of menstruation recovery timing in premenopausal breast cancer patients taking tamoxifen after chemotherapy: An ASTRRA substudy. <i>Cancer Research</i> , 2022 , 82, P2-10-07-P2-10-07	10.1