

Johan Reutfors

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7837801/publications.pdf>

Version: 2024-02-01

53
papers

1,574
citations

393982

19
h-index

329751

37
g-index

54
all docs

54
docs citations

54
times ranked

2420
citing authors

#	ARTICLE	IF	CITATIONS
1	Risks of adverse pregnancy and birth outcomes in women treated or not treated with mood stabilisers for bipolar disorder: population based cohort study. <i>BMJ</i> , The, 2012, 345, e7085-e7085.	3.0	185
2	International trends in antipsychotic use: A study in 16 countries, 2005–2014. <i>European Neuropsychopharmacology</i> , 2017, 27, 1064-1076.	0.3	185
3	Antipsychotics During Pregnancy. <i>Archives of General Psychiatry</i> , 2012, 69, 715-21.	13.8	111
4	Association Between Methylphenidate and Amphetamine Use in Pregnancy and Risk of Congenital Malformations. <i>JAMA Psychiatry</i> , 2018, 75, 167.	6.0	93
5	Early non-adherence to medication and other risk factors for rehospitalization in schizophrenia and schizoaffective disorder. <i>Schizophrenia Research</i> , 2011, 133, 36-41.	1.1	72
6	Risk factors for suicide in schizophrenia: Findings from a Swedish population-based case-control study. <i>Schizophrenia Research</i> , 2009, 108, 231-237.	1.1	70
7	Mortality in treatment-resistant unipolar depression: A register-based cohort study in Sweden. <i>Journal of Affective Disorders</i> , 2018, 238, 674-679.	2.0	64
8	Suicide and hospitalization for mental disorders in Sweden: A population-based case-control study. <i>Journal of Psychiatric Research</i> , 2010, 44, 741-747.	1.5	48
9	Antidepressant use in Denmark, Germany, Spain, and Sweden between 2009 and 2014: Incidence and comorbidities of antidepressant initiators. <i>Journal of Affective Disorders</i> , 2019, 249, 242-252.	2.0	47
10	Seasonality of suicide in Sweden: Relationship with psychiatric disorder. <i>Journal of Affective Disorders</i> , 2009, 119, 59-65.	2.0	45
11	All-cause mortality in patients with treatment-resistant depression: a cohort study in the US population. <i>Annals of General Psychiatry</i> , 2019, 18, 23.	1.2	44
12	Trends in utilization and dosing of antipsychotic drugs in Scandinavia: Comparison of 2006 and 2016. <i>British Journal of Clinical Pharmacology</i> , 2019, 85, 1598-1606.	1.1	38
13	Medication and suicide risk in schizophrenia: A nested case-control study. <i>Schizophrenia Research</i> , 2013, 150, 416-420.	1.1	37
14	Serotonergic medication enhances the association between suicide and sunshine. <i>Journal of Affective Disorders</i> , 2016, 189, 276-281.	2.0	35
15	Humanistic outcomes in treatment resistant depression: a secondary analysis of the STAR*D study. <i>BMC Psychiatry</i> , 2018, 18, 352.	1.1	34
16	Risk Factors for Suicide and Suicide Attempts Among Patients With Treatment-Resistant Depression: Nested Case-Control Study. <i>Archives of Suicide Research</i> , 2021, 25, 424-438.	1.2	31
17	Antipsychotic drug use in pregnancy: A multinational study from ten countries. <i>Schizophrenia Research</i> , 2020, 220, 106-115.	1.1	30
18	Risk of Acute Liver Injury in Agomelatine and Other Antidepressant Users in Four European Countries: A Cohort and Nested Case-Control Study Using Automated Health Data Sources. <i>CNS Drugs</i> , 2019, 33, 383-395.	2.7	23

#	ARTICLE	IF	CITATIONS
19	Prevalence trends and individual patterns of antiepileptic drug use in pregnancy 2006–2016: A study in the five Nordic countries, United States, and Australia. <i>Pharmacoepidemiology and Drug Safety</i> , 2020, 29, 913-922.	0.9	23
20	Initiation and long-term use of benzodiazepines and Z-drugs in bipolar disorder. <i>Bipolar Disorders</i> , 2018, 20, 634-646.	1.1	22
21	Treatment-resistant depression as risk factor for substance use disorders—a nationwide register-based cohort study. <i>Addiction</i> , 2019, 114, 1274-1282.	1.7	21
22	Substance use disorders and risk for treatment resistant depression: a population-based, nested case-control study. <i>Addiction</i> , 2020, 115, 768-777.	1.7	20
23	Risk of breast cancer in risperidone users: A nationwide cohort study. <i>Schizophrenia Research</i> , 2017, 182, 98-103.	1.1	19
24	Use of hypnotic drugs among children, adolescents, and young adults in Scandinavia. <i>Acta Psychiatrica Scandinavica</i> , 2021, 144, 100-112.	2.2	17
25	Monotherapy vs. combination therapy for post mania maintenance treatment: A population based cohort study. <i>European Neuropsychopharmacology</i> , 2019, 29, 691-700.	0.3	16
26	Reducing the rehospitalization risk after a manic episode: A population based cohort study of lithium, valproate, olanzapine, quetiapine and aripiprazole in monotherapy and combinations. <i>Journal of Affective Disorders</i> , 2017, 217, 16-23.	2.0	15
27	A register-based approach to identifying treatment-resistant depression—Comparison with clinical definitions. <i>PLoS ONE</i> , 2020, 15, e0236434.	1.1	15
28	Suicide risk and antipsychotic side effects in schizophrenia: nested case-control study. <i>Human Psychopharmacology</i> , 2016, 31, 341-345.	0.7	14
29	Diagnostic profile and suicide risk in schizophrenia spectrum disorder. <i>Schizophrenia Research</i> , 2010, 123, 251-256.	1.1	13
30	Suicide and all-cause mortality in Swedish deployed military veterans: a population-based matched cohort study. <i>BMJ Open</i> , 2017, 7, e014034.	0.8	13
31	Antipsychotic use in pregnancy and risk of attention/deficit-hyperactivity disorder and autism spectrum disorder: a Nordic cohort study. <i>Evidence-Based Mental Health</i> , 2022, 25, 54-62.	2.2	13
32	Long-Term Risk of Skin Cancer and Lymphoma in Users of Topical Tacrolimus and Pimecrolimus: Final Results from the Extension of the Cohort Study Protopic Joint European Longitudinal Lymphoma and Skin Cancer Evaluation (JOELLE). <i>Clinical Epidemiology</i> , 2021, Volume 13, 1141-1153.	1.5	13
33	Central nervous system tumours in children in Costa Rica, 1981-96. <i>Paediatric and Perinatal Epidemiology</i> , 2002, 16, 219-225.	0.8	12
34	Suicide Ideation and Behavior as Risk Factors for Subsequent Suicide in Schizophrenia: A Nested Case-Control Study. <i>Suicide and Life-Threatening Behavior</i> , 2019, 49, 996-1005.	0.9	12
35	Risk and predictors of attempted and completed suicide in patients with hematological malignancies. <i>Psycho-Oncology</i> , 2014, 23, 1276-1282.	1.0	11
36	Potentially addictive drugs dispensing to patients receiving opioid agonist therapy: a register-based prospective cohort study in Norway and Sweden from 2015 to 2017. <i>BMJ Open</i> , 2020, 10, e036860.	0.8	11

#	ARTICLE	IF	CITATIONS
37	A retrospective analysis to estimate the healthcare resource utilization and cost associated with treatment-resistant depression in commercially insured US patients. PLoS ONE, 2020, 15, e0238843.	1.1	11
38	Risk and risk factors for disability pension among patients with treatment resistant depression – a matched cohort study. BMC Psychiatry, 2020, 20, 232.	1.1	11
39	Predictors for initiation of pharmacological prophylaxis in patients with newly diagnosed bipolar disorder – A nationwide cohort study. Journal of Affective Disorders, 2015, 172, 204-210.	2.0	10
40	Marriage and divorce after military deployment to Afghanistan: A matched cohort study from Sweden. PLoS ONE, 2019, 14, e0207981.	1.1	10
41	Response rate and subjective memory after electroconvulsive therapy in depressive disorders with psychiatric comorbidity. Journal of Affective Disorders, 2021, 292, 276-283.	2.0	10
42	Antipsychotic Prescription Filling in Patients With Schizophrenia or Schizoaffective Disorder. Journal of Clinical Psychopharmacology, 2013, 33, 759-765.	0.7	8
43	<p>Maternal bereavement shortly before or during pregnancy and risk of postpartum psychotic illness: a population-based study from Denmark and Sweden</p>. Clinical Epidemiology, 2019, Volume 11, 285-298.	1.5	7
44	Diabetes and glucose disturbances in patients with psychosis in Sweden. BMJ Open Diabetes Research and Care, 2015, 3, e000120.	1.2	6
45	Excess deaths in treatment-resistant depression. Therapeutic Advances in Psychopharmacology, 2021, 11, 204512532110065.	1.2	6
46	A Post-Authorization Safety Study of Quetiapine as Antidepressant Treatment in Sweden: Nested Case – Control Analyses of Select Outcomes. Drug Safety, 2020, 43, 135-145.	1.4	5
47	Risk of depression in multiple sclerosis across disease-modifying therapies. Multiple Sclerosis Journal, 2022, 28, 632-641.	1.4	5
48	Violent crime among Swedish military veterans after deployment to Afghanistan: a population-based matched cohort study. International Journal of Epidemiology, 2019, 48, 1604-1613.	0.9	3
49	Sunshine, temperature and suicidal behaviour in patients treated with antidepressants: an explorative nested case – control study. Scientific Reports, 2021, 11, 10178.	1.6	3
50	Health care utilisation in treatment-resistant depression: a Swedish population-based cohort study. International Journal of Psychiatry in Clinical Practice, 2022, 26, 251-258.	1.2	3
51	Trends in use of attention deficit hyperactivity disorder medication among children and adolescents in Scandinavia in 2010 – 2020. European Child and Adolescent Psychiatry, 0, , .	2.8	2
52	Season of treatment initiation with antidepressants and suicidal behavior: A population-based cohort study in Sweden. Journal of Affective Disorders, 2017, 215, 245-255.	2.0	1
53	The use of uncertain exposure – A method to define switching and add – on in pharmacoepidemiology. Pharmacoepidemiology and Drug Safety, 2021, , .	0.9	1