

# Johan Reutfors

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7837801/publications.pdf>

Version: 2024-02-01

53  
papers

1,574  
citations

393982

19  
h-index

329751

37  
g-index

54  
all docs

54  
docs citations

54  
times ranked

2420  
citing authors

#	ARTICLE	IF	CITATIONS
1	Risk of depression in multiple sclerosis across disease-modifying therapies. <i>Multiple Sclerosis Journal</i> , 2022, 28, 632-641.	1.4	5
2	Antipsychotic use in pregnancy and risk of attention/deficit-hyperactivity disorder and autism spectrum disorder: a Nordic cohort study. <i>Evidence-Based Mental Health</i> , 2022, 25, 54-62.	2.2	13
3	Health care utilisation in treatment-resistant depression: a Swedish population-based cohort study. <i>International Journal of Psychiatry in Clinical Practice</i> , 2022, 26, 251-258.	1.2	3
4	Risk Factors for Suicide and Suicide Attempts Among Patients With Treatment-Resistant Depression: Nested Case-Control Study. <i>Archives of Suicide Research</i> , 2021, 25, 424-438.	1.2	31
5	Excess deaths in treatment-resistant depression. <i>Therapeutic Advances in Psychopharmacology</i> , 2021, 11, 204512532110065.	1.2	6
6	Sunshine, temperature and suicidal behaviour in patients treated with antidepressants: an explorative nested case-control study. <i>Scientific Reports</i> , 2021, 11, 10178.	1.6	3
7	Use of hypnotic drugs among children, adolescents, and young adults in Scandinavia. <i>Acta Psychiatrica Scandinavica</i> , 2021, 144, 100-112.	2.2	17
8	Response rate and subjective memory after electroconvulsive therapy in depressive disorders with psychiatric comorbidity. <i>Journal of Affective Disorders</i> , 2021, 292, 276-283.	2.0	10
9	The use of uncertain exposure—A method to define switching and add-on in pharmacoepidemiology. <i>Pharmacoepidemiology and Drug Safety</i> , 2021, , .	0.9	1
10	Long-Term Risk of Skin Cancer and Lymphoma in Users of Topical Tacrolimus and Pimecrolimus: Final Results from the Extension of the Cohort Study Protopic Joint European Longitudinal Lymphoma and Skin Cancer Evaluation (JOELLE). <i>Clinical Epidemiology</i> , 2021, Volume 13, 1141-1153.	1.5	13
11	Substance use disorders and risk for treatment resistant depression: a population-based, nested case-control study. <i>Addiction</i> , 2020, 115, 768-777.	1.7	20
12	A Post-Authorization Safety Study of Quetiapine as Antidepressant Treatment in Sweden: Nested Case-Control Analyses of Select Outcomes. <i>Drug Safety</i> , 2020, 43, 135-145.	1.4	5
13	Potentially addictive drugs dispensing to patients receiving opioid agonist therapy: a register-based prospective cohort study in Norway and Sweden from 2015 to 2017. <i>BMJ Open</i> , 2020, 10, e036860.	0.8	11
14	A register-based approach to identifying treatment-resistant depression—Comparison with clinical definitions. <i>PLoS ONE</i> , 2020, 15, e0236434.	1.1	15
15	A retrospective analysis to estimate the healthcare resource utilization and cost associated with treatment-resistant depression in commercially insured US patients. <i>PLoS ONE</i> , 2020, 15, e0238843.	1.1	11
16	Risk and risk factors for disability pension among patients with treatment resistant depression—a matched cohort study. <i>BMC Psychiatry</i> , 2020, 20, 232.	1.1	11
17	Prevalence trends and individual patterns of antiepileptic drug use in pregnancy 2006–2016: A study in the five Nordic countries, United States, and Australia. <i>Pharmacoepidemiology and Drug Safety</i> , 2020, 29, 913-922.	0.9	23
18	Antipsychotic drug use in pregnancy: A multinational study from ten countries. <i>Schizophrenia Research</i> , 2020, 220, 106-115.	1.1	30

#	ARTICLE	IF	CITATIONS
19	Suicide Ideation and Behavior as Risk Factors for Subsequent Suicide in Schizophrenia: A Nested Caseâ€”Control Study. <i>Suicide and Life-Threatening Behavior</i> , 2019, 49, 996-1005.	0.9	12
20	All-cause mortality in patients with treatment-resistant depression: a cohort study in the US population. <i>Annals of General Psychiatry</i> , 2019, 18, 23.	1.2	44
21	&lt;p&gt;Maternal bereavement shortly before or during pregnancy and risk of postpartum psychotic illness: a population-based study from Denmark and Sweden&lt;/p&gt;. <i>Clinical Epidemiology</i> , 2019, Volume 11, 285-298.	1.5	7
22	Monotherapy vs. combination therapy for post mania maintenance treatment: A population based cohort study. <i>European Neuropsychopharmacology</i> , 2019, 29, 691-700.	0.3	16
23	Violent crime among Swedish military veterans after deployment to Afghanistan: a population-based matched cohort study. <i>International Journal of Epidemiology</i> , 2019, 48, 1604-1613.	0.9	3
24	Risk of Acute Liver Injury in Agomelatine and Other Antidepressant Users in Four European Countries: A Cohort and Nested Caseâ€”Control Study Using Automated Health Data Sources. <i>CNS Drugs</i> , 2019, 33, 383-395.	2.7	23
25	Marriage and divorce after military deployment to Afghanistan: A matched cohort study from Sweden. <i>PLoS ONE</i> , 2019, 14, e0207981.	1.1	10
26	Trends in utilization and dosing of antipsychotic drugs in Scandinavia: Comparison of 2006 and 2016. <i>British Journal of Clinical Pharmacology</i> , 2019, 85, 1598-1606.	1.1	38
27	Treatmentâ€”resistant depression as risk factor for substance use disordersâ€”a nationâ€”wide registerâ€”based cohort study. <i>Addiction</i> , 2019, 114, 1274-1282.	1.7	21
28	Antidepressant use in Denmark, Germany, Spain, and Sweden between 2009 and 2014: Incidence and comorbidities of antidepressant initiators. <i>Journal of Affective Disorders</i> , 2019, 249, 242-252.	2.0	47
29	Initiation and longâ€”term use of benzodiazepines and Zâ€”drugs in bipolar disorder. <i>Bipolar Disorders</i> , 2018, 20, 634-646.	1.1	22
30	Association Between Methylphenidate and Amphetamine Use in Pregnancy and Risk of Congenital Malformations. <i>JAMA Psychiatry</i> , 2018, 75, 167.	6.0	93
31	Humanistic outcomes in treatment resistant depression: a secondary analysis of the STAR*D study. <i>BMC Psychiatry</i> , 2018, 18, 352.	1.1	34
32	Mortality in treatment-resistant unipolar depression: A register-based cohort study in Sweden. <i>Journal of Affective Disorders</i> , 2018, 238, 674-679.	2.0	64
33	Reducing the rehospitalization risk after a manic episode: A population based cohort study of lithium, valproate, olanzapine, quetiapine and aripiprazole in monotherapy and combinations. <i>Journal of Affective Disorders</i> , 2017, 217, 16-23.	2.0	15
34	Season of treatment initiation with antidepressants and suicidal behavior: A population-based cohort study in Sweden. <i>Journal of Affective Disorders</i> , 2017, 215, 245-255.	2.0	1
35	Suicide and all-cause mortality in Swedish deployed military veterans: a population-based matched cohort study. <i>BMJ Open</i> , 2017, 7, e014034.	0.8	13
36	International trends in antipsychotic use: A study in 16 countries, 2005â€”2014. <i>European Neuropsychopharmacology</i> , 2017, 27, 1064-1076.	0.3	185

#	ARTICLE	IF	CITATIONS
37	Risk of breast cancer in risperidone users: A nationwide cohort study. <i>Schizophrenia Research</i> , 2017, 182, 98-103.	1.1	19
38	Suicide risk and antipsychotic side effects in schizophrenia: nested case-control study. <i>Human Psychopharmacology</i> , 2016, 31, 341-345.	0.7	14
39	Serotonergic medication enhances the association between suicide and sunshine. <i>Journal of Affective Disorders</i> , 2016, 189, 276-281.	2.0	35
40	Diabetes and glucose disturbances in patients with psychosis in Sweden. <i>BMJ Open Diabetes Research and Care</i> , 2015, 3, e000120.	1.2	6
41	Predictors for initiation of pharmacological prophylaxis in patients with newly diagnosed bipolar disorder—A nationwide cohort study. <i>Journal of Affective Disorders</i> , 2015, 172, 204-210.	2.0	10
42	Risk and predictors of attempted and completed suicide in patients with hematological malignancies. <i>Psycho-Oncology</i> , 2014, 23, 1276-1282.	1.0	11
43	Medication and suicide risk in schizophrenia: A nested case-control study. <i>Schizophrenia Research</i> , 2013, 150, 416-420.	1.1	37
44	Antipsychotic Prescription Filling in Patients With Schizophrenia or Schizoaffective Disorder. <i>Journal of Clinical Psychopharmacology</i> , 2013, 33, 759-765.	0.7	8
45	Risks of adverse pregnancy and birth outcomes in women treated or not treated with mood stabilisers for bipolar disorder: population based cohort study. <i>BMJ, The</i> , 2012, 345, e7085-e7085.	3.0	185
46	Antipsychotics During Pregnancy. <i>Archives of General Psychiatry</i> , 2012, 69, 715-21.	13.8	111
47	Early non-adherence to medication and other risk factors for rehospitalization in schizophrenia and schizoaffective disorder. <i>Schizophrenia Research</i> , 2011, 133, 36-41.	1.1	72
48	Suicide and hospitalization for mental disorders in Sweden: A population-based case-control study. <i>Journal of Psychiatric Research</i> , 2010, 44, 741-747.	1.5	48
49	Diagnostic profile and suicide risk in schizophrenia spectrum disorder. <i>Schizophrenia Research</i> , 2010, 123, 251-256.	1.1	13
50	Seasonality of suicide in Sweden: Relationship with psychiatric disorder. <i>Journal of Affective Disorders</i> , 2009, 119, 59-65.	2.0	45
51	Risk factors for suicide in schizophrenia: Findings from a Swedish population-based case-control study. <i>Schizophrenia Research</i> , 2009, 108, 231-237.	1.1	70
52	Central nervous system tumours in children in Costa Rica, 1981-96. <i>Paediatric and Perinatal Epidemiology</i> , 2002, 16, 219-225.	0.8	12
53	Trends in use of attention deficit hyperactivity disorder medication among children and adolescents in Scandinavia in 2010–2020. <i>European Child and Adolescent Psychiatry</i> , 0, , .	2.8	2