

Filippo Maggi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7839270/filippo-maggi-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

375
papers

7,454
citations

42
h-index

59
g-index

396
ext. papers

9,476
ext. citations

4.1
avg, IF

6.56
L-index

#	Paper	IF	Citations
375	Optimization of Solvent-Free Microwave-Assisted Hydrodiffusion and Gravity Extraction of L. Fruits Maximizing Polyphenols, Sugar Content, and Biological Activities Using Central Composite Design.. <i>Pharmaceuticals</i> , 2022 , 15,	5.2	2
374	Ethnobotanical investigation of L. grown in El Kala (Algeria), and phytochemical study and antioxidant activity of its essential oil and extracts.. <i>Natural Product Research</i> , 2022 , 1-6	2.3	0
373	Phytotoxic activity of Moldavian dragonhead (<i>Dracocephalum moldavica</i> L.) essential oil and its possible use as bio-herbicide. <i>Process Biochemistry</i> , 2022 , 114, 86-92	4.8	2
372	Evaluation of the chemical constituents, antioxidant and enzyme inhibitory activities of six Yemeni green coffee beans varieties. <i>Food Bioscience</i> , 2022 , 46, 101552	4.9	1
371	Lethal and sublethal effects of essential oil-loaded zein nanocapsules on a zoonotic disease vector mosquito, and their non-target impact. <i>Industrial Crops and Products</i> , 2022 , 176, 114413	5.9	1
370	Lethal and behavioural effects of a green insecticide against an invasive polyphagous fruit fly pest and its safety to mammals. <i>Chemosphere</i> , 2022 , 287, 132089	8.4	5
369	A new HPLC-MS/MS method for the simultaneous determination of 36 polyphenols in blueberry, strawberry and their commercial products and determination of antioxidant activity. <i>Food Chemistry</i> , 2022 , 367, 130743	8.5	15
368	A Comprehensive Phytochemical Analysis of Terpenes, Polyphenols and Cannabinoids, and Micromorphological Characterization of 9 Commercial Varieties of L.. <i>Plants</i> , 2022 , 11,	4.5	2
367	Natural diversity in phenolic components and antioxidant properties of oregano (<i>Origanum vulgare</i> L.) accessions, grown under the same conditions.. <i>Scientific Reports</i> , 2022 , 12, 5813	4.9	4
366	Piperitenone oxide-rich <i>Mentha longifolia</i> essential oil and its nanoemulsion to manage different developmental stages of insect and mite pests attacking stored wheat. <i>Industrial Crops and Products</i> , 2022 , 178, 114600	5.9	3
365	Enhanced Anticancer Activity of <i>Hymenocardia acida</i> Stem Bark Extract Loaded into PLGA Nanoparticles. <i>Pharmaceuticals</i> , 2022 , 15, 535	5.2	0
364	Introducing Three New Fruit-Scented Mints to Farmlands: Insights on Drug Yield, Essential-Oil Quality, and Antioxidant Properties. <i>Antioxidants</i> , 2022 , 11, 866	7.1	3
363	Two Iranian <i>Scrophularia striata</i> Boiss. Ecotypes under UV-B radiation: Germination and initial growth perspective. <i>South African Journal of Botany</i> , 2022 , 148, 460-468	2.9	0
362	Alkaloids and sesquiterpenes from roots and leaves of L. (<i>Solanaceae</i>) with antioxidant and anti-acetylcholinesterase activities. <i>Natural Product Research</i> , 2021 , 35, 2784-2788	2.3	5
361	The essential oil of <i>Moris</i> and its antioxidant and antimicrobial activities. <i>Natural Product Research</i> , 2021 , 35, 5452-5458	2.3	1
360	Essential oil compositions of , subsp. and growing in Sicily and Malta. <i>Natural Product Research</i> , 2021 , 35, 3460-3469	2.3	16
359	Insecticidal activity of two essential oils used in perfumery (ylang ylang and frankincense). <i>Natural Product Research</i> , 2021 , 35, 4746-4752	2.3	6

358	Essential oil variability in Benth populations: a narrow endemic species of Iran. <i>Natural Product Research</i> , 2021 , 35, 2588-2592	2.3	4
357	Chemical Variability in the Composition of <i>Zhumeria majdae</i> (Rech. F. & Wendelbo) Essential Oil According to Storage Time and Temperature. <i>Horticulturae</i> , 2021 , 7, 463	2.5	1
356	Natural diversity in fatty acids profiles and antioxidant properties of sumac fruits (<i>Rhus coriaria</i> L.): Selection of preferable populations for food industries.. <i>Food Chemistry</i> , 2021 , 374, 131757	8.5	1
355	A vibrational in vitro approach to evaluate the potential of monoolein nanoparticles as isofuranodiene carrier in MDA-MB 231 breast cancer cell line: New insights from Infrared and Raman microspectroscopies.. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2021 , 269, 120735	4.4	1
354	Valorization of CBD-hemp through distillation to provide essential oil and improved cannabinoids profile. <i>Scientific Reports</i> , 2021 , 11, 19890	4.9	1
353	Effect of Roasting, Boiling, and Frying Processing on 29 Polyphenolics and Antioxidant Activity in Seeds and Shells of Sweet Chestnut (Mill.). <i>Plants</i> , 2021 , 10,	4.5	3
352	Therapeutic Effects of Hydroalcoholic Extracts from the Ancient Apple <i>Mela Rosa dei Monti Sibillini</i> in Transient Global Ischemia in Rats. <i>Pharmaceuticals</i> , 2021 , 14,	5.2	1
351	Developing a Essential Oil Nanoemulsion for the Eco-Friendly Management of and Larvae and Adults on Stored Wheat. <i>Molecules</i> , 2021 , 26,	4.8	14
350	<i>Carlina acaulis</i> and <i>Trachyspermum ammi</i> essential oils formulated in protein baits are highly toxic and reduce aggressiveness in the medfly, <i>Ceratitis capitata</i> . <i>Industrial Crops and Products</i> , 2021 , 161, 113191	5.9	14
349	An insight into <i>Verbascum bombyciferum</i> extracts: Different extraction methodologies, biological abilities and chemical profiles. <i>Industrial Crops and Products</i> , 2021 , 161, 113201	5.9	7
348	Improvement of dragonhead (<i>Dracocephalum moldavica</i> L.) yield quality through a coupled intercropping system and vermicompost application along with maintenance of soil microbial activity. <i>Land Degradation and Development</i> , 2021 , 32, 2833-2848	4.4	10
347	Isofuranodiene, a Natural Sesquiterpene Isolated from Wild Celery (L.), Protects Rats against Acute Ischemic Stroke. <i>Pharmaceuticals</i> , 2021 , 14,	5.2	3
346	Chemical Composition and Broad-Spectrum Insecticidal Activity of the Flower Essential Oil from an Ancient Sicilian Food Plant, <i>Ridolfia segetum</i> . <i>Agriculture (Switzerland)</i> , 2021 , 11, 304	3	14
345	Antimicrobial Activity and Chemical Composition of Essential Oil from <i>Thymus daenensis</i> and <i>Thymus fedtschenkoi</i> During Phenological Stages. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2021 , 24, 469-479	1.7	2
344	<i>Vepris macrophylla</i> Essential Oil Produces Notable Antiproliferative Activity and Morphological Alterations in Human Breast Adenocarcinoma Cells. <i>Applied Sciences (Switzerland)</i> , 2021 , 11, 4369	2.6	0
343	Sub-Tissue Localization of Phytochemicals in (L.) J. Presl. Growing in Northern Italy. <i>Plants</i> , 2021 , 10,	4.5	2
342	A new chemotype with high tricyclene content from the essential oil of L. growing in Algerian Pre-Sahara. <i>Natural Product Research</i> , 2021 , 1-6	2.3	1
341	Occurrence of flavonoids in different Lamiaceae taxa for a preliminary study on their evolution based on phytochemistry. <i>Biochemical Systematics and Ecology</i> , 2021 , 96, 104247	1.4	5

340	Insecticidal, antibacterial and dye adsorbent properties of <i>Sargassum muticum</i> decorated nano-silver particles. <i>South African Journal of Botany</i> , 2021 , 139, 432-441	2.9	3
339	Influence of Freezing and Different Drying Methods on Volatile Profiles of Strawberry and Analysis of Volatile Compounds of Strawberry Commercial Jams. <i>Molecules</i> , 2021 , 26,	4.8	3
338	Composition and biological activities of the essential oil from a Sicilian accession of (L.) Lindl. <i>Natural Product Research</i> , 2021 , 35, 733-743	2.3	12
337	Vermicompost Application in Different Intercropping Patterns Improves the Mineral Nutrient Uptake and Essential Oil Compositions of Sweet Basil (<i>Ocimum basilicum</i> L.). <i>Journal of Soil Science and Plant Nutrition</i> , 2021 , 21, 450-466	3.2	11
336	Comparison of drying methods for the extraction of essential oil from dragonhead (<i>Dracocephalum moldavica</i> L., Lamiaceae). <i>Journal of Essential Oil Research</i> , 2021 , 33, 162-170	2.3	4
335	Coumarin (2H-1-benzopyran-2-one): a novel and eco-friendly aphicide. <i>Natural Product Research</i> , 2021 , 35, 1566-1571	2.3	1
334	Chemical compositions and biological activity of essential oils from four populations of <i>Satureja macrantha</i> C.A.Mey. <i>Journal of Essential Oil Research</i> , 2021 , 33, 133-142	2.3	4
333	Chemical constituents and anticholinesterase activity of the essential oil of Algerian (<i>Desf.</i>) maire. <i>Natural Product Research</i> , 2021 , 1-6	2.3	1
332	Phytochemical Profile and Biological Activities of Crude and Purified Extracts. <i>Plants</i> , 2021 , 10,	4.5	3
331	Effects of Essential Oils from spp. and on Biofilm and Virulence Properties of O157:H7. <i>Antibiotics</i> , 2021 , 10,	4.9	5
330	Phytotoxic Potential and Phenolic Profile of Extracts from. <i>Plants</i> , 2021 , 10,	4.5	8
329	Enhancement of In Vitro Production of Volatile Organic Compounds by Shoot Differentiation in. <i>Plants</i> , 2021 , 10,	4.5	4
328	Effect of Active-Edible Coating and Essential Oils on Lamb Patties Oxidation during Display. <i>Foods</i> , 2021 , 10,	4.9	10
327	Essential Oils as Natural Sources of Fragrance Compounds for Cosmetics and Cosmeceuticals. <i>Molecules</i> , 2021 , 26,	4.8	83
326	Encapsulation of <i>Carlina acaulis</i> essential oil and carlina oxide to develop long-lasting mosquito larvicides: microemulsions versus nanoemulsions. <i>Journal of Pest Science</i> , 2021 , 94, 899-915	5.5	12
325	Composition and profiling of essential oil, volatile and crude extract constituents of <i>Micromeria inodora</i> growing in western Algeria. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021 , 195, 113856	3.5	1
324	<i>Funneliformis mosseae</i> inoculation under water deficit stress improves the yield and phytochemical characteristics of thyme in intercropping with soybean. <i>Scientific Reports</i> , 2021 , 11, 15279	4.9	9
323	Efficacy of 12 commercial essential oils as wheat protectants against stored-product beetles, and their acetylcholinesterase inhibitory activity. <i>Entomologia Generalis</i> , 2021 , 41, 385-414	5.3	5

322	A Design of Experiment (DoE) Approach to Model the Yield and Chemical Composition of Ajowan (L.) Essential Oil Obtained by Microwave-Assisted Extraction. <i>Pharmaceuticals</i> , 2021 , 14,	5.2	3
321	Volatile Organic Compounds of the Glandular Trichomes of <i>Ocimum basilicum</i> and Artifacts during the Distillation of the Leaves. <i>Applied Sciences (Switzerland)</i> , 2021 , 11, 7312	2.6	1
320	Two Medicinal Plants (<i>Alkanna trichophila</i> and <i>Convolvulus galaticus</i>) from Turkey: Chemical Characterization and Biological Perspectives. <i>Chemistry and Biodiversity</i> , 2021 , 18, e2100356	2.5	0
319	Isofuranodiene-based nanoemulsion: larvicidal and adulticidal activity against tenebrionid beetles attacking stored wheat. <i>Journal of Stored Products Research</i> , 2021 , 93, 101859	2.5	2
318	Evaluation of chemical constituents and biological properties of two endemic <i>Verbascum</i> species. <i>Process Biochemistry</i> , 2021 , 108, 110-120	4.8	6
317	Bioactivity of Essential Oil and Its Main Component towards the Olive Fruit Fly, δ : Ingestion Toxicity, Electrophysiological and Behavioral Insights. <i>Insects</i> , 2021 , 12,	2.8	2
316	The chemical composition of the aerial parts essential oil of subsp. (Lamiaceae) growing in Sicily (Italy). <i>Natural Product Research</i> , 2021 , 1-5	2.3	
315	Chemical composition, antioxidant and anticholinesterase activity of the essential oil of algerian L. <i>Natural Product Research</i> , 2021 , 1-9	2.3	3
314	Intraspecific divergence in phytochemical characteristics and drought tolerance of two carvacrol-rich <i>Origanum vulgare</i> subspecies: subsp. <i>hirtum</i> and subsp. <i>gracile</i> . <i>Industrial Crops and Products</i> , 2021 , 168, 113557	5.9	7
313	Toxics or Lures? Biological and Behavioral Effects of Plant Essential Oils on Tephritidae Fruit Flies. <i>Molecules</i> , 2021 , 26,	4.8	4
312	Prolonged sublethal effects of essential oils from non-wood parts of nine conifers on key insect pests and vectors. <i>Industrial Crops and Products</i> , 2021 , 168, 113590	5.9	12
311	Spilanthol-rich essential oil obtained by microwave-assisted extraction from <i>Acmella oleracea</i> (L.) R.K. Jansen and its nanoemulsion: Insecticidal, cytotoxic and anti-inflammatory activities. <i>Industrial Crops and Products</i> , 2021 , 172, 114027	5.9	2
310	Apiaceae essential oils and their constituents as insecticides against mosquitoes: A review. <i>Industrial Crops and Products</i> , 2021 , 171, 113892	5.9	8
309	<i>Tanacetum vulgare</i> essential oil as grain protectant against adults and larvae of four major stored-product insect pests. <i>Journal of Stored Products Research</i> , 2021 , 94, 101882	2.5	4
308	Essential oils from three Algerian medicinal plants (<i>Artemisia campestris</i> , <i>Pulicaria arabica</i> , and <i>Saccocalyx saturoioides</i>) as new botanical insecticides?. <i>Environmental Science and Pollution Research</i> , 2020 , 27, 26594-26604	5.1	12
307	Chemical Composition, Antifungal and Insecticidal Activities of the Essential Oils from Tunisian Subsp. and Subsp.. <i>Molecules</i> , 2020 , 25,	4.8	8
306	Fiber-Sample Distance, An Important Parameter To Be Considered in Headspace Solid-Phase Microextraction Applications. <i>Analytical Chemistry</i> , 2020 , 92, 7478-7484	7.8	11
305	Essential Oil Chemical Variability in (Apiaceae) from Different Regions of Iran and Its Relationship with Environmental Factors. <i>Plants</i> , 2020 , 9,	4.5	6

304	Chemical Composition, Antioxidant and Enzyme Inhibitory Properties of Different Extracts Obtained from Spent Coffee Ground and Coffee Silverskin. <i>Foods</i> , 2020 , 9,	4.9	15
303	Comprehensive characterization of phytochemicals and biological activities of the Italian ancient apple 'Mela Rosa dei Monti Sibillini'. <i>Food Research International</i> , 2020 , 137, 109422	7	8
302	Hairy Garlic () from Sicily (Italy): LC-DAD-MS Analysis of Secondary Metabolites and In Vitro Biological Properties. <i>Molecules</i> , 2020 , 25,	4.8	12
301	Cannabidiol-enriched hemp essential oil obtained by an optimized microwave-assisted extraction using a central composite design. <i>Industrial Crops and Products</i> , 2020 , 154, 112688	5.9	35
300	Effectiveness of eight essential oils against two key stored-product beetles, <i>Prostephanus truncatus</i> (Horn) and <i>Trogoderma granarium</i> Everts. <i>Food and Chemical Toxicology</i> , 2020 , 139, 111255	4.7	26
299	Effect of different fertilizer sources and harvesting time on the growth characteristics, nutrient uptakes, essential oil productivity and composition of <i>Mentha x piperita</i> L.. <i>Industrial Crops and Products</i> , 2020 , 148, 112290	5.9	26
298	Combustion of Nanoaluminum and Magnesium in Fuel-Rich Propellants. <i>Propellants, Explosives, Pyrotechnics</i> , 2020 , 45, 724-729	1.7	5
297	Exploring essential oils of Slovak medicinal plants for insecticidal activity: The case of <i>Thymus alternans</i> and <i>Teucrium montanum</i> subsp. <i>jailae</i> . <i>Food and Chemical Toxicology</i> , 2020 , 138, 111203	4.7	9
296	Acaricidal activity, mode of action, and persistent efficacy of selected essential oils on the poultry red mite (<i>Dermanyssus gallinae</i>). <i>Food and Chemical Toxicology</i> , 2020 , 138, 111207	4.7	10
295	Efficacy of the furanosesquiterpene isofuranodiene against the stored-product insects <i>Prostephanus truncatus</i> (Coleoptera: Bostrychidae) and <i>Trogoderma granarium</i> (Coleoptera: Dermestidae). <i>Journal of Stored Products Research</i> , 2020 , 86, 101553	2.5	14
294	Acaricidal properties of hemp (<i>Cannabis sativa</i> L.) essential oil against <i>Dermanyssus gallinae</i> and <i>Hyalomma dromedarii</i> . <i>Industrial Crops and Products</i> , 2020 , 147, 112238	5.9	20
293	Ascaridole-rich essential oil from marsh rosemary (<i>Ledum palustre</i>) growing in Poland exerts insecticidal activity on mosquitoes, moths and flies without serious effects on non-target organisms and human cells. <i>Food and Chemical Toxicology</i> , 2020 , 138, 111184	4.7	11
292	Optimization of espresso coffee extraction through variation of particle sizes, perforated disk height and filter basket aimed at lowering the amount of ground coffee used. <i>Food Chemistry</i> , 2020 , 314, 126220	8.5	11
291	Chemical Composition and Antibacterial Activity of Essential Oils from the Algerian Endemic <i>Desf.</i> against Multidrug-Resistant Uropathogenic Isolates. <i>Antibiotics</i> , 2020 , 9,	4.9	13
290	The volatile oils from the oleo-gum-resins of <i>Ferula assa-foetida</i> and <i>Ferula gummosa</i> : A comprehensive investigation of their insecticidal activity and eco-toxicological effects. <i>Food and Chemical Toxicology</i> , 2020 , 140, 111312	4.7	15
289	Quantification of 2- and 3-isopropylmalic acids in forty Italian wines by UHPLC-MS/MS triple quadrupole and evaluation of their antimicrobial, antioxidant activities and biocompatibility. <i>Food Chemistry</i> , 2020 , 321, 126726	8.5	7
288	Chemical Composition, Antibacterial and Radical Scavenging Activity of Essential Oils from <i>C.A.Mey.</i> at Different Growth Stages. <i>Foods</i> , 2020 , 9,	4.9	12
287	(Baker) I. Verd Essential Oil: An Antifungal Agent against Phytopathogenic Fungi. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	4

286	Hepatoprotective Effects of Standardized Extracts from an Ancient Italian Apple Variety (Mela Rosa dei Monti Sibillini) against Carbon Tetrachloride (CCl ₄)-Induced Hepatotoxicity in Rats. <i>Molecules</i> , 2020 , 25,	4.8	4
285	A new analytical method for the simultaneous quantification of isoflavones and lignans in 25 green coffee samples by HPLC-MS/MS. <i>Food Chemistry</i> , 2020 , 325, 126924	8.5	8
284	Total phytochemical analysis of <i>Thymus munbyanus</i> subsp. <i>coloratus</i> from Algeria by HS-SPME-GC-MS, NMR and HPLC-MS studies. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020 , 186, 113330	3.5	10
283	Phenolic monoterpene-rich essential oils from Apiaceae and Lamiaceae species: insecticidal activity and safety evaluation on non-target earthworms. <i>Entomologia Generalis</i> , 2020 , 40, 421-435	5.3	20
282	Intercropping fennel (<i>Foeniculum vulgare</i> L.) with common bean (<i>Phaseolus vulgaris</i> L.) as affected by PGPR inoculation: A strategy for improving yield, essential oil and fatty acid composition. <i>Scientia Horticulturae</i> , 2020 , 261, 108951	4.1	29
281	Effects of active edible coating based on thyme and garlic essential oils on lamb meat shelf life after long-term frozen storage. <i>Journal of the Science of Food and Agriculture</i> , 2020 , 100, 656-664	4.3	21
280	Essential oils of hedgenettles (<i>Stachys inflata</i> , <i>S. lavandulifolia</i> , and <i>S. byzantina</i>) have antioxidant, anti-Alzheimer, antidiabetic, and anti-obesity potential: A comparative study. <i>Industrial Crops and Products</i> , 2020 , 145, 112089	5.9	8
279	Developing green insecticides to manage olive fruit flies? Ingestion toxicity of four essential oils in protein baits on <i>Bactrocera oleae</i> . <i>Industrial Crops and Products</i> , 2020 , 143, 111884	5.9	20
278	Outstanding insecticidal activity and sublethal effects of <i>Carlina acaulis</i> root essential oil on the housefly, <i>Musca domestica</i> , with insights on its toxicity on human cells. <i>Food and Chemical Toxicology</i> , 2020 , 136, 111037	4.7	40
277	Developing a Highly Stable Essential Oil Nanoemulsion for Managing. <i>Nanomaterials</i> , 2020 , 10,	5.4	29
276	Chemical Composition and Antiproliferative Effect of Essential Oils of Four <i>Solidago</i> Species (<i>S. canadensis</i> , <i>S. gigantea</i> , <i>S. virgaurea</i> and <i>S. Biederederi</i>). <i>Chemistry and Biodiversity</i> , 2020 , 17, e2000685	2.5	1
275	Exploring the bio-control efficacy of <i>Artemisia fragrans</i> essential oil on the perennial weed <i>Convolvulus arvensis</i> : Inhibitory effects on the photosynthetic machinery and induction of oxidative stress. <i>Industrial Crops and Products</i> , 2020 , 155, 112785	5.9	20
274	Chitosan nanoemulsions of cold-pressed orange essential oil to preserve fruit juices. <i>International Journal of Food Microbiology</i> , 2020 , 331, 108786	5.8	15
273	Phytochemical Analysis and Trypanocidal Activity of Desr. <i>Molecules</i> , 2020 , 25,	4.8	2
272	Characterization of Odor-Active Compounds, Polyphenols, and Fatty Acids in Coffee Silverskin. <i>Molecules</i> , 2020 , 25,	4.8	9
271	Comparative Study of the Chemical Compositions and Antioxidant Activities of Fresh Juices from Romanian Cucurbitaceae Varieties. <i>Molecules</i> , 2020 , 25,	4.8	10
270	Mosquitocidal and Anti-Inflammatory Properties of The Essential Oils Obtained from Monoecious, Male, and Female Inflorescences of Hemp (<i>L.</i>) and Their Encapsulation in Nanoemulsions. <i>Molecules</i> , 2020 , 25,	4.8	11
269	Phytol, (E)-nerolidol and spathulenol from <i>Stevia rebaudiana</i> leaf essential oil as effective and eco-friendly botanical insecticides against <i>Metopolophium dirhodum</i> . <i>Industrial Crops and Products</i> , 2020 , 155, 112844	5.9	21

268	A new ionone derivative from Boiss. (Solanaceae). <i>Natural Product Research</i> , 2020 , 1-8	2.3	2
267	Subsp. (Guss.) Troia & Raimondo from Sicily (Italy): Isolation of Essential Oil and Evaluation of Its Bioactivity. <i>Molecules</i> , 2020 , 25,	4.8	15
266	The Variability of Thymol and Carvacrol Contents Reveals the Level of Antibacterial Activity of the Essential Oils from Different Accessions of. <i>Antibiotics</i> , 2020 , 9,	4.9	7
265	In Vitro Scolicidal Activity of the Sesquiterpenes Isofuranodiene, β Bisabolol and Farnesol on Protozoa. <i>Molecules</i> , 2020 , 25,	4.8	3
264	Essential Oil Nanoemulsion Toxicity against ? Shedding Light on Its Interactions with Aspartate Aminotransferase and Alanine Aminotransferase by Molecular Docking. <i>Molecules</i> , 2020 , 25,	4.8	6
263	Butter oil (ghee) enrichment with aromatic plants: Chemical characterization and effects on fibroblast migration in an in-vitro wound healing model. <i>Arabian Journal of Chemistry</i> , 2020 , 13, 8909-8919	5.9	3
262	Encapsulation of sea fennel (<i>Crithmum maritimum</i>) essential oil in nanoemulsion and SiO ₂ nanoparticles for treatment of the crop pest <i>Spodoptera litura</i> and the dengue vector <i>Aedes aegypti</i> . <i>Industrial Crops and Products</i> , 2020 , 158, 113033	5.9	12
261	The aromatic ginger <i>Kaempferia galanga</i> L. (Zingiberaceae) essential oil and its main compounds are effective larvicidal agents against <i>Aedes vittatus</i> and <i>Anopheles maculatus</i> without toxicity on the non-target aquatic fauna. <i>Industrial Crops and Products</i> , 2020 , 158, 113012	5.9	10
260	Essential oil composition and biological activities of Poir (Fabaceae). <i>Natural Product Research</i> , 2020 , 1-6	2.3	2
259	Chemical composition of the essential oil of (<i>L.</i>) Bertol subsp. (Desf.) Fiori (Umbelliferae) collected wild in Central Sicily and its antimicrobial activity. <i>Natural Product Research</i> , 2020 , 1-9	2.3	10
258	Chemical composition and biological activities of the essential oil from (<i>L.</i>) C. A. Mey. growing wild in Egypt. <i>Natural Product Research</i> , 2020 , 34, 2358-2362	2.3	13
257	Quality assessment of commercial samples. <i>Natural Product Research</i> , 2020 , 34, 3154-3157	2.3	0
256	Promising insecticidal efficacy of the essential oils from the halophyte <i>Echinophora spinosa</i> (Apiaceae) growing in Corsica Island, France. <i>Environmental Science and Pollution Research</i> , 2020 , 27, 14454-14464	5.1	9
255	Phytochemical analysis of <i>Rhazya stricta</i> extract and its use in fabrication of silver nanoparticles effective against mosquito vectors and microbial pathogens. <i>Science of the Total Environment</i> , 2020 , 700, 134443	10.2	24
254	Insecticidal and mosquito repellent efficacy of the essential oils from stem bark and wood of <i>Hazomalania voyronii</i> . <i>Journal of Ethnopharmacology</i> , 2020 , 248, 112333	5	17
253	Nanostructured liquid crystalline particles as delivery vectors for isofuranodiene: Characterization and in-vitro anticancer activity. <i>Colloids and Surfaces B: Biointerfaces</i> , 2020 , 192, 111050	6	6
252	Spent coffee grounds: A potential commercial source of phytosterols. <i>Food Chemistry</i> , 2020 , 325, 126836	6.5	12
251	Optimization of edible <i>Alyssum homalocarpum</i> seed gum-chitosan coating formulation to improve the postharvest storage potential and quality of apricot (<i>Prunus armeniaca</i> L.). <i>Journal of Food Safety</i> , 2020 , 40, e12805	2	4

250	Himalayan Nettle as a Candidate Ingredient for Pharmaceutical and Nutraceutical Applications-Phytochemical Analysis and In Vitro Bioassays. <i>Molecules</i> , 2020 , 25,	4.8	8
249	The Essential Oil of and its Application as A Biocide on Stone and Derived Surfaces. <i>Plants</i> , 2019 , 8,	4.5	20
248	Green Micro- and Nanoemulsions for Managing Parasites, Vectors and Pests. <i>Nanomaterials</i> , 2019 , 9,	5.4	62
247	Sesquiterpene rich essential oil from Nepalese Bael tree (Aegle marmelos (L.) Correa) as potential antiproliferative agent. <i>Phytotherapy</i> 2019 , 138, 104266	3.2	5
246	Plant extracts for developing mosquito larvicides: From laboratory to the field, with insights on the modes of action. <i>Acta Tropica</i> , 2019 , 193, 236-271	3.2	108
245	Rationale for developing novel mosquito larvicides based on isofuranodiene microemulsions. <i>Journal of Pest Science</i> , 2019 , 92, 909-921	5.5	41
244	Carlina oxide from Carlina acaulis root essential oil acts as a potent mosquito larvicide. <i>Industrial Crops and Products</i> , 2019 , 137, 356-366	5.9	35
243	Chemical profiles and insecticidal efficacy of the essential oils from four Thymus taxa growing in central-southern Italy. <i>Industrial Crops and Products</i> , 2019 , 138, 111460	5.9	24
242	Evaluation of common bean (Phaseolus vulgaris L.) seed yield and quali-quantitative production of the essential oils from fennel (Foeniculum vulgare Mill.) and dragonhead (Dracocephalum moldavica L.) in intercropping system under humic acid application. <i>Journal of Cleaner Production</i> , 2019 , 235, 112-122	10.3	25
241	Microemulsions: An effective encapsulation tool to enhance the antimicrobial activity of selected EOs. <i>Journal of Drug Delivery Science and Technology</i> , 2019 , 53, 101101	4.5	22
240	In Vitro and In Vivo Effectiveness of Carvacrol, Thymol and Linalool against. <i>Molecules</i> , 2019 , 24,	4.8	25
239	Innate positive chemotaxis to paeonal from highly attractive Chinese medicinal herbs in the cigarette beetle, Lasioderma serricorne. <i>Scientific Reports</i> , 2019 , 9, 6995	4.9	3
238	Efficacy of Two Monoterpenoids, Carvacrol and Thymol, and Their Combinations against Eggs and Larvae of the West Nile Vector. <i>Molecules</i> , 2019 , 24,	4.8	29
237	Effect of salinity stress on the physiological characteristics, phenolic compounds and antioxidant activity of Thymus vulgaris L. and Thymus daenensis Celak. <i>Industrial Crops and Products</i> , 2019 , 135, 311-320	5.9	94
236	Green nanoemulsion interventions for biopesticide formulations 2019 , 133-160		5
235	Characterization of nutrients, polyphenols and volatile components of the ancient apple cultivar 'Mela Rosa Dei Monti Sibillini' from Marche region, central Italy. <i>International Journal of Food Sciences and Nutrition</i> , 2019 , 70, 796-812	3.7	8
234	The Nonvolatile and Volatile Metabolites of Prangos ferulacea and Their Biological Properties. <i>Planta Medica</i> , 2019 , 85, 815-824	3.1	7
233	Triterpene Acid and Phenolics from Ancient Apples of Friuli Venezia Giulia as Nutraceutical Ingredients: LC-MS Study and In Vitro Activities. <i>Molecules</i> , 2019 , 24,	4.8	27

232	Antioxidant and Anti-Inflammatory Properties of Oil in Human Pre-Adipocytes. <i>Antioxidants</i> , 2019 , 8,	7.1	49
231	Exploring the Insecticidal Potential of Boldo () Essential Oil: Toxicity to Pests and Vectors and Non-target Impact on the Microcrustacean. <i>Molecules</i> , 2019 , 24,	4.8	9
230	Insecticidal activity of the essential oil and polar extracts from <i>Ocimum gratissimum</i> grown in Ivory Coast: Efficacy on insect pests and vectors and impact on non-target species. <i>Industrial Crops and Products</i> , 2019 , 132, 377-385	5.9	31
229	Antimicrobial efficacy of <i>Thymbra capitata</i> (L.) Cav. essential oil loaded in self-assembled zein nanoparticles in combination with heat. <i>Industrial Crops and Products</i> , 2019 , 133, 98-104	5.9	32
228	Curcumin: Total-Scale Analysis of the Scientific Literature. <i>Molecules</i> , 2019 , 24,	4.8	32
227	<i>Origanum syriacum</i> subsp. <i>syracum</i> : From an ingredient of Lebanese <i>hanousheh</i> to a source of effective and eco-friendly botanical insecticides. <i>Industrial Crops and Products</i> , 2019 , 134, 26-32	5.9	29
226	Enhancement of the antifungal activity of thyme and dill essential oils against <i>Colletotrichum nympheae</i> by nano-encapsulation with copper NPs. <i>Industrial Crops and Products</i> , 2019 , 132, 213-225	5.9	21
225	Fixed oil from seeds of narrow-leaved ash (<i>F. angustifolia</i> subsp. <i>angustifolia</i>): Chemical profile, antioxidant and antiproliferative activities. <i>Food Research International</i> , 2019 , 119, 369-377	7	9
224	Anti- activity of hemlock () essential oil. <i>Natural Product Research</i> , 2019 , 33, 3436-3440	2.3	11
223	Insecticidal efficacy of the essential oil of jamb[<i>Acmella oleracea</i> (L.) R.K. Jansen) cultivated in central Italy against filariasis mosquito vectors, houseflies and moth pests. <i>Journal of Ethnopharmacology</i> , 2019 , 229, 272-279	5	28
222	Secondary metabolites, secretory structures and biological activity of water celery (<i>Apium nodiflorum</i> (L.) Lag.) growing in central Italy. <i>Plant Biosystems</i> , 2019 , 153, 325-335	1.6	7
221	Preliminary evaluation of quince (<i>Cydonia oblonga</i> Mill.) fruit as extraction source of antioxidant phytoconstituents for nutraceutical and functional food applications. <i>Journal of the Science of Food and Agriculture</i> , 2019 , 99, 1046-1054	4.3	11
220	Exploring new applications of tulip tree (<i>Liriodendron tulipifera</i> L.): leaf essential oil as apoptotic agent for human glioblastoma. <i>Environmental Science and Pollution Research</i> , 2019 , 26, 30485-30497	5.1	10
219	Enhanced Duration of Truffle Sauce Preservation due to Addition of Linoleic Acid. <i>Journal of Food Quality</i> , 2019 , 2019, 1-10	2.7	3
218	Species Secondary Metabolites Chemodiversity and Bioactivities. <i>Frontiers in Plant Science</i> , 2019 , 10, 834	6.2	19
217	Efficacy of Essential Oil against the Mosquito Vector and the Gastrointestinal Parasite , with Insights on Acetylcholinesterase Inhibition. <i>Molecules</i> , 2019 , 24,	4.8	10
216	Lyme disease is on the rise [How about tick repellents? A global view. <i>Entomologia Generalis</i> , 2019 , 39, 61-72	5.3	16
215	Antioxidant and Enzyme Inhibitory Properties of the Polyphenolic-Rich Extract from an Ancient Apple Variety of Central Italy (Mela Rosa dei Monti Sibillini). <i>Plants</i> , 2019 , 9,	4.5	8

214	The emulsion made with essential oil and aromatic water from <i>Oliveria decumbens</i> protects murine macrophages from LPS-induced oxidation and exerts relevant radical scavenging activities. <i>Biocatalysis and Agricultural Biotechnology</i> , 2019 , 17, 538-544	4.2	12
213	Medicinal plants and their traditional uses in the highland region of Bordj Bou Arreridj (Northeast Algeria). <i>Journal of Herbal Medicine</i> , 2019 , 16, 100262	2.3	28
212	Chemical Characterization of Leaves, Male and Female Flowers from Spontaneous Cannabis (<i>Cannabis sativa</i> L.) Growing in Hungary. <i>Chemistry and Biodiversity</i> , 2019 , 16, e1800562	2.5	32
211	Protective effects of hydroalcoholic extracts from an ancient apple variety 'Mela Rosa dei Monti Sibillini' against renal ischemia/reperfusion injury in rats. <i>Food and Function</i> , 2019 , 10, 7544-7552	6.1	7
210	Microemulsions for delivery of Apiaceae essential oils: Towards highly effective and eco-friendly mosquito larvicides?. <i>Industrial Crops and Products</i> , 2019 , 129, 631-640	5.9	76
209	Preliminary study on the phytochemical evolution of different Lamiaceae species based on iridoids. <i>Biochemical Systematics and Ecology</i> , 2019 , 82, 44-51	1.4	6
208	Comparison of chemical composition and antioxidant activities of two Winter savory subspecies (subsp. and subsp.) cultivated in Northern Italy. <i>Natural Product Research</i> , 2019 , 33, 3143-3147	2.3	11
207	<i>Paeonia arietina</i> and <i>Paeonia kesrounansis</i> bioactive constituents: NMR, LC-DAD-MS fingerprinting and in vitro assays. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2019 , 165, 1-11	3.5	13
206	Towards green drugs against cestodes: Effectiveness of <i>Pelargonium roseum</i> and <i>Ferula gummosa</i> essential oils and their main component on <i>Echinococcus granulosus</i> protoscoleces. <i>Veterinary Parasitology</i> , 2019 , 266, 84-87	2.8	8
205	Essential oil composition and total metabolite content of a chemotype of <i>Ajuga reptans</i> L. (Lamiaceae) collected in Central Italy. <i>Plant Biosystems</i> , 2019 , 153, 552-558	1.6	5
204	Valorizing industrial hemp (<i>Cannabis sativa</i> L.) by-products: Cannabidiol enrichment in the inflorescence essential oil optimizing sample pre-treatment prior to distillation. <i>Industrial Crops and Products</i> , 2019 , 128, 581-589	5.9	65
203	Isofuranodiene synergizes with temozolomide in inducing glioma cells death. <i>Phytomedicine</i> , 2019 , 52, 51-59	6.5	17
202	Evaluation of two invasive plant invaders in Europe (<i>Solidago canadensis</i> and <i>Solidago gigantea</i>) as possible sources of botanical insecticides. <i>Journal of Pest Science</i> , 2019 , 92, 805-821	5.5	19
201	Essential oil composition of aerial parts from Algerian <i>Anacyclus monanthos</i> subsp. <i>cyrtolepidioides</i> (Pomel) Humphries. <i>Natural Product Research</i> , 2019 , 33, 292-295	2.3	1
200	The water extract of tutsan (<i>Hypericum androsaemum</i> L.) red berries exerts antidepressive-like effects and in vivo antioxidant activity in a mouse model of post-stroke depression. <i>Biomedicine and Pharmacotherapy</i> , 2018 , 99, 290-298	7.5	23
199	Aniseed (<i>Pimpinella anisum</i> L.) essential oil reduces pro-inflammatory cytokines and stimulates mucus secretion in primary airway bronchial and tracheal epithelial cell lines. <i>Industrial Crops and Products</i> , 2018 , 114, 81-86	5.9	18
198	Anxiolytic and antidepressant activities of <i>Pelargonium roseum</i> essential oil on Swiss albino mice: Possible involvement of serotonergic transmission. <i>Phytotherapy Research</i> , 2018 , 32, 1014-1022	6.7	16
197	Chemical composition and antibacterial activity of seven uncommon essential oils. <i>Journal of Essential Oil Research</i> , 2018 , 30, 233-243	2.3	13

196	Volatile components, polar constituents and biological activity of tansy daisy (<i>Tanacetum macrophyllum</i> (Waldst. et Kit.) Schultz Bip.). <i>Industrial Crops and Products</i> , 2018 , 118, 225-235	5.9	23
195	Green drugs in the fight against <i>Anisakis simplex</i> -larvicidal activity and acetylcholinesterase inhibition of <i>Origanum compactum</i> essential oil. <i>Parasitology Research</i> , 2018 , 117, 861-867	2.4	33
194	High efficacy of (Z)- β -bisabolene from the essential oil of <i>Galinsoga parviflora</i> (Asteraceae) as larvicide and oviposition deterrent against six mosquito vectors. <i>Environmental Science and Pollution Research</i> , 2018 , 25, 10555-10566	5.1	16
193	The desert wormwood (<i>Artemisia herba-alba</i>) - From Arabian folk medicine to a source of green and effective nanoinsecticides against mosquito vectors. <i>Journal of Photochemistry and Photobiology B: Biology</i> , 2018 , 180, 225-234	6.7	23
192	Acute and sub-lethal toxicity of eight essential oils of commercial interest against the filariasis mosquito <i>Culex quinquefasciatus</i> and the housefly <i>Musca domestica</i> . <i>Industrial Crops and Products</i> , 2018 , 112, 668-680	5.9	82
191	<i>Pimpinella anisum</i> essential oil nanoemulsions against <i>Tribolium castaneum</i> -insecticidal activity and mode of action. <i>Environmental Science and Pollution Research</i> , 2018 , 25, 18802-18812	5.1	100
190	Essential oil profile of oregano (<i>Origanum vulgare</i> L.) populations grown under similar soil and climate conditions. <i>Industrial Crops and Products</i> , 2018 , 119, 183-190	5.9	33
189	Essential oils (EOs), pressurized liquid extracts (PLE) and carbon dioxide supercritical fluid extracts (SFE-CO ₂) from Algerian <i>Thymus munbyanus</i> as valuable sources of antioxidants to be used on an industrial level. <i>Food Chemistry</i> , 2018 , 260, 289-298	8.5	26
188	Identification of highly effective antitrypanosomal compounds in essential oils from the Apiaceae family. <i>Ecotoxicology and Environmental Safety</i> , 2018 , 156, 154-165	7	41
187	Oviposition inhibitory activity of the Mexican sunflower <i>Tithonia diversifolia</i> (Asteraceae) polar extracts against the two-spotted spider mite <i>Tetranychus urticae</i> (Tetranychidae). <i>Physiological and Molecular Plant Pathology</i> , 2018 , 101, 85-92	2.6	12
186	High toxicity of camphene and β -elemene from <i>Wedelia prostrata</i> essential oil against larvae of <i>Spodoptera litura</i> (Lepidoptera: Noctuidae). <i>Environmental Science and Pollution Research</i> , 2018 , 25, 10383-10391	5.1	25
185	Mosquito control with green nanopesticides: towards the One Health approach? A review of non-target effects. <i>Environmental Science and Pollution Research</i> , 2018 , 25, 10184-10206	5.1	82
184	Chemical composition and antioxidant activity of essential oils in <i>Origanum vulgare</i> subsp. <i>gracile</i> at different phenological stages and plant parts. <i>Journal of Food Processing and Preservation</i> , 2018 , 42, e13516	2.1	27
183	<i>Clausena anisata</i> and <i>Dysphania ambrosioides</i> essential oils: from ethno-medicine to modern uses as effective insecticides. <i>Environmental Science and Pollution Research</i> , 2018 , 25, 10493-10503	5.1	49
182	Larvicidal Activity of Essential Oils of Five Apiaceae Taxa and Some of Their Main Constituents Against <i>Culex quinquefasciatus</i> . <i>Chemistry and Biodiversity</i> , 2018 , 15, e1700382	2.5	37
181	Poly(Styrene Sulfonate)/Poly(Allylamine Hydrochloride) Encapsulation of TiO ₂ Nanoparticles Boosts Their Toxic and Repellent Activity Against Zika Virus Mosquito Vectors. <i>Journal of Cluster Science</i> , 2018 , 29, 27-39	3	10
180	Variation in the essential oil yields and compositions of Myrtle (<i>Myrtus communis</i> L.) Populations collected from natural habitats of Southern Iran. <i>Journal of Essential Oil Research</i> , 2018 , 30, 369-378	2.3	1
179	Supercritical CO ₂ extracts and essential oils from <i>Teucrium polium</i> L. growing in Algeria: chemical composition and antioxidant activity. <i>Journal of Essential Oil Research</i> , 2018 , 30, 488-497	2.3	7

178	Phytochemical analysis, antioxidant and antimicrobial activity of wild and derived plants of Hook - An endemic species from Western Ghats, India. <i>Journal of Genetic Engineering and Biotechnology</i> , 2018 , 16, 621-630	3.1	7
177	Thyme extract increases mucociliary-beating frequency in primary cell lines from chronic obstructive pulmonary disease patients. <i>Biomedicine and Pharmacotherapy</i> , 2018 , 105, 1248-1253	7.5	11
176	Not just popular spices! Essential oils from Cuminum cyminum and Pimpinella anisum are toxic to insect pests and vectors without affecting non-target invertebrates. <i>Industrial Crops and Products</i> , 2018 , 124, 236-243	5.9	47
175	Thymus lanceolatus ethanolic extract protects human cells from t-BHP induced oxidative damage. <i>Food and Function</i> , 2018 , 9, 3665-3672	6.1	10
174	The essential oil from industrial hemp (Cannabis sativa L.) by-products as an effective tool for insect pest management in organic crops. <i>Industrial Crops and Products</i> , 2018 , 122, 308-315	5.9	107
173	γ-Aminobutyric acid treatment confers decay tolerance in strawberry fruit by warranting sufficient cellular energy providing. <i>Scientia Horticulturae</i> , 2018 , 240, 249-257	4.1	17
172	New Drugs from Old Natural Compounds: Scarcely Investigated Sesquiterpenes as New Possible Therapeutic Agents. <i>Current Medicinal Chemistry</i> , 2018 , 25, 1241-1258	4.3	26
171	Sumac (Rhus coriaria L.) fruit: Essential oil variability in Iranian populations. <i>Industrial Crops and Products</i> , 2018 , 111, 1-7	5.9	37
170	Evaluation of yield, essential oil content and compositions of peppermint (Mentha piperita L.) intercropped with faba bean (Vicia faba L.). <i>Journal of Cleaner Production</i> , 2018 , 171, 529-537	10.3	30
169	Chemical composition and insecticidal activity of the essential oil from Helichrysum faradifani endemic to Madagascar. <i>Natural Product Research</i> , 2018 , 32, 1690-1698	2.3	9
168	Identification of tagitinin C from Tithonia diversifolia as antitrypanosomal compound using bioactivity-guided fractionation. <i>Fitoterapia</i> , 2018 , 124, 145-151	3.2	16
167	Evaluation of competition, essential oil quality and quantity of peppermint intercropped with soybean. <i>Industrial Crops and Products</i> , 2018 , 111, 743-754	5.9	21
166	Supercritical CO2 extraction of Rosmarinus eriocalyx growing in Algeria: Chemical composition and antioxidant activity of extracts and their solid plant materials. <i>Industrial Crops and Products</i> , 2018 , 111, 768-774	5.9	18
165	Chemical characterization of the essential oil compositions and antioxidant activity from Iranian populations of Achillea wilhelmsii K.Koch. <i>Industrial Crops and Products</i> , 2018 , 112, 274-280	5.9	24
164	Insecticidal activity of camphene, zerbombone and thumulene from Cheilocostus speciosus rhizome essential oil against the Old-World bollworm, Helicoverpa armigera. <i>Ecotoxicology and Environmental Safety</i> , 2018 , 148, 781-786	7	34
163	The crop-residue of fiber hemp cv. Futura 75: from a waste product to a source of botanical insecticides. <i>Environmental Science and Pollution Research</i> , 2018 , 25, 10515-10525	5.1	51
162	Essential Oil of Achillea ligustica (Asteraceae) as an Antifungal Agent against Phytopathogenic Fungi. <i>Natural Product Communications</i> , 2018 , 13, 1934578X1801300	0.9	2
161	Essential Oils from Aromatic and Medicinal Plants as Effective Weapons Against Mosquito Vectors of Public Health Importance. <i>Parasitology Research Monographs</i> , 2018 , 69-129	0.3	6

160	A novel herbal product based on Piper betle and Sphaeranthus indicus essential oils: Toxicity, repellent activity and impact on detoxifying enzymes GST and CYP450 of Aedes aegypti Liston (Diptera: Culicidae). <i>Journal of Asia-Pacific Entomology</i> , 2018 , 21, 1466-1472	1.4	18
159	Bioactive Constituents of Juniperus turbinata Guss. from La Maddalena Archipelago. <i>Chemistry and Biodiversity</i> , 2018 , 15, e1800148	2.5	22
158	Application of combined fertilizers improves biomass, essential oil yield, aroma profile, and antioxidant properties of Thymus daenensis Celak.. <i>Industrial Crops and Products</i> , 2018 , 121, 434-440	5.9	51
157	In vitro antioxidant activity, α -glucosidase inhibitory potential and in vivo protective effect of Asparagus stipularis Forssk aqueous extract against high-fructose diet-induced metabolic syndrome in rats. <i>Journal of Functional Foods</i> , 2018 , 47, 521-530	5.1	7
156	Essential oil composition, total phenolic and flavonoids contents, and antioxidant activity of Oliveria decumbens Vent. (Apiaceae) at different phenological stages. <i>Journal of Cleaner Production</i> , 2018 , 198, 91-95	10.3	41
155	Phenolic acids, antioxidant and antiproliferative activities of Naviglio extracts from Schizogyne sericea (Asteraceae). <i>Natural Product Research</i> , 2017 , 31, 515-522	2.3	15
154	Acute larvicidal toxicity of five essential oils (Pinus nigra, Hyssopus officinalis, Satureja montana, Aloysia citrodora and Pelargonium graveolens) against the filariasis vector Culex quinquefasciatus: Synergistic and antagonistic effects. <i>Parasitology International</i> , 2017 , 66, 166-171	2.1	98
153	Stabilization of the cyclodecadiene derivative isofuranodiene by silver (I) coordination. Mechanistic and biological aspects. <i>Fluoroplasma</i> , 2017 , 117, 52-60	3.2	9
152	An overlooked horticultural crop, Smyrniolus olusatrum, as a potential source of compounds effective against African trypanosomiasis. <i>Parasitology International</i> , 2017 , 66, 146-151	2.1	20
151	Variation in Chemical Composition and Antibacterial Activity of the Essential Oil of Wild Populations of Phlomis olivieri. <i>Chemistry and Biodiversity</i> , 2017 , 14, e1600444	2.5	6
150	Comparative toxicity of Helosciadium nodiflorum essential oils and combinations of their main constituents against the cabbage looper, Trichoplusia ni (Lepidoptera). <i>Industrial Crops and Products</i> , 2017 , 98, 46-52	5.9	45
149	Phytochemical investigations and antiproliferative secondary metabolites from Thymus alternans growing in Slovakia. <i>Pharmaceutical Biology</i> , 2017 , 55, 1162-1170	3.8	30
148	Volatile components of horsetail (Hippuris vulgaris L.) growing in central Italy. <i>Natural Product Research</i> , 2017 , 31, 2316-2320	2.3	1
147	Valorizing overlooked local crops in the era of globalization: the case of aniseed (Pimpinella anisum L.) from Castignano (central Italy). <i>Industrial Crops and Products</i> , 2017 , 104, 99-110	5.9	14
146	Effect of prolonged water stress on essential oil content, compositions and gene expression patterns of mono- and sesquiterpene synthesis in two oregano (Origanum vulgare L.) subspecies. <i>Plant Physiology and Biochemistry</i> , 2017 , 111, 119-128	5.4	91
145	Chemical variability in volatile composition between several Italian accessions of Siler montanum (S. montanum subsp. montanum and S. montanum subsp. siculum). <i>Biochemical Systematics and Ecology</i> , 2017 , 70, 14-21	1.4	3
144	Kundmannia sicula (L.) DC: a rich source of germacrene D. <i>Journal of Essential Oil Research</i> , 2017 , 29, 437-442	2.3	33
143	Antioxidant and antibacterial activities of the essential oils obtained from seven Iranian populations of Rosmarinus officinalis. <i>Industrial Crops and Products</i> , 2017 , 107, 305-311	5.9	58

142	Cytotoxic Essential Oils from <i>Eryngium campestre</i> and <i>Eryngium amethystinum</i> (Apiaceae) Growing in Central Italy. <i>Chemistry and Biodiversity</i> , 2017 , 14, e1700096	2.5	20
141	Chemical constituents, radical scavenging activity and enzyme inhibitory capacity of fruits from <i>Cotoneaster pannosus</i> Franch. <i>Food and Function</i> , 2017 , 8, 1775-1784	6.1	7
140	Not ordinary antimalarial drugs: Madagascar plant decoctions potentiating the chloroquine action against <i>Plasmodium</i> parasites. <i>Industrial Crops and Products</i> , 2017 , 103, 19-38	5.9	22
139	Chemical composition, antioxidant activity and cytotoxicity on tumour cells of the essential oil from flowers of <i>Magnolia grandiflora</i> cultivated in Iran. <i>Natural Product Research</i> , 2017 , 31, 2857-2864	2.3	14
138	Microemulsions enhance the shelf-life and processability of <i>Smyrniolum olusatrum</i> L. essential oil. <i>Flavour and Fragrance Journal</i> , 2017 , 32, 159-164	2.5	23
137	Polar Constituents, Essential Oil and Antioxidant Activity of Marsh Woundwort (<i>Stachys palustris</i> L.). <i>Chemistry and Biodiversity</i> , 2017 , 14, e1600401	2.5	22
136	Synergized mixtures of Apiaceae essential oils and related plant-borne compounds: Larvicidal effectiveness on the filariasis vector <i>Culex quinquefasciatus</i> Say. <i>Industrial Crops and Products</i> , 2017 , 96, 186-195	5.9	113
135	Isobutyrylshikonin and isovalerylshikonin from the roots of <i>Onosma visianii</i> inhibit larval growth of the tobacco cutworm <i>Spodoptera littoralis</i> . <i>Industrial Crops and Products</i> , 2017 , 109, 266-273	5.9	15
134	Efficacy of sea fennel (<i>Crithmum maritimum</i> L., Apiaceae) essential oils against <i>Culex quinquefasciatus</i> Say and <i>Spodoptera littoralis</i> (Boisd.). <i>Industrial Crops and Products</i> , 2017 , 109, 603-610	5.9	59
133	<i>Trypanosoma brucei</i> Inhibition by Essential Oils from Medicinal and Aromatic Plants Traditionally Used in Cameroon (<i>Azadirachta indica</i> , <i>Aframomum melegueta</i> , <i>Aframomum daniellii</i> , <i>Clausena anisata</i> , <i>Dichrostachys cinerea</i> and <i>Echinops giganteus</i>). <i>International Journal of Environmental Research and Public Health</i> , 2017 , 14,	4.6	13
132	Nanoparticles as effective acaricides against ticks-A review. <i>Ticks and Tick-borne Diseases</i> , 2017 , 8, 821-826	3.6	53
131	Efficacy of the Volatile Oil from Water Celery (<i>Helosciadium nodiflorum</i> , Apiaceae) against the Filariasis Vector <i>Culex quinquefasciatus</i> , the Housefly <i>Musca domestica</i> , and the African Cotton Leafworm <i>Spodoptera littoralis</i> . <i>Chemistry and Biodiversity</i> , 2017 , 14, e1700376	2.5	17
130	Toxic and repellent activity of selected monoterpenoids (thymol, carvacrol and linalool) against the castor bean tick, <i>Ixodes ricinus</i> (Acari: Ixodidae). <i>Veterinary Parasitology</i> , 2017 , 245, 86-91	2.8	81
129	Isofuranodiene and germacrone from <i>Smyrniolum olusatrum</i> essential oil as acaricides and oviposition inhibitors against <i>Tetranychus urticae</i> : impact of chemical stabilization of isofuranodiene by interaction with silver triflate. <i>Journal of Pest Science</i> , 2017 , 90, 693-699	5.5	23
128	NMR, HS-SPME-GC/MS, and HPLC/MS Analyses of Phytoconstituents and Aroma Profile of <i>Rosmarinus eriocalyx</i> . <i>Chemistry and Biodiversity</i> , 2017 , 14, e1700248	2.5	6
127	Effect of the Leaf Essential Oil from <i>Cinnamosma madagascariensis</i> Danguy on Pentylene-tetrazol-induced Seizure in Rats. <i>Chemistry and Biodiversity</i> , 2017 , 14, e1700256	2.5	8
126	Effect of <i>Rosmarinus officinalis</i> L. essential oil combined with different packaging conditions to extend the shelf life of refrigerated beef meat. <i>Food Chemistry</i> , 2017 , 221, 1069-1076	8.5	79
125	Analysis of Food Supplement with Unusual Raspberry Ketone Content. <i>Journal of Food Processing and Preservation</i> , 2017 , 41, e13019	2.1	2

124	Antimicrobial and antioxidant activity of the essential oil from the Carpathian <i>Thymus alternans</i> Klokov. <i>Natural Product Research</i> , 2017 , 31, 1121-1130	2.3	8
123	Essential Oil of <i>Thymus munbyanus</i> subsp. <i>coloratus</i> from Algeria: Chemotypification and in vitro Biological Activities. <i>Chemistry and Biodiversity</i> , 2017 , 14, e1600299	2.5	11
122	Commentary: Making Green Pesticides Greener? The Potential of Plant Products for Nanosynthesis and Pest Control. <i>Journal of Cluster Science</i> , 2017 , 28, 3-10	3	132
121	Polar constituents, protection against reactive oxygen species, and nutritional value of Chinese artichoke (<i>Stachys affinis</i> Bunge). <i>Food Chemistry</i> , 2017 , 221, 473-481	8.5	21
120	Polar extracts from the berry-like fruits of <i>Hypericum androsaemum</i> L. as a promising ingredient in skin care formulations. <i>Journal of Ethnopharmacology</i> , 2017 , 195, 255-265	5	15
119	<i>Rosmarinus eriocalyx</i> : An alternative to <i>Rosmarinus officinalis</i> as a source of antioxidant compounds. <i>Food Chemistry</i> , 2017 , 218, 78-88	8.5	31
118	Chemical composition of <i>Cinnamosma madagascariensis</i> (Cannellaceae) essential oil and its larvicidal potential against the filariasis vector <i>Culex quinquefasciatus</i> Say. <i>South African Journal of Botany</i> , 2017 , 108, 359-363	2.9	16
117	Identification of <i>Onosma visianii</i> Roots Extract and Purified Shikonin Derivatives as Potential Acaricidal Agents against <i>Tetranychus urticae</i> . <i>Molecules</i> , 2017 , 22,	4.8	14
116	Reassessment of the polar fraction of (<i>L.</i>) Benth. subsp. (Ten.) Grande (Lamiaceae) from the Monti Sibillini National Park: A potential source of bioactive compounds. <i>Journal of Intercultural Ethnopharmacology</i> , 2017 , 6, 144-153		13
115	Essential oil composition and biological activity from <i>Artemisia caerulescens</i> subsp. <i>densiflora</i> (Viv.) Gamisans ex Kerguelen & Lambinon (Asteraceae), an endemic species in the habitat of La Maddalena Archipelago. <i>Natural Product Research</i> , 2016 , 30, 1802-9	2.3	16
114	Chemical composition of the essential oil of <i>Kaliphora madagascariensis</i> Hook. f. <i>Natural Product Research</i> , 2016 , 30, 960-6	2.3	2
113	Chemical analysis of essential oils from different parts of <i>Ferula communis</i> L. growing in central Italy. <i>Natural Product Research</i> , 2016 , 30, 806-13	2.3	12
112	Reassessment of <i>Melittis melissophyllum</i> L. subsp. <i>melissophyllum</i> iridoidic fraction. <i>Natural Product Research</i> , 2016 , 30, 218-22	2.3	14
111	Isofuranodiene: A neurotoxic compound isolated from wild celery (<i>Smyrniololus atratum</i> L., Apiaceae). <i>Food Chemistry</i> , 2016 , 192, 782-7	8.5	21
110	Secondary Metabolites, Glandular Trichomes and Biological Activity of <i>Sideritis montana</i> L. subsp. <i>montana</i> from Central Italy. <i>Chemistry and Biodiversity</i> , 2016 , 13, 1380-1390	2.5	18
109	Evaluations of thyme extract effects in human normal bronchial and tracheal epithelial cell lines and in human lung cancer cell line. <i>Chemico-Biological Interactions</i> , 2016 , 256, 125-33	5	32
108	Isofuranodiene, the main volatile constituent of wild celery (<i>Smyrniololus atratum</i> L.), protects d-galactosamin/lipopopolysacchride-induced liver injury in rats. <i>Natural Product Research</i> , 2016 , 30, 1162-5	2.3	14
107	Comparative Analysis of the Volatile Profiles from Wild, Cultivated, and Commercial Roots of <i>Gentiana lutea</i> L. by Headspace Solid Phase Microextraction (HSBPME) Coupled to Gas Chromatography Mass Spectrometry (GCMS). <i>Food Analytical Methods</i> , 2016 , 9, 311-321	3.4	7

106	Chemical and biological analysis of the by-product obtained by processing <i>Gentiana lutea</i> L. and other herbs during production of bitter liqueurs. <i>Industrial Crops and Products</i> , 2016 , 80, 131-140	5.9	12
105	Diverse biological effects of the essential oil from Iranian <i>Trachyspermum ammi</i> . <i>Arabian Journal of Chemistry</i> , 2016 , 9, 775-786	5.9	68
104	Volatile profile, nutritional value and secretory structures of the berry-like fruits of <i>Hypericum androsaemum</i> L. <i>Food Research International</i> , 2016 , 79, 1-10	7	16
103	Blue honeysuckle fruit (<i>Lonicera caerulea</i> L.) from eastern Russia: phenolic composition, nutritional value and biological activities of its polar extracts. <i>Food and Function</i> , 2016 , 7, 1892-903	6.1	31
102	Phytochemical analysis of the labdanum-poor <i>Cistus creticus</i> subsp. <i>eriocephalus</i> (Viv.) Greuter et Burdet growing in central Italy. <i>Biochemical Systematics and Ecology</i> , 2016 , 66, 50-57	1.4	13
101	Mexican sunflower (<i>Tithonia diversifolia</i> , Asteraceae) volatile oil as a selective inhibitor of <i>Staphylococcus aureus</i> nicotinate mononucleotide adenyltransferase (NadD). <i>Industrial Crops and Products</i> , 2016 , 85, 181-189	5.9	19
100	Chemical analysis of the essential oils from <i>Schizogyne sericea</i> growing in different areas of Tenerife (Spain). <i>Biochemical Systematics and Ecology</i> , 2016 , 65, 192-197	1.4	5
99	A new glucosidic phthalide from <i>Helichrysum microphyllum</i> subsp. <i>tyrrhenicum</i> from La Maddalena Island (Sardinia, Italy). <i>Natural Product Research</i> , 2016 , 30, 789-95	2.3	19
98	Biological Activities of the Essential Oil from <i>Erigeron floribundus</i> . <i>Molecules</i> , 2016 , 21,	4.8	16
97	Polar Constituents and Biological Activity of the Berry-Like Fruits from <i>Hypericum androsaemum</i> L. <i>Frontiers in Plant Science</i> , 2016 , 7, 232	6.2	34
96	Bioactive Secondary Metabolites from <i>Schizogyne sericea</i> (Asteraceae) Endemic to Canary Islands. <i>Chemistry and Biodiversity</i> , 2016 , 13, 826-36	2.5	8
95	Phytochemistry, micromorphology and bioactivities of <i>Ajuga chamaepitys</i> (L.) Schreb. (Lamiaceae, Ajugoideae): Two new harpagide derivatives and an unusual iridoid glycosides pattern. <i>Phytotherapy Research</i> , 2016 , 113, 35-43	3.2	42
94	The Chemical Constituents and the Hepato-protective Effect of the Essential Oil of <i>Ferulago campestris</i> (Besser) Grecescu (Apiaceae). <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2016 , 19, 1701-1708	1.7	4
93	Evaluation of the anticonvulsant activity of the essential oil of <i>Myrothamnus moschatus</i> in convulsion induced by pentylenetetrazole and picrotoxin. <i>Asian Pacific Journal of Tropical Biomedicine</i> , 2016 , 6, 501-505	1.4	7
92	Nutritional composition, bioactive compounds and volatile profile of cocoa beans from different regions of Cameroon. <i>International Journal of Food Sciences and Nutrition</i> , 2016 , 67, 422-30	3.7	23
91	Traditional herbal remedies and dietary spices from Cameroon as novel sources of larvicides against filariasis mosquitoes?. <i>Parasitology Research</i> , 2016 , 115, 4617-4626	2.4	14
90	Methanolic extract from red berry-like fruits of <i>Hypericum androsaemum</i> : Chemical characterization and inhibitory potential of central nervous system enzymes. <i>Industrial Crops and Products</i> , 2016 , 94, 363-367	5.9	12
89	Antioxidant activity and cytotoxicity on tumour cells of the essential oil from <i>Cedronella canariensis</i> var. <i>canariensis</i> (L.) Webb & Berthel. (Lamiaceae). <i>Natural Product Research</i> , 2015 , 29, 1641-9	2.3	7

88	Chemical characterization of the essential oil compositions from Iranian populations of <i>Hypericum perforatum</i> L.. <i>Industrial Crops and Products</i> , 2015 , 76, 565-573	5.9	31
87	Antioxidant and α -glucosidase inhibitory activities of <i>Achillea tenorii</i> . <i>Pharmaceutical Biology</i> , 2015 , 53, 1505-10	3.8	35
86	Essential oil composition, polar compounds, glandular trichomes and biological activity of <i>Hyssopus officinalis</i> subsp. <i>aristatus</i> (Godr.) Nyman from central Italy. <i>Industrial Crops and Products</i> , 2015 , 77, 353-363	5.9	47
85	Evaluation of neurotogenic activity of cultivated, wild and commercial roots of <i>Gentiana lutea</i> L.. <i>Journal of Functional Foods</i> , 2015 , 19, 164-173	5.1	15
84	Phytochemical analysis and in vitro biological activity of three <i>Hypericum</i> species from the Canary Islands (<i>Hypericum reflexum</i> , <i>Hypericum canariense</i> and <i>Hypericum grandifolium</i>). <i>Flavour and Fragrance Journal</i> , 2015 , 100, 95-109	3.2	46
83	Effective clean-up and ultra high-performance liquid chromatography-tandem mass spectrometry for isoflavone determination in legumes. <i>Food Chemistry</i> , 2015 , 174, 487-94	8.5	16
82	Comparative HPLC/ESI-MS and HPLC/DAD study of different populations of cultivated, wild and commercial <i>Gentiana lutea</i> L. <i>Food Chemistry</i> , 2015 , 174, 426-33	8.5	39
81	Phytochemical Analysis, Biological Activity, and Secretory Structures of <i>Stachys annua</i> (L.) L. subsp. <i>annua</i> (Lamiaceae) from Central Italy. <i>Chemistry and Biodiversity</i> , 2015 , 12, 1172-83	2.5	23
80	Chemical composition and biological activity of the essential oil from <i>Helichrysum microphyllum</i> Cambess. ssp. <i>tyrrhenicum</i> Bacch., Brullo e Giusso growing in La Maddalena Archipelago, Sardinia. <i>Journal of Oleo Science</i> , 2015 , 64, 19-26	1.6	24
79	Secondary Metabolites of <i>Alchemilla persica</i> Growing in Iran (East Azarbaijan). <i>Natural Product Communications</i> , 2015 , 10, 1934578X1501001	0.9	0
78	Essential oil chemotypification and secretory structures of the neglected vegetable <i>Smyrniolus atratum</i> L. (Apiaceae) growing in central Italy. <i>Flavour and Fragrance Journal</i> , 2015 , 30, 139-159	2.5	37
77	Quantitative Profiling of Volatile and Phenolic Substances in the Wine Vernaccia di Serrapetrona by Development of an HS-SPME-GC-FID/MS Method and HPLC-MS. <i>Food Analytical Methods</i> , 2014 , 7, 1651-1660	3.4	16
76	Rapid Quantification of Soyasaponins I and II in Italian Lentils by High-Performance Liquid Chromatography (HPLC) Tandem Mass Spectrometry (MS/MS). <i>Food Analytical Methods</i> , 2014 , 7, 1024-1031	3.4	9
75	In vitro biological activities of seed essential oils from the Cameroonian spices <i>Afrotyrax lepidophyllus</i> MILDBR. and <i>Scorodophloeus zenkeri</i> HARMS rich in sulfur-containing compounds. <i>Chemistry and Biodiversity</i> , 2014 , 11, 161-9	2.5	22
74	Simultaneous Determination of 18 Bioactive Compounds in Italian Bitter Liqueurs by Reversed-Phase High-Performance Liquid Chromatography Diode Array Detection. <i>Food Analytical Methods</i> , 2014 , 7, 697-705	3.4	6
73	Volatile oil from striped African pepper (<i>Xylopiopsis parviflora</i> , Annonaceae) possesses notable chemopreventive, anti-inflammatory and antimicrobial potential. <i>Food Chemistry</i> , 2014 , 149, 183-9	8.5	35
72	Natural daucane esters induces apoptosis in leukaemic cells through ROS production. <i>Phytochemistry</i> , 2014 , 108, 147-56	4	18
71	HPTLC determination of chemical composition variability in raw materials used in botanicals. <i>Natural Product Research</i> , 2014 , 28, 119-26	2.3	39

70	Composition and biological activities of hogweed [<i>Heracleum sphondylium</i> L. subsp. <i>ternatum</i> (Velen.) Brummitt] essential oil and its main components octyl acetate and octyl butyrate. <i>Natural Product Research</i> , 2014 , 28, 1354-63	2.3	23
69	Volatile compounds from <i>Achillea tenorii</i> (Grande) growing in the Majella National Park (Italy). <i>Natural Product Research</i> , 2014 , 28, 1699-704	2.3	11
68	Intra-population chemical polymorphism in <i>Thymus pannonicus</i> All. growing in Slovakia. <i>Natural Product Research</i> , 2014 , 28, 1557-66	2.3	7
67	Characterization of secondary metabolites, biological activity and glandular trichomes of <i>Stachys tymphaea</i> Hausskn. from the Monti Sibillini National Park (Central Apennines, Italy). <i>Chemistry and Biodiversity</i> , 2014 , 11, 245-61	2.5	46
66	Wild celery (<i>Smyrniolum olusatrum</i> L.) oil and isofuranodiene induce apoptosis in human colon carcinoma cells. <i>Phytotherapy</i> , 2014 , 97, 133-41	3.2	42
65	Ascorbic acid content, fatty acid composition and nutritional value of the neglected vegetable Alexanders (<i>Smyrniolum olusatrum</i> L., Apiaceae). <i>Journal of Food Composition and Analysis</i> , 2014 , 35, 30-36 ^{4.1}		5
64	Antiproliferative evaluation of isofuranodiene on breast and prostate cancer cell lines. <i>Scientific World Journal, The</i> , 2014 , 2014, 264829	2.2	15
63	Quantification of caffeine, trigonelline and nicotinic acid in espresso coffee: the influence of espresso machines and coffee cultivars. <i>International Journal of Food Sciences and Nutrition</i> , 2014 , 65, 465-9	3.7	47
62	Volatile profiles of flavedo, pulp and seeds in <i>Poncirus trifoliata</i> fruits. <i>Journal of the Science of Food and Agriculture</i> , 2014 , 94, 2874-87	4.3	5
61	Biogenic amines as freshness index of meat wrapped in a new active packaging system formulated with essential oils of <i>Rosmarinus officinalis</i> . <i>International Journal of Food Sciences and Nutrition</i> , 2013 , 64, 921-8	3.7	36
60	Phytochemical analysis, biological evaluation and micromorphological study of <i>Stachys alopecuroides</i> (L.) Benth. subsp. <i>divulsa</i> (Ten.) Grande endemic to central Apennines, Italy. <i>Phytotherapy</i> , 2013 , 90, 94-103	3.2	47
59	In vitro biological activity of essential oils and isolated furanosesquiterpenes from the neglected vegetable <i>Smyrniolum olusatrum</i> L. (Apiaceae). <i>Food Chemistry</i> , 2013 , 138, 808-13	8.5	44
58	Polar constituents composition of endemic <i>Sideritis italica</i> (MILL.) GREUTER et BURTER from Central Italy. <i>Natural Product Research</i> , 2013 , 27, 1408-12	2.3	26
57	Secondary metabolites from <i>Pinus mugo</i> Turra subsp. <i>mugo</i> growing in the Majella National Park (Central Apennines, Italy). <i>Chemistry and Biodiversity</i> , 2013 , 10, 2091-100	2.5	20
56	Chemopreventive and antioxidant activity of the chamazulene-rich essential oil obtained from <i>Artemisia arborescens</i> L. growing on the Isle of La Maddalena, Sardinia, Italy. <i>Chemistry and Biodiversity</i> , 2013 , 10, 1464-74	2.5	44
55	Congruence of phytochemical and morphological profiles along an altitudinal gradient in <i>Origanum vulgare</i> ssp. <i>vulgare</i> from Venetian Region (NE Italy). <i>Chemistry and Biodiversity</i> , 2013 , 10, 569-83	2.5	21
54	Simultaneous Determination of Squalene, Tocopherol and Carotene in Table Olives by Solid Phase Extraction and High-Performance Liquid Chromatography with Diode Array Detection. <i>Food Analytical Methods</i> , 2013 , 6, 54-60	3.4	16
53	Chemical composition and in vitro biological activities of the essential oil of <i>Vepris macrophylla</i> (BAKER) I.VERD. endemic to Madagascar. <i>Chemistry and Biodiversity</i> , 2013 , 10, 356-66	2.5	22

52	Antioxidant and antiproliferative activity of <i>Hypericum hircinum</i> L. subsp. <i>majus</i> (Aiton) N. Robson essential oil. <i>Natural Product Research</i> , 2013 , 27, 862-8	2.3	63
51	Determination of soyasaponins I and II in raw and cooked legumes by solid phase extraction (SPE) coupled to liquid chromatography (LC)-mass spectrometry (MS) and assessment of their bioaccessibility by an in vitro digestion model. <i>Journal of Agricultural and Food Chemistry</i> , 2013 , 61, 1702-9	5.7	25
50	Central nervous system activities of <i>Hypericum organifolium</i> extract via GABAergic and opioidergic mechanisms. <i>Phytotherapy Research</i> , 2013 , 27, 877-84	6.7	13
49	Essential-oil polymorphism in the 'resurrection plant' <i>Myrothamnus moschatus</i> and associated ethnobotanical knowledge. <i>Chemistry and Biodiversity</i> , 2013 , 10, 1987-98	2.5	4
48	Chemical composition and biological activities of the essential oil of <i>Athanasia brownii</i> Hochr. (Asteraceae) endemic to Madagascar. <i>Chemistry and Biodiversity</i> , 2013 , 10, 1876-86	2.5	17
47	Antioxidant, Antiproliferative and Antimicrobial Activities of the Volatile Oil from the Wild Pepper <i>Piper capense</i> Used in Cameroon as a Culinary Spice. <i>Natural Product Communications</i> , 2013 , 8, 1934578X1300801	0.9	0.1
46	Characterization and biological activity of essential oils from fruits of <i>Zanthoxylum xanthoxyloides</i> Lam. and <i>Z. leprieurii</i> Guill. & Perr., two culinary plants from Cameroon. <i>Flavour and Fragrance Journal</i> , 2012 , 27, 171-179	2.5	22
45	Antimicrobial efficacy of <i>Achillea ligustica</i> All. (Asteraceae) essential oils against reference and isolated oral microorganisms. <i>Chemistry and Biodiversity</i> , 2012 , 9, 12-24	2.5	28
44	In vitro biological activities of the essential oil from the 'resurrection plant' <i>Myrothamnus moschatus</i> (Baillon) Niedenzu endemic to Madagascar. <i>Natural Product Research</i> , 2012 , 26, 2291-300	2.3	16
43	Optimization of espresso machine parameters through the analysis of coffee odorants by HS-SPME-GC/MS. <i>Food Chemistry</i> , 2012 , 135, 1127-33	8.5	59
42	A forgotten vegetable (<i>Smyrniololus atrum</i> L., Apiaceae) as a rich source of isofuranodiene. <i>Food Chemistry</i> , 2012 , 135, 2852-62	8.5	39
41	Comparative study of aroma profile and phenolic content of Montepulciano monovarietal red wines from the Marche and Abruzzo regions of Italy using HS-SPME-GC-MS and HPLC-MS. <i>Food Chemistry</i> , 2012 , 132, 1592-1599	8.5	61
40	Phytochemical investigation of the essential oil from the resurrection plant <i>Myrothamnus moschatus</i> (Baillon) Niedenzu endemic to Madagascar. <i>Journal of Essential Oil Research</i> , 2012 , 24, 299-304	2.3	2
39	Analysis of the volatile compounds of <i>Teucrium flavum</i> L. subsp. <i>flavum</i> (Lamiaceae) by headspace solid-phase microextraction coupled to gas chromatography with flame ionisation and mass spectrometric detection. <i>Natural Product Research</i> , 2012 , 26, 1339-47	2.3	6
38	Gas chromatography for the characterization of the mushroom-like flavor in <i>Melittis melissophyllum</i> L. (Lamiaceae). <i>Journal of Essential Oil Research</i> , 2012 , 24, 321-337	2.3	13
37	Effects of treatment with St. John's Wort on blood glucose levels and pain perceptions of streptozotocin-diabetic rats. <i>Phytotherapy Research</i> , 2011 , 25, 576-84	3.2	27
36	HPLC quantification of coumarin in bastard balm (<i>Melittis melissophyllum</i> L., Lamiaceae). <i>Phytotherapy Research</i> , 2011 , 25, 1215-21	3.2	30
35	Natural daucane sesquiterpenes with antiproliferative and proapoptotic activity against human tumor cells. <i>Bioorganic and Medicinal Chemistry</i> , 2011 , 19, 5876-85	3.4	30

34	Molecular mediators involved in <i>Ferulago campestris</i> essential oil effects on osteoblast metabolism. <i>Journal of Cellular Biochemistry</i> , 2011 , 112, 3742-54	4.7	10
33	Solid-phase microextraction (SPME) analysis of six Italian populations of <i>Ephedra nebrodensis</i> Tineo ex Guss. subsp. <i>nebrodensis</i> . <i>Chemistry and Biodiversity</i> , 2011 , 8, 95-114	2.5	
32	Chemical differences in volatiles between <i>Melittis melissophyllum</i> L. subsp. <i>melissophyllum</i> and subsp. <i>albida</i> (Guss) P. W. Ball (Lamiaceae) determined by solid-phase microextraction (SPME) coupled with GC/FID and GC/MS. <i>Chemistry and Biodiversity</i> , 2011 , 8, 325-43	2.5	14
31	Glandular trichomes and essential oil composition of endemic <i>Sideritis italica</i> (Mill.) Greuter et Burdet from central Italy. <i>Chemistry and Biodiversity</i> , 2011 , 8, 2179-94	2.5	10
30	Volatile components of whole and different plant parts of bastard balm (<i>Melittis melissophyllum</i> L., Lamiaceae) collected in Central Italy and Slovakia. <i>Chemistry and Biodiversity</i> , 2011 , 8, 2057-79	2.5	15
29	New antidepressant drug candidate: <i>Hypericum montbretti</i> extract. <i>Natural Product Research</i> , 2011 , 25, 1469-72	2.3	11
28	Evaluation of the wound healing potentials of two subspecies of <i>Hypericum perforatum</i> on cultured NIH3T3 fibroblasts. <i>Phytotherapy Research</i> , 2011 , 25, 208-14	6.7	29
27	Qualitative Analysis of the Smoke-Stream of Different Kinds of Incense by SPME/GC-MS. <i>Natural Product Communications</i> , 2010 , 5, 1934578X1000500	0.9	
26	Essential Oil Composition of <i>Ephedra nebrodensis</i> Tineo ex Guss. subsp. <i>nebrodensis</i> from Central Italy. <i>Journal of Essential Oil Research</i> , 2010 , 22, 354-357	2.3	3
25	Chemical Composition and Seasonal Variation of <i>Hypericum hircinum</i> L. subsp. <i>majus</i> (Aiton) N. Robson Essential Oil. <i>Journal of Essential Oil Research</i> , 2010 , 22, 434-443	2.3	3
24	Chemical composition and antimicrobial activity of <i>Hypericum hircinum</i> L. Subsp. <i>majus</i> essential oil. <i>Chemistry of Natural Compounds</i> , 2010 , 46, 125-129	0.7	8
23	Identification of non-alkaloid acetylcholinesterase inhibitors from <i>Ferulago campestris</i> (Besser) Grecescu (Apiaceae). <i>Phytotherapy</i> 2010 , 81, 1208-12	3.2	40
22	Histochemical localization of secretion and composition of the essential oil in <i>Melittis melissophyllum</i> L. subsp. <i>melissophyllum</i> from Central Italy. <i>Flavour and Fragrance Journal</i> , 2010 , 25, 63-70	2.5	28
21	Essential oil from fruits and roots of <i>Ferulago campestris</i> (Besser) Grecescu (Apiaceae): composition and antioxidant and anti- <i>Candida</i> activity. <i>Flavour and Fragrance Journal</i> , 2010 , 25, 493-502	2.5	24
20	Chemical composition and antimicrobial activity of the essential oils from several <i>Hypericum</i> taxa (Guttiferae) growing in central Italy (Appennino Umbro-Marchigiano). <i>Chemistry and Biodiversity</i> , 2010 , 7, 447-66	2.5	39
19	Characterisation of the mushroom-like flavour of <i>Melittis melissophyllum</i> L. subsp. <i>melissophyllum</i> by headspace solid-phase microextraction (HS-SPME) coupled with gas chromatography (GC/FID) and gas chromatography-mass spectrometry (GC/MS). <i>Food Chemistry</i> , 2010 , 123, 983-992	8.5	35
18	Chemical composition and antimicrobial activity of the essential oil from <i>Ferula glauca</i> L. (F. <i>communis</i> L. subsp. <i>glauca</i>) growing in Marche (central Italy). <i>Phytotherapy</i> 2009 , 80, 68-72	3.2	60
17	Composition and biological activity of essential oil of <i>Achillea ligustica</i> All. (Asteraceae) naturalized in central Italy: ideal candidate for anti-cariogenic formulations. <i>Phytotherapy</i> 2009 , 80, 313-9	3.2	45

16	Chemical analysis of the essential oil of <i>Ferula glauca</i> L. (Apiaceae) growing in Marche (central Italy). <i>Biochemical Systematics and Ecology</i> , 2009 , 37, 432-441	1.4	11
15	Chemical composition and antimicrobial activity of the essential oil of <i>Ferulago campestris</i> (Besser) Grecescu growing in central Italy. <i>Flavour and Fragrance Journal</i> , 2009 , 24, 309-315	2.5	17
14	Comparison of the characterisation of the fruit-like aroma of <i>Teucrium flavum</i> L. subsp. <i>flavum</i> by hydrodistillation and solid-phase micro-extraction. <i>Journal of the Science of Food and Agriculture</i> , 2009 , 89, 2505-2518	4.3	17
13	<i>Melittis melissophyllum</i> L. subsp. <i>melissophyllum</i> (Lamiaceae) from central Italy: A new source of a mushroom-like flavour. <i>Food Chemistry</i> , 2009 , 113, 216-221	8.5	22
12	Analysis of the Volatile Components of <i>Onosma echioides</i> (L.) L. var. <i>columnae</i> Lacaita Growing in Central Italy. <i>Journal of Essential Oil Research</i> , 2009 , 21, 441-447	2.3	23
11	Quantification of soyasaponins I and betag in Italian lentil seeds by solid-phase extraction (SPE) and high-performance liquid chromatography-mass spectrometry (HPLC-MS). <i>Journal of Agricultural and Food Chemistry</i> , 2009 , 57, 11226-33	5.7	22
10	Phytochemical and antioxidant analysis of eight <i>Hypericum</i> taxa from Central Italy. <i>Phytotherapy</i> 2008 , 79, 210-3	3.2	48
9	Essential Oil Composition of <i>Hypericum</i> <i>hidcotei</i> <i>Journal of Essential Oil Research</i> , 2008 , 20, 539-541	2.3	2
8	Alkannin/shikonin mixture from roots of <i>Onosma echioides</i> (L.) L.: extraction method study and quantification. <i>Journal of Separation Science</i> , 2008 , 31, 945-52	3.4	26
7	Palynological analysis of five selected <i>Onosma</i> taxa. <i>Biologia (Poland)</i> , 2008 , 63, 183-186	1.5	7
6	Antimicrobial activity of seven hypericum entities from central Italy. <i>Planta Medica</i> , 2007 , 73, 564-6	3.1	46
5	Essential oil composition of <i>Hypericum richeri</i> Vill. from Italy. <i>Flavour and Fragrance Journal</i> , 2005 , 20, 295-298	2.5	22
4	Morphological, histochemical and phytochemical investigation of the genus <i>Hypericum</i> of the Central Italy. <i>Phytotherapy</i> 2004 , 75, 702-11	3.2	51
3	Visual and olfactory preferences of <i>Frankliniella occidentalis</i> (Thysanoptera: Thripidae) for color and volatiles of different <i>Rosa chinensis</i> (Rosales: Rosaceae) cultivars. <i>Oriental Insects</i> , 1-17	0.3	0
2	Antiproliferative, antimicrobial and antioxidant properties of <i>Cedrus libani</i> and <i>Pinus pinea</i> wood oils and <i>Juniperus excelsa</i> berry oil. <i>Plant Biosystems</i> , 1-12	1.6	5
1	Variability in chemical composition and antibacterial activity of <i>Salvia majdae</i> essential oil under various extraction techniques. <i>Journal of Essential Oil Research</i> , 1-11	2.3	0