List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7841822/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Rapid identification and quantification of methamphetamine and amphetamine in hair by gas chromatography/mass spectrometry coupled with micropulverized extraction, aqueous acetylation and microextraction by packed sorbent. Journal of Chromatography A, 2009, 1216, 4063-4070. | 3.7 | 87 |
| 2 | Morphological and chemical analysis of magic mushrooms in Japan. Forensic Science International, 2003, 138, 85-90. | 2.2 | 58 |
| 3 | Chiral analysis of amphetamine-type stimulants using reversed-polarity capillary electrophoresis/positive ion electrospray ionization tandem mass spectrometry. Electrophoresis, 2003, 24, 1770-1776. | 2.4 | 58 |
| 4 | <i>In vitro</i> stability and metabolism of salvinorin A in rat plasma. Xenobiotica, 2009, 39, 391-398. | 1.1 | 55 |
| 5 | Degradation pathways of 4-methylmethcathinone in alkaline solution and stability of methcathinone analogs in various pH solutions. Forensic Science International, 2012, 220, 103-110. | 2.2 | 53 |
| 6 | A method for screening for various sedative-hypnotics in serum by liquid chromatography/single quadrupole mass spectrometry. Forensic Science International, 2006, 157, 57-70. | 2.2 | 48 |
| 7 | Methamphetamine impurity profiling using a 0.32 mm i.d. nonpolar capillary column. Forensic Science International, 2003, 135, 42-47. | 2.2 | 44 |
| 8 | Analysis of hallucinogenic constituents in Amanita mushrooms circulated in Japan. Forensic Science International, 2006, 164, 172-178. | 2.2 | 42 |
| 9 | Determination of muscimol and ibotenic acid in Amanita mushrooms by high-performance liquid chromatography and liquid chromatography-tandem mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2007, 852, 430-435. | 2.3 | 40 |
| 10 | Forensic application of chiral separation of amphetamine-type stimulants to impurity analysis of seized methamphetamine by capillary electrophoresis. Forensic Science International, 2006, 161, 92-96. | 2.2 | 39 |
| 11 | The use of a sulfonated capillary on chiral capillary electrophoresis/mass spectrometry of amphetamine-type stimulants for methamphetamine impurity profiling. Forensic Science International, 2015, 249, 59-65. | 2.2 | 39 |
| 12 | Time-course measurements of caffeine and its metabolites extracted from fingertips after coffee intake: a preliminary study for the detection of drugs from fingerprints. Analytical and Bioanalytical Chemistry, 2013, 405, 3945-3952. | 3.7 | 35 |
| 13 | Thermal degradation of a new synthetic cannabinoid QUPIC during analysis by gas chromatography–mass spectrometry. Forensic Toxicology, 2014, 32, 201-207. | 2.4 | 35 |
| 14 | Comparison and classification of methamphetamine seized in Japan and Thailand using gas chromatography with liquid–liquid extraction and solid-phase microextraction. Forensic Science International, 2008, 175, 85-92. | 2.2 | 33 |
| 15 | Three-step drug extraction from a single sub-millimeter segment of hair and nail to determine the exact day of drug intake. Analytica Chimica Acta, 2016, 948, 40-47. | 5.4 | 33 |
| 16 | Chemical profiling of seized methamphetamine putatively synthesized from phenylacetic acid derivatives. Forensic Science International, 2013, 227, 42-44. | 2.2 | 32 |
| 17 | Differentiation of regioisomeric fluoroamphetamine analogs by gas chromatography–mass spectrometry and liquid chromatography–tandem mass spectrometry. Forensic Toxicology, 2013, 31, 241-250. | 2.4 | 32 |
| 18 | ldentification of impurities and the statistical classification of methamphetamine using headspace solid phase microextraction and gas chromatography–mass spectrometry. Forensic Science International. 2006. 160. 44-52. | 2.2 | 29 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Simple and simultaneous detection of methamphetamine and dimethyl sulfone in crystalline methamphetamine seizures by fast gas chromatography. Forensic Toxicology, 2008, 26, 19-22. | 2.4 | 29 |
| 20 | Micro-segmental hair analysis for proving drug-facilitated crimes: Evidence that a victim ingested a sleeping aid, diphenhydramine, on a specific day. Forensic Science International, 2018, 288, 23-28. | 2.2 | 29 |
| 21 | Metabolism of Fentanyl and Acetylfentanyl in Human-Induced Pluripotent Stem Cell-Derived Hepatocytes. Biological and Pharmaceutical Bulletin, 2018, 41, 106-114. | 1.4 | 29 |
| 22 | Application of a portable near infrared spectrometer for presumptive identification of psychoactive drugs. Forensic Science International, 2014, 242, 162-171. | 2.2 | 28 |
| 23 | Rapid, simple, and highly sensitive analysis of drugs in biological samples using thin-layer chromatography coupled with matrix-assisted laser desorption/ionization mass spectrometry. Analytical and Bioanalytical Chemistry, 2012, 402, 1257-1267. | 3.7 | 27 |
| 24 | Excretory Profile of 4-Bromo-2,5-dimethoxyphenethylamine (2C-B) in Rat Journal of Health Science, 2003, 49, 166-169. | 0.9 | 25 |
| 25 | Determination of salvinorin A and salvinorin B in Salvia divinorum-related products circulated in Japan. Forensic Science International, 2008, 180, 105-109. | 2.2 | 25 |
| 26 | Uptake of 3,4-methylenedioxymethamphetamine and its related compounds by a proton-coupled transport system in Caco-2 cells. Biochimica Et Biophysica Acta - Biomembranes, 2008, 1778, 42-50. | 2.6 | 25 |
| 27 | Profiling of seized methamphetamine putatively synthesized by reductive amination of 1-phenyl-2-propanone. Forensic Toxicology, 2012, 30, 70-75. | 2.4 | 25 |
| 28 | Thermal degradation of α-pyrrolidinopentiophenone during injection in gas chromatography/mass spectrometry. Forensic Science International, 2013, 231, 296-299. | 2.2 | 25 |
| 29 | Effectiveness of saliva and fingerprints as alternative specimens to urine and blood in forensic drug testing. Drug Testing and Analysis, 2016, 8, 644-651. | 2.6 | 25 |
| 30 | Time ourse measurements of drug concentrations in hair and toenails after single administrations of pharmaceutical products. Drug Testing and Analysis, 2017, 9, 571-577. | 2.6 | 25 |
| 31 | Use of hepatocytes isolated from a liver-humanized mouse for studies on the metabolism of drugs: application to the metabolism of fentanyl and acetylfentanyl. Forensic Toxicology, 2018, 36, 467-475. | 2.4 | 25 |
| 32 | Potentiation of domperidone-induced catalepsy by a P-glycoprotein inhibitor, cyclosporin A. Biopharmaceutics and Drug Disposition, 2003, 24, 105-114. | 1.9 | 24 |
| 33 | Rapid detection of hypnotics using surface-enhanced Raman scattering based on gold nanoparticle co-aggregation in a wet system. Analyst, The, 2019, 144, 2158-2165. | 3.5 | 23 |
| 34 | Micro-pulverized extraction pretreatment for highly sensitive analysis of 11-nor-9-carboxy-î" ⁹ -tetrahydrocannabinol in hair by liquid chromatography/tandem mass spectrometry. Rapid Communications in Mass Spectrometry, 2015, 29, 2158-2166. | 1.5 | 22 |
| 35 | Enantioseparation of methamphetamine by supercritical fluid chromatography with cellulose-based packed column. Forensic Science International, 2017, 273, 39-44. | 2.2 | 22 |
| 36 | Strong evidence of drug-facilitated crimes by hair analysis using LC–MS/MS after micro-segmentation. Forensic Toxicology, 2019, 37, 480-487. | 2.4 | 22 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Detection of main metabolites of XLRâ€11 and its thermal degradation product in human hepatoma HepaRG cells and human urine. Drug Testing and Analysis, 2015, 7, 341-345. | 2.6 | 21 |
| 38 | Analysis of amphetamine-type stimulants and their metabolites in plasma, urine and bile by liquid chromatography with a strong cation-exchange column-tandem mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2008, 867, 78-83. | 2.3 | 20 |
| 39 | Development of an on-site screening system for amphetamine-type stimulant tablets with a portable attenuated total reflection Fourier transform infrared spectrometer. Analytica Chimica Acta, 2008, 608, 95-103. | 5.4 | 20 |
| 40 | Distribution measurements of 3,4-methylenedioxymethamphetamine and its metabolites in organs by matrix-assisted laser desorption/ionization imaging mass spectrometry using an automatic matrix spraying system with an air brush and a turntable. Analytical and Bioanalytical Chemistry, 2012, 404, 1823-1830. | 3.7 | 20 |
| 41 | Evaluation method for linking methamphetamine seizures using stable carbon and nitrogen isotopic compositions: a complementary study with impurity profiling. Rapid Communications in Mass Spectrometry, 2008, 22, 3816-3822. | 1.5 | 19 |
| 42 | Approaching over 10Â000â€fold sensitivity increase in chiral capillary electrophoresis: Cationâ€selective exhaustive injection and sweeping cyclodextrinâ€modified micellar electrokinetic chromatography. Electrophoresis, 2016, 37, 2970-2976. | 2.4 | 19 |
| 43 | Characterization and Differentiation of Geometric Isomers of 3â€methylfentanyl Analogs by Gas Chromatography/Mass Spectrometry, Liquid Chromatography/Mass Spectrometry, and Nuclear Magnetic Resonance Spectroscopy. Journal of Forensic Sciences, 2017, 62, 1472-1478. | 1.6 | 19 |
| 44 | Different localizations of drugs simultaneously administered in a strand of hair by microâ€segmental analysis. Drug Testing and Analysis, 2018, 10, 750-760. | 2.6 | 19 |
| 45 | In Vivo Metabolism of 5-Methoxy-N,N-diisopropyltryptamine in Rat. Journal of Health Science, 2006, 52, 425-430. | 0.9 | 18 |
| 46 | Applicability of chemically modified capillaries in chiral capillary electrophoresis for methamphetamine profiling. Forensic Science International, 2013, 226, 235-239. | 2.2 | 18 |
| 47 | Analysis of 4â€Bromoâ€2,5â€Dimethoxyphenethylamine <scp>A</scp> buser's Urine: Identification and Quantitation of Urinary Metabolites. Journal of Forensic Sciences, 2013, 58, 279-287. | 1.6 | 18 |
| 48 | Time-course measurements of drugs and metabolites transferred from fingertips after drug administration: usefulness of fingerprints for drug testing. Forensic Toxicology, 2014, 32, 235-242. | 2.4 | 18 |
| 49 | Differentiation of regioisomeric chloroamphetamine analogs using gas chromatography–chemical ionization-tandem mass spectrometry. Forensic Toxicology, 2015, 33, 338-347. | 2.4 | 18 |
| 50 | Identification and differentiation of methcathinone analogs by gas chromatographyâ€mass spectrometry. Drug Testing and Analysis, 2013, 5, 670-677. | 2.6 | 17 |
| 51 | Metabolism of Butyrylfentanyl in Fresh Human Hepatocytes: Chemical Synthesis of Authentic Metabolite Standards for Definitive Identification. Biological and Pharmaceutical Bulletin, 2019, 42, 623-630. | 1.4 | 17 |
| 52 | Protease-Sensitive Urinary Pheromones Induce Region-Specific Fos-Expression in Rat Accessory Olfactory Bulb. Biochemical and Biophysical Research Communications, 1999, 260, 222-224. | 2.1 | 16 |
| 53 | Development of a new field-test procedure for cocaine. Forensic Science International, 2017, 270, 267-274. | 2.2 | 16 |
| 54 | Accurate Estimation of Drug Intake Day by Microsegmental Analysis of a Strand of Hair by Use of Internal Temporal Markers. journal of applied laboratory medicine, The, 2018, 3, 37-47. | 1.3 | 16 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Development of simple and accurate detection systems for Cannabis sativa using DNA chromatography. Forensic Science International, 2018, 291, 68-75. | 2.2 | 16 |
| 56 | A Fatal Case of Suspected Anaphylaxis with Cefoperazone and Sulbactam: LCâ€MS Analysis. Journal of Forensic Sciences, 2008, 53, 226-231. | 1.6 | 15 |
| 57 | Thermal desorption counterâ€flow introduction atmospheric pressure chemical ionization for direct mass spectrometry of ecstasy tablets. Journal of Mass Spectrometry, 2009, 44, 1300-1307. | 1.6 | 14 |
| 58 | Recreational drugs, 3,4-Methylenedioxymethamphetamine(MDMA), 3,4-methylenedioxyamphetamine (MDA) and diphenylprolinol, inhibit neurite outgrowth in PC12 cells. Journal of Toxicological Sciences, 2010, 35, 375-381. | 1.5 | 14 |
| 59 | Differentiation of ringâ€substituted regioisomers of amphetamine and methamphetamine by supercritical fluid chromatography. Drug Testing and Analysis, 2017, 9, 389-398. | 2.6 | 14 |
| 60 | Simultaneous chiral impurity analysis of methamphetamine and its precursors by supercritical fluid chromatography–tandem mass spectrometry. Forensic Toxicology, 2019, 37, 145-153. | 2.4 | 14 |
| 61 | In vivometabolism of 2,5-dimethoxy-4-propylthiophenethylamine in rat. Xenobiotica, 2007, 37, 679-692. | 1.1 | 13 |
| 62 | Urinary Excretion Profiles of Two Major Triazolam Metabolites α-Hydroxytriazolam and 4-Hydroxytnazolam. Journal of Analytical Toxicology, 2005, 29, 240-243. | 2.8 | 12 |
| 63 | Interactions between 3,4-methylenedioxymethamphetamine, methamphetamine, ketamine, and caffeine in human intestinal Caco-2 cells and in oral administration to rats. Forensic Science International, 2007, 170, 183-188. | 2.2 | 12 |
| 64 | Seized methamphetamine samples with unique profiles of stable nitrogen isotopic composition documented by stable isotope ratio mass spectrometry. Forensic Toxicology, 2010, 28, 119-123. | 2.4 | 12 |
| 65 | Development of a novel immunoassay for herbal cannabis using a new fluorescent antibody probe, "Ultra Quenchbody― Forensic Science International, 2016, 266, 541-548. | 2.2 | 12 |
| 66 | Rapid detection of synthetic cannabinoids in herbal highs using surface-enhanced Raman scattering produced by gold nanoparticle co-aggregation in a wet system. Analyst, The, 2019, 144, 6928-6935. | 3.5 | 12 |
| 67 | Estimation of day of death using micro-segmental hair analysis based on drug use history: a case of lidocaine use as a marker. International Journal of Legal Medicine, 2019, 133, 117-122. | 2.2 | 12 |
| 68 | Micro-segmental hair analysis: detailed procedures and applications in forensic toxicology. Forensic Toxicology, 2022, 40, 215-233. | 2.4 | 12 |
| 69 | Contribution of thermal desorption and liquid–liquid extraction for identification and profiling of impurities in methamphetamine by gas chromatography–mass spectrometry. Forensic Science International, 2007, 171, 9-15. | 2.2 | 11 |
| 70 | Interaction of 3,4â€Methylenedioxymethamphetamine and Methamphetamine During Metabolism by <i>In Vitro</i> Human Metabolic Enzymes and in Rats*. Journal of Forensic Sciences, 2012, 57, 1008-1013. | 1.6 | 11 |
| 71 | Development of a Library Searchâ€Based Screening System for 3,4â€Methylenedioxymethamphetamine in Ecstasy Tablets Using a Portable Nearâ€Infrared Spectrometer. Journal of Forensic Sciences, 2016, 61, 1208-1214. | 1.6 | 11 |
| 72 | Differentiation of ring-substituted bromoamphetamine analogs by gas chromatography–tandem mass spectrometry. Forensic Toxicology, 2016, 34, 125-132. | 2.4 | 10 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Analysis of Benzylpiperazine-like Compounds. Japanese Journal of Science and Technology for Identification, 2004, 9, 165-184. | 0.2 | 9 |
| 74 | Measurement of three-dimensional distributions of drugs in nails using liquid chromatography/tandem mass spectrometry after micro-segmentation to elucidate drug uptake routes. Analytica Chimica Acta, 2020, 1108, 89-97. | 5.4 | 9 |
| 75 | In vivometabolism ofα-methyltryptamine in rats: Identification of urinary metabolites. Xenobiotica, 2008, 38, 1476-1486. | 1.1 | 8 |
| 76 | Synthesis and Identification of Urinary Metabolites of 4-Iodo-2,5-dimethoxyphenethylamine. Journal of Forensic Sciences, 2011, 56, 1319-1323. | 1.6 | 8 |
| 77 | Distribution measurement of amphetamineâ€type stimulants in organs using micropulverized extraction and liquid chromatography/tandem mass spectrometry to complement drug distribution using mass spectrometry imaging. Rapid Communications in Mass Spectrometry, 2011, 25, 2397-2406. | 1.5 | 8 |
| 78 | Phosgene in deteriorated chloroform: presumptive cause of production of 3,4-dimethyl-5-phenyl-2-oxazolidones in methamphetamine. Forensic Toxicology, 2020, 38, 475-480. | 2.4 | 8 |
| 79 | Studies on the phase I metabolites of the new designer drug 1-(2,3-dihydro-1H-inden-5-yl)-2-(pyrrolidine-1-yl)butan-1-one (5-PPDI) in human urine. Forensic Science International, 2020, 310, 110214. | 2.2 | 8 |
| 80 | Stereoselective analysis of ephedrine and its stereoisomers as impurities and/or by-products in seized methamphetamine by supercritical fluid chromatography/tandem mass spectrometry. Forensic Science International, 2021, 318, 110591. | 2.2 | 8 |
| 81 | Development of an improved method to estimate the days of continuous drug ingestion, based on the microâ€segmental hair analysis. Drug Testing and Analysis, 2021, 13, 1295-1304. | 2.6 | 8 |
| 82 | Thermal decomposition of CBD to Δ9-THC during GC-MS analysis: A potential cause of Δ9-THC misidentification. Forensic Science International, 2022, 337, 111366. | 2.2 | 8 |
| 83 | Determination of 4-Hydroxy-3-methoxymethamphetamine as a Metabolite of Methamphetamine in Rats and Human Liver Microsomes Using Gas Chromatography-Mass Spectrometry and Liquid Chromatography-Tandem Mass Spectrometry. Journal of Analytical Toxicology, 2009, 33, 266-271. | 2.8 | 7 |
| 84 | Degradation of N-hydroxy-3,4-methylenedioxymethamphetamine in aqueous solution and its prevention. Forensic Science International, 2009, 193, 106-111. | 2.2 | 7 |
| 85 | Simultaneous determination of tryptamine analogues in designer drugs using gas chromatography–mass spectrometry and liquid chromatography–tandem mass spectrometry. Forensic Toxicology, 2014, 32, 154-161. | 2.4 | 7 |
| 86 | Utilization of matrix-assisted laser desorption/ionization imaging mass spectrometry to search for cannabis in herb mixtures. Analytical and Bioanalytical Chemistry, 2014, 406, 4789-4794. | 3.7 | 7 |
| 87 | Highly sensitive quantification of unconjugated 11-nor-9-carboxy-î"9-tetrahydrocannabinol in a cannabis user's hair using micropulverized extraction. Forensic Science International, 2016, 262, e34-e36. | 2.2 | 7 |
| 88 | Metabolism of a new synthetic opioid tetrahydrofuranylfentanyl in fresh isolated human hepatocytes: Detection and confirmation of ringâ€opened metabolites. Drug Testing and Analysis, 2020, 12, 439-448. | 2.6 | 7 |
| 89 | Development of the "selective concentration―analytical method for drug-containing hair regions based on micro-segmental analysis to identify a trace amount of drug in hair: hair analysis following single-dose ingestion of midazolam. Forensic Toxicology, 2021, 39, 156-166. | 2.4 | 7 |
| 90 | Rapid identification of drug-type and fiber-type cannabis by allele specific duplex PCR. Forensic Science International, 2021, 318, 110634. | 2.2 | 6 |

| # | Article | IF | CITATIONS |
|-----|---|-------------------|---------------------|
| 91 | Distribution profiles of diphenhydramine and lidocaine in scalp, axillary, and pubic hairs measured by micro-segmental hair analysis: good indicator for discrimination between administration and external contamination of the drugs. Forensic Toxicology, 2022, 40, 64-74. | 2.4 | 6 |
| 92 | Thin-layer chromatography on silver nitrate-impregnated silica gel for analysis of homemade tetrahydrocannabinol mixtures. Forensic Toxicology, 2022, 40, 125-131. | 2.4 | 6 |
| 93 | Differentiation of regioisomeric methylamphetamines by GC/MS. Japanese Journal of Forensic Science and Technology, 2014, 19, 111-119. | 0.1 | 5 |
| 94 | Comments on "Characterization of four new designer drugs, 5-chloro-NNEI, NNEI indazole analog, α-PHPP and α-POP, with 11 newly distributed designer drugs in illegal products― Forensic Science International, 2015, 251, e15-e17. | 2.2 | 5 |
| 95 | Instability of the hydrochloride salts of cathinone derivatives in air. Forensic Science International, 2015, 248, 48-54. | 2.2 | 5 |
| 96 | Development of rapid and simple method for DNA extraction from cannabis resin based on the evaluation of relative PCR amplification ability. Forensic Science International, 2018, 287, 176-182. | 2.2 | 5 |
| 97 | Evaluation of Agonistic Activity of Fluorinated and Nonfluorinated Fentanyl Analogs on μ-Opioid Receptor Using a Cell-Based Assay System. Biological and Pharmaceutical Bulletin, 2021, 44, 159-161. | 1.4 | 5 |
| 98 | Agonistic activity of fentanyl analogs and their metabolites on opioid receptors. Forensic Toxicology, 2022, 40, 156-162. | 2.4 | 5 |
| 99 | Detection and confirmation of the ring-opened carboxylic acid metabolite of a new synthetic opioid furanylfentanyl. Forensic Toxicology, 2021, 39, 114-122. | 2.4 | 4 |
| 100 | Analysis of potential phenylacetone precursors (ethyl 3â€oxoâ€2â€phenylbutyrate, methyl) Tj ETQqO 0 0 rgBT / their conversion to phenylacetone. Drug Testing and Analysis, 2021, , . | Overlock 1 2.6 | .0 Tf 50 387 T 4 |
| 101 | Title is missing!. Japanese Journal of Science and Technology for Identification, 2004, 9, 71-78. | 0.2 | 4 |
| 102 | Optimized Conditions for the Enzymatic Hydrolysis of .ALPHAHydroxytriazolam-Glucuronide in Human Urine. Journal of Health Science, 2004, 50, 286-289. | 0.9 | 3 |
| 103 | Increase in split ratio enables detection of underivatized N-hydroxy-3,4-methylenedioxymethamphetamine and N-hydroxy-3,4-methylenedioxyamphetamine by capillary GC-MS. Forensic Toxicology, 2010, 28, 55-57. | 2.4 | 3 |
| 104 | A model system for prediction of the in vivo metabolism of designer drugs using three-dimensional culture of rat and human hepatocytes. Forensic Toxicology, 2011, 29, 142-151. | 2.4 | 3 |
| 105 | Rapid Chemical Examinations of Cannabis and Its Related Herbal Products. Japanese Journal of Forensic Science and Technology, 2013, 18, 143-153. | 0.1 | 3 |
| 106 | Profiling of Methamphetamine. Bunseki Kagaku, 2014, 63, 221-231. | 0.2 | 3 |
| 107 | Synthesis and Analysis of Glucuronic Acidâ€Conjugated Metabolites of 4â€Bromoâ€2,5â€Dimethoxyphenethylamine. Journal of Forensic Sciences, 2017, 62, 488-492. | 1.6 | 3 |
| 108 | DNA testing of suspected cannabis samples with exceptional morphology using a simple detection kit. Forensic Toxicology, 2021, 39, 266-274. | 2.4 | 3 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Degradation of 1-phenyl-2-propanone during long-term storage: useful information for methamphetamine impurity profiling. Forensic Toxicology, 2021, 39, 405-416. | 2.4 | 3 |
| 110 | Formation of Oxazolidine Derivatives by Reaction with Ephedrines and Aldehyde Impurities in Ethyl Acetate. Journal of Chromatographic Science, 2022, 60, 316-323. | 1.4 | 3 |
| 111 | Differentiation of ringâ€substituted regioisomers of cathinone analogs by supercritical fluid chromatography. Analytical Science Advances, 2020, 1, 22. | 2.8 | 2 |
| 112 | Effects of the Various Preparation Procedures of Dragendorff Reagent on Sensitivity for Thin Layer Chromatography. Japanese Journal of Forensic Science and Technology, 2005, 10, 127-133. | 0.1 | 1 |
| 113 | Urinary Excretion Profiles of 2,5-Dimethoxy-4-alkylthiophenethylamine Analogs in Rats. Biological and Pharmaceutical Bulletin, 2016, 39, 883-886. | 1.4 | 1 |
| 114 | Evaluation of drug identification and discrimination ability of portable spectrometers. Japanese Journal of Forensic Science and Technology, 2017, 22, 9-24. | 0.1 | 1 |
| 115 | Development and demonstration of cannabis DNA detection kit using DNA chromatography chip. Japanese Journal of Forensic Science and Technology, 2021, 26, 29-48. | 0.1 | 1 |
| 116 | Identification of the metabolites of 2,5-dimethoxy-4-ethylthiophenethylamine (2C-T-2) and 2,5-dimethoxy-4-isopropylthiophenethylamine (2C-T-4) in rat urine. Japanese Journal of Forensic Science and Technology, 2014, 19, 91-101. | 0.1 | 1 |
| 117 | Urinary excretion profiles ofN-hydroxy-3,4-methylenedioxymethamphetamine in rats. Xenobiotica, 2011, 41, 578-584. | 1.1 | Ο |
| 118 | Comparison and evaluation of the quick purification methods of methamphetamine hydrochloride from dimethyl sulfone for spectroscopic identification. Forensic Science International, 2018, 282, 86-91. | 2.2 | 0 |
| 119 | Chiral Capillary Electrophoresis of Amphetamine-Type Stimulants. Denki Eido, 2015, 59, 64-66. | 0.0 | Ο |
| 120 | Expediting cannabis seed examination by combining color reaction and DNA testing. Japanese Journal of Forensic Science and Technology, 2021, , . | 0.1 | 0 |
| 121 | Evaluation of a cannabis seed examination method without cultivation process. Japanese Journal of Forensic Science and Technology, 2022, , . | 0.1 | 0 |