

Kenji tsujikawa

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7841822/kenji-tsuji-kawa-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

116
papers

1,570
citations

21
h-index

30
g-index

121
ext. papers

1,815
ext. citations

2.6
avg, IF

4.32
L-index

#	Paper	IF	Citations
116	Degradation of 1-phenyl-2-propanone during long-term storage: useful information for methamphetamine impurity profiling. <i>Forensic Toxicology</i> , 2021 , 39, 405-416	2.6	0
115	Development of an improved method to estimate the days of continuous drug ingestion, based on the micro-segmental hair analysis. <i>Drug Testing and Analysis</i> , 2021 , 13, 1295-1304	3.5	3
114	DNA testing of suspected cannabis samples with exceptional morphology using a simple detection kit. <i>Forensic Toxicology</i> , 2021 , 39, 266-274	2.6	0
113	Rapid identification of drug-type and fiber-type cannabis by allele specific duplex PCR. <i>Forensic Science International</i> , 2021 , 318, 110634	2.6	1
112	Stereoselective analysis of ephedrine and its stereoisomers as impurities and/or by-products in seized methamphetamine by supercritical fluid chromatography/tandem mass spectrometry. <i>Forensic Science International</i> , 2021 , 318, 110591	2.6	4
111	Detection and confirmation of the ring-opened carboxylic acid metabolite of a new synthetic opioid furanylfentanyl. <i>Forensic Toxicology</i> , 2021 , 39, 114-122	2.6	2
110	Development of the selective concentration analytical method for drug-containing hair regions based on micro-segmental analysis to identify a trace amount of drug in hair: hair analysis following single-dose ingestion of midazolam. <i>Forensic Toxicology</i> , 2021 , 39, 156-166	2.6	4
109	Evaluation of Agonistic Activity of Fluorinated and Nonfluorinated Fentanyl Analogs on μ Opioid Receptor Using a Cell-Based Assay System. <i>Biological and Pharmaceutical Bulletin</i> , 2021 , 44, 159-161	2.3	2
108	Development and demonstration of cannabis DNA detection kit using DNA chromatography chip. <i>Japanese Journal of Forensic Science and Technology</i> , 2021 , 26, 29-48	0.1	
107	Differentiation of ring-substituted regioisomers of cathinone analogs by supercritical fluid chromatography. <i>Analytical Science Advances</i> , 2020 , 1, 22	1.1	1
106	Measurement of three-dimensional distributions of drugs in nails using liquid chromatography/tandem mass spectrometry after micro-segmentation to elucidate drug uptake routes. <i>Analytica Chimica Acta</i> , 2020 , 1108, 89-97	6.6	3
105	Studies on the phase I metabolites of the new designer drug 1-(2,3-dihydro-1H-inden-5-yl)-2-(pyrrolidine-1-yl)butan-1-one (5-PPDI) in human urine. <i>Forensic Science International</i> , 2020 , 310, 110214	2.6	5
104	Phosgene in deteriorated chloroform: presumptive cause of production of 3,4-dimethyl-5-phenyl-2-oxazolidones in methamphetamine. <i>Forensic Toxicology</i> , 2020 , 38, 475-480	2.6	3
103	Metabolism of a new synthetic opioid tetrahydrofuranylfentanyl in fresh isolated human hepatocytes: Detection and confirmation of ring-opened metabolites. <i>Drug Testing and Analysis</i> , 2020 , 12, 439-448	3.5	5
102	Metabolism of Butyrylfentanyl in Fresh Human Hepatocytes: Chemical Synthesis of Authentic Metabolite Standards for Definitive Identification. <i>Biological and Pharmaceutical Bulletin</i> , 2019 , 42, 623-630	2.3	13
101	Strong evidence of drug-facilitated crimes by hair analysis using LCMS/MS after micro-segmentation. <i>Forensic Toxicology</i> , 2019 , 37, 480-487	2.6	13
100	Rapid detection of hypnotics using surface-enhanced Raman scattering based on gold nanoparticle co-aggregation in a wet system. <i>Analyst, The</i> , 2019 , 144, 2158-2165	5	14

99	Rapid detection of synthetic cannabinoids in herbal highs using surface-enhanced Raman scattering produced by gold nanoparticle co-aggregation in a wet system. <i>Analyst, The</i> , 2019 , 144, 6928-6935	5	7
98	Estimation of day of death using micro-segmental hair analysis based on drug use history: a case of lidocaine use as a marker. <i>International Journal of Legal Medicine</i> , 2019 , 133, 117-122	3.1	6
97	Simultaneous chiral impurity analysis of methamphetamine and its precursors by supercritical fluid chromatography tandem mass spectrometry. <i>Forensic Toxicology</i> , 2019 , 37, 145-153	2.6	11
96	Development of rapid and simple method for DNA extraction from cannabis resin based on the evaluation of relative PCR amplification ability. <i>Forensic Science International</i> , 2018 , 287, 176-182	2.6	4
95	Comparison and evaluation of the quick purification methods of methamphetamine hydrochloride from dimethyl sulfone for spectroscopic identification. <i>Forensic Science International</i> , 2018 , 282, 86-91	2.6	
94	Micro-segmental hair analysis for proving drug-facilitated crimes: Evidence that a victim ingested a sleeping aid, diphenhydramine, on a specific day. <i>Forensic Science International</i> , 2018 , 288, 23-28	2.6	18
93	Metabolism of Fentanyl and Acetylfentanyl in Human-Induced Pluripotent Stem Cell-Derived Hepatocytes. <i>Biological and Pharmaceutical Bulletin</i> , 2018 , 41, 106-114	2.3	18
92	Different localizations of drugs simultaneously administered in a strand of hair by micro-segmental analysis. <i>Drug Testing and Analysis</i> , 2018 , 10, 750-760	3.5	14
91	Use of hepatocytes isolated from a liver-humanized mouse for studies on the metabolism of drugs: application to the metabolism of fentanyl and acetylfentanyl. <i>Forensic Toxicology</i> , 2018 , 36, 467-475	2.6	16
90	Development of simple and accurate detection systems for Cannabis sativa using DNA chromatography. <i>Forensic Science International</i> , 2018 , 291, 68-75	2.6	8
89	Accurate Estimation of Drug Intake Day by Microsegmental Analysis of a Strand of Hair by Use of Internal Temporal Markers. <i>Journal of Applied Laboratory Medicine</i> , 2018 , 3, 37-47	2	10
88	Time-course measurements of drug concentrations in hair and toenails after single administrations of pharmaceutical products. <i>Drug Testing and Analysis</i> , 2017 , 9, 571-577	3.5	17
87	Enantioseparation of methamphetamine by supercritical fluid chromatography with cellulose-based packed column. <i>Forensic Science International</i> , 2017 , 273, 39-44	2.6	20
86	Characterization and Differentiation of Geometric Isomers of 3-methylfentanyl Analogs by Gas Chromatography/Mass Spectrometry, Liquid Chromatography/Mass Spectrometry, and Nuclear Magnetic Resonance Spectroscopy. <i>Journal of Forensic Sciences</i> , 2017 , 62, 1472-1478	1.8	13
85	Differentiation of ring-substituted regioisomers of amphetamine and methamphetamine by supercritical fluid chromatography. <i>Drug Testing and Analysis</i> , 2017 , 9, 389-398	3.5	12
84	Synthesis and Analysis of Glucuronic Acid-Conjugated Metabolites of 4-Bromo-2,5-Dimethoxyphenethylamine. <i>Journal of Forensic Sciences</i> , 2017 , 62, 488-492	1.8	3
83	Development of a new field-test procedure for cocaine. <i>Forensic Science International</i> , 2017 , 270, 267-274	4.6	10
82	Evaluation of drug identification and discrimination ability of portable spectrometers. <i>Japanese Journal of Forensic Science and Technology</i> , 2017 , 22, 9-24	0.1	0

81	Approaching over 10000-fold sensitivity increase in chiral capillary electrophoresis: Cation-selective exhaustive injection and sweeping cyclodextrin-modified micellar electrokinetic chromatography. <i>Electrophoresis</i> , 2016 , 37, 2970-2976	3.6	14
80	Urinary Excretion Profiles of 2,5-Dimethoxy-4-alkylthiophenethylamine Analogs in Rats. <i>Biological and Pharmaceutical Bulletin</i> , 2016 , 39, 883-6	2.3	0
79	Development of a Library Search-Based Screening System for 3,4-Methylenedioxyamphetamine in Ecstasy Tablets Using a Portable Near-Infrared Spectrometer. <i>Journal of Forensic Sciences</i> , 2016 , 61, 1208-14	1.8	7
78	Highly sensitive quantification of unconjugated 11-nor-9-carboxy-(Δ)-tetrahydrocannabinol in a cannabis user's hair using micropulverized extraction. <i>Forensic Science International</i> , 2016 , 262, e34-6	2.6	6
77	Differentiation of ring-substituted bromoamphetamine analogs by gas chromatography-tandem mass spectrometry. <i>Forensic Toxicology</i> , 2016 , 34, 125-132	2.6	9
76	Effectiveness of saliva and fingerprints as alternative specimens to urine and blood in forensic drug testing. <i>Drug Testing and Analysis</i> , 2016 , 8, 644-51	3.5	15
75	Three-step drug extraction from a single sub-millimeter segment of hair and nail to determine the exact day of drug intake. <i>Analytica Chimica Acta</i> , 2016 , 948, 40-47	6.6	19
74	Development of a novel immunoassay for herbal cannabis using a new fluorescent antibody probe, "Ultra Quenchbody". <i>Forensic Science International</i> , 2016 , 266, 541-548	2.6	11
73	Comments on "Characterization of four new designer drugs, 5-chloro-NNEI, NNEI indazole analog, Δ PHPP and Δ POP, with 11 newly distributed designer drugs in illegal products". <i>Forensic Science International</i> , 2015 , 251, e15-7	2.6	4
72	Differentiation of regioisomeric chloroamphetamine analogs using gas chromatography-chemical ionization-tandem mass spectrometry. <i>Forensic Toxicology</i> , 2015 , 33, 338-347	2.6	13
71	Instability of the hydrochloride salts of cathinone derivatives in air. <i>Forensic Science International</i> , 2015 , 248, 48-54	2.6	5
70	Micro-pulverized extraction pretreatment for highly sensitive analysis of 11-nor-9-carboxy-(Δ)-tetrahydrocannabinol in hair by liquid chromatography/tandem mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2015 , 29, 2158-66	2.2	16
69	The use of a sulfonated capillary on chiral capillary electrophoresis/mass spectrometry of amphetamine-type stimulants for methamphetamine impurity profiling. <i>Forensic Science International</i> , 2015 , 249, 59-65	2.6	35
68	Detection of main metabolites of XLR-11 and its thermal degradation product in human hepatoma HepaRG cells and human urine. <i>Drug Testing and Analysis</i> , 2015 , 7, 341-5	3.5	19
67	Chiral Capillary Electrophoresis of Amphetamine-Type Stimulants 2015 , 59, 64-66		
66	Thermal degradation of a new synthetic cannabinoid QUPIC during analysis by gas chromatography-mass spectrometry. <i>Forensic Toxicology</i> , 2014 , 32, 201-207	2.6	31
65	Application of a portable near infrared spectrometer for presumptive identification of psychoactive drugs. <i>Forensic Science International</i> , 2014 , 242, 162-171	2.6	23
64	Time-course measurements of drugs and metabolites transferred from fingertips after drug administration: usefulness of fingerprints for drug testing. <i>Forensic Toxicology</i> , 2014 , 32, 235-242	2.6	15

63	Utilization of matrix-assisted laser desorption/ionization imaging mass spectrometry to search for cannabis in herb mixtures. <i>Analytical and Bioanalytical Chemistry</i> , 2014 , 406, 4789-94	4.4	5
62	Profiling of Methamphetamine. <i>Bunseki Kagaku</i> , 2014 , 63, 221-231	0.2	1
61	Differentiation of regioisomeric methylamphetamines by GC/MS. <i>Japanese Journal of Forensic Science and Technology</i> , 2014 , 19, 111-119	0.1	5
60	Simultaneous determination of tryptamine analogues in designer drugs using gas chromatography-mass spectrometry and liquid chromatography-tandem mass spectrometry. <i>Forensic Toxicology</i> , 2014 , 32, 154-161	2.6	7
59	Identification of the metabolites of 2,5-dimethoxy-4-ethylthiophenethylamine (2C-T-2) and 2,5-dimethoxy-4-isopropylthiophenethylamine (2C-T-4) in rat urine. <i>Japanese Journal of Forensic Science and Technology</i> , 2014 , 19, 91-101	0.1	1
58	Time-course measurements of caffeine and its metabolites extracted from fingertips after coffee intake: a preliminary study for the detection of drugs from fingerprints. <i>Analytical and Bioanalytical Chemistry</i> , 2013 , 405, 3945-52	4.4	27
57	Identification and differentiation of methcathinone analogs by gas chromatography-mass spectrometry. <i>Drug Testing and Analysis</i> , 2013 , 5, 670-7	3.5	12
56	Applicability of chemically modified capillaries in chiral capillary electrophoresis for methamphetamine profiling. <i>Forensic Science International</i> , 2013 , 226, 235-9	2.6	17
55	Chemical profiling of seized methamphetamine putatively synthesized from phenylacetic acid derivatives. <i>Forensic Science International</i> , 2013 , 227, 42-4	2.6	25
54	Analysis of 4-bromo-2,5-dimethoxyphenethylamine abuser's urine: identification and quantitation of urinary metabolites. <i>Journal of Forensic Sciences</i> , 2013 , 58, 279-87	1.8	17
53	Differentiation of regioisomeric fluoroamphetamine analogs by gas chromatography-mass spectrometry and liquid chromatography-tandem mass spectrometry. <i>Forensic Toxicology</i> , 2013 , 31, 241-250	2.6	28
52	Thermal degradation of pyrrolidinopentiophenone during injection in gas chromatography/mass spectrometry. <i>Forensic Science International</i> , 2013 , 231, 296-9	2.6	16
51	Rapid Chemical Examinations of Cannabis and Its Related Herbal Products. <i>Japanese Journal of Forensic Science and Technology</i> , 2013 , 18, 143-153	0.1	1
50	Interaction of 3,4-methylenedioxymethamphetamine and methamphetamine during metabolism by in vitro human metabolic enzymes and in rats. <i>Journal of Forensic Sciences</i> , 2012 , 57, 1008-13	1.8	11
49	Profiling of seized methamphetamine putatively synthesized by reductive amination of 1-phenyl-2-propanone. <i>Forensic Toxicology</i> , 2012 , 30, 70-75	2.6	21
48	Rapid, simple, and highly sensitive analysis of drugs in biological samples using thin-layer chromatography coupled with matrix-assisted laser desorption/ionization mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2012 , 402, 1257-67	4.4	21
47	Degradation pathways of 4-methylmethcathinone in alkaline solution and stability of methcathinone analogs in various pH solutions. <i>Forensic Science International</i> , 2012 , 220, 103-10	2.6	46
46	Distribution measurements of 3,4-methylenedioxymethamphetamine and its metabolites in organs by matrix-assisted laser desorption/ionization imaging mass spectrometry using an automatic matrix spraying system with an air brush and a turntable. <i>Analytical and Bioanalytical Chemistry</i> , 2012 , 404, 1883-89	4.4	18

45	Synthesis and identification of urinary metabolites of 4-iodo-2,5-dimethoxyphenethylamine. <i>Journal of Forensic Sciences</i> , 2011 , 56, 1319-23	1.8	8
44	A model system for prediction of the in vivo metabolism of designer drugs using three-dimensional culture of rat and human hepatocytes. <i>Forensic Toxicology</i> , 2011 , 29, 142-151	2.6	3
43	Distribution measurement of amphetamine-type stimulants in organs using micropulverized extraction and liquid chromatography/tandem mass spectrometry to complement drug distribution using mass spectrometry imaging. <i>Rapid Communications in Mass Spectrometry</i> , 2011 , 25, 2397-406	2.2	7
42	Urinary excretion profiles of N-hydroxy-3,4-methylenedioxymethamphetamine in rats. <i>Xenobiotica</i> , 2011 , 41, 578-84	2	
41	Recreational drugs, 3,4-Methylenedioxymethamphetamine(MDMA), 3,4-methylenedioxyamphetamine (MDA) and diphenylprolinol, inhibit neurite outgrowth in PC12 cells. <i>Journal of Toxicological Sciences</i> , 2010 , 35, 375-81	1.9	13
40	Increase in split ratio enables detection of underivatized N-hydroxy-3,4-methylenedioxymethamphetamine and N-hydroxy-3,4-methylenedioxyamphetamine by capillary GC-MS. <i>Forensic Toxicology</i> , 2010 , 28, 55-57	2.6	2
39	Seized methamphetamine samples with unique profiles of stable nitrogen isotopic composition documented by stable isotope ratio mass spectrometry. <i>Forensic Toxicology</i> , 2010 , 28, 119-123	2.6	12
38	Determination of 4-hydroxy-3-methoxymethamphetamine as a metabolite of methamphetamine in rats and human liver microsomes using gas chromatography-mass spectrometry and liquid chromatography-tandem mass spectrometry. <i>Journal of Analytical Toxicology</i> , 2009 , 33, 266-71	2.9	4
37	Thermal desorption counter-flow introduction atmospheric pressure chemical ionization for direct mass spectrometry of ecstasy tablets. <i>Journal of Mass Spectrometry</i> , 2009 , 44, 1300-7	2.2	12
36	Degradation of N-hydroxy-3,4-methylenedioxymethamphetamine in aqueous solution and its prevention. <i>Forensic Science International</i> , 2009 , 193, 106-11	2.6	5
35	Rapid identification and quantification of methamphetamine and amphetamine in hair by gas chromatography/mass spectrometry coupled with micropulverized extraction, aqueous acetylation and microextraction by packed sorbent. <i>Journal of Chromatography A</i> , 2009 , 1216, 4063-70	4.5	80
34	In vitro stability and metabolism of salvinorin A in rat plasma. <i>Xenobiotica</i> , 2009 , 39, 391-8	2	44
33	Uptake of 3,4-methylenedioxymethamphetamine and its related compounds by a proton-coupled transport system in Caco-2 cells. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2008 , 1778, 42-50	3.8	22
32	In vivo metabolism of alpha-methyltryptamine in rats: identification of urinary metabolites. <i>Xenobiotica</i> , 2008 , 38, 1476-86	2	7
31	Simple and simultaneous detection of methamphetamine and dimethyl sulfone in crystalline methamphetamine seizures by fast gas chromatography. <i>Forensic Toxicology</i> , 2008 , 26, 19-22	2.6	24
30	Evaluation method for linking methamphetamine seizures using stable carbon and nitrogen isotopic compositions: a complementary study with impurity profiling. <i>Rapid Communications in Mass Spectrometry</i> , 2008 , 22, 3816-22	2.2	15
29	Analysis of amphetamine-type stimulants and their metabolites in plasma, urine and bile by liquid chromatography with a strong cation-exchange column-tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008 , 867, 78-83	3.2	14
28	Development of an on-site screening system for amphetamine-type stimulant tablets with a portable attenuated total reflection Fourier transform infrared spectrometer. <i>Analytica Chimica Acta</i> , 2008 , 608, 95-103	6.6	18

27	Determination of salvinorin A and salvinorin B in <i>Salvia divinorum</i> -related products circulated in Japan. <i>Forensic Science International</i> , 2008 , 180, 105-9	2.6	20
26	Comparison and classification of methamphetamine seized in Japan and Thailand using gas chromatography with liquid-liquid extraction and solid-phase microextraction. <i>Forensic Science International</i> , 2008 , 175, 85-92	2.6	29
25	A fatal case of suspected anaphylaxis with cefoperazone and sulbactam: LC-MS analysis. <i>Journal of Forensic Sciences</i> , 2008 , 53, 226-31	1.8	12
24	Contribution of thermal desorption and liquid-liquid extraction for identification and profiling of impurities in methamphetamine by gas chromatography-mass spectrometry. <i>Forensic Science International</i> , 2007 , 171, 9-15	2.6	10
23	Interactions between 3,4-methylenedioxymethamphetamine, methamphetamine, ketamine, and caffeine in human intestinal Caco-2 cells and in oral administration to rats. <i>Forensic Science International</i> , 2007 , 170, 183-8	2.6	12
22	Determination of muscimol and ibotenic acid in <i>Amanita</i> mushrooms by high-performance liquid chromatography and liquid chromatography-tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2007 , 852, 430-5	3.2	31
21	In vivo metabolism of 2,5-dimethoxy-4-propylthiophenethylamine in rat. <i>Xenobiotica</i> , 2007 , 37, 679-92	2	11
20	Analysis of hallucinogenic constituents in <i>Amanita</i> mushrooms circulated in Japan. <i>Forensic Science International</i> , 2006 , 164, 172-8	2.6	34
19	In Vivo Metabolism of 5-Methoxy-N,N-diisopropyltryptamine in Rat. <i>Journal of Health Science</i> , 2006 , 52, 425-430		17
18	Forensic application of chiral separation of amphetamine-type stimulants to impurity analysis of seized methamphetamine by capillary electrophoresis. <i>Forensic Science International</i> , 2006 , 161, 92-6	2.6	35
17	A method for screening for various sedative-hypnotics in serum by liquid chromatography/single quadrupole mass spectrometry. <i>Forensic Science International</i> , 2006 , 157, 57-70	2.6	42
16	Identification of impurities and the statistical classification of methamphetamine using headspace solid phase microextraction and gas chromatography-mass spectrometry. <i>Forensic Science International</i> , 2006 , 160, 44-52	2.6	22
15	Effects of the Various Preparation Procedures of Dragendorff Reagent on Sensitivity for Thin Layer Chromatography. <i>Japanese Journal of Forensic Science and Technology</i> , 2005 , 10, 127-133	0.1	1
14	Urinary excretion profiles of two major triazolam metabolites, alpha-hydroxytriazolam and 4-hydroxytriazolam. <i>Journal of Analytical Toxicology</i> , 2005 , 29, 240-3	2.9	11
13	Analysis of Benzylpiperazine-like Compounds. <i>Japanese Journal of Science and Technology for Identification</i> , 2004 , 9, 165-184		8
12	Optimized Conditions for the Enzymatic Hydrolysis of .ALPHA.-Hydroxytriazolam-Glucuronide in Human Urine. <i>Journal of Health Science</i> , 2004 , 50, 286-289		2
11	. <i>Japanese Journal of Science and Technology for Identification</i> , 2004 , 9, 71-78		4
10	Excretory Profile of 4-Bromo-2,5-dimethoxyphenethylamine (2C-B) in Rat.. <i>Journal of Health Science</i> , 2003 , 49, 166-169		21

9	Morphological and chemical analysis of magic mushrooms in Japan. <i>Forensic Science International</i> , 2003 , 138, 85-90	2.6	46
8	Chiral analysis of amphetamine-type stimulants using reversed-polarity capillary electrophoresis/positive ion electrospray ionization tandem mass spectrometry. <i>Electrophoresis</i> , 2003 , 24, 1770-6	3.6	51
7	Potentiation of domperidone-induced catalepsy by a P-glycoprotein inhibitor, cyclosporin A. <i>Biopharmaceutics and Drug Disposition</i> , 2003 , 24, 105-14	1.7	23
6	Methamphetamine impurity profiling using a 0.32 mm i.d. nonpolar capillary column. <i>Forensic Science International</i> , 2003 , 135, 42-7	2.6	39
5	Protease-sensitive urinary pheromones induce region-specific Fos-expression in rat accessory olfactory bulb. <i>Biochemical and Biophysical Research Communications</i> , 1999 , 260, 222-4	3.4	15
4	Agonistic activity of fentanyl analogs and their metabolites on opioid receptors. <i>Forensic Toxicology</i> ,1	2.6	1
3	Distribution profiles of diphenhydramine and lidocaine in scalp, axillary, and pubic hairs measured by micro-segmental hair analysis: good indicator for discrimination between administration and external contamination of the drugs. <i>Forensic Toxicology</i> ,1	2.6	1
2	Thin-layer chromatography on silver nitrate-impregnated silica gel for analysis of homemade tetrahydrocannabinol mixtures. <i>Forensic Toxicology</i> ,1	2.6	0
1	Micro-segmental hair analysis: detailed procedures and applications in forensic toxicology. <i>Forensic Toxicology</i> ,1	2.6	1