

# William Deraedt

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7857872/publications.pdf>

Version: 2024-02-01

10  
papers

2,189  
citations

1040056

9  
h-index

1372567

10  
g-index

10  
all docs

10  
docs citations

10  
times ranked

2032  
citing authors

#	ARTICLE	IF	CITATIONS
1	Ciltacabtagene Autoleucel, an Anti-BCMA B-cell Maturation Antigen Chimeric Antigen Receptor T-Cell Therapy, for Relapsed/Refractory Multiple Myeloma: CARTITUDE-1 2-Year Follow-Up. <i>Journal of Clinical Oncology</i> , 2023, 41, 1265-1274.	1.6	160
2	Ciltacabtagene autoleucel, a B-cell maturation antigen (BCMA)-directed chimeric antigen receptor T-cell (CAR-T) therapy, in relapsed/refractory multiple myeloma (R/R MM): Updated results from CARTITUDE-1. <i>Journal of Clinical Oncology</i> , 2021, 39, 8005-8005.	1.6	23
3	Matching-adjusted indirect comparison of efficacy outcomes for ciltacabtagene autoleucel in CARTITUDE-1 versus idcabtagene vicleucel in KarMMa for the treatment of patients with relapsed or refractory multiple myeloma. <i>Current Medical Research and Opinion</i> , 2021, 37, 1779-1788.	1.9	21
4	Ciltacabtagene autoleucel, a B-cell maturation antigen-directed chimeric antigen receptor T-cell therapy in patients with relapsed or refractory multiple myeloma (CARTITUDE-1): a phase 1b/2 open-label study. <i>Lancet</i> , 2021, 398, 314-324.	13.7	711
5	The effects of different schedules of bortezomib, melphalan, and prednisone for patients with newly diagnosed multiple myeloma who are transplant ineligible: a matching-adjusted indirect comparison. <i>Leukemia and Lymphoma</i> , 2020, 61, 680-690.	1.3	9
6	Daratumumab plus Bortezomib, Melphalan, and Prednisone for Untreated Myeloma. <i>New England Journal of Medicine</i> , 2018, 378, 518-528.	27.0	747
7	Persistent Overall Survival Benefit and No Increased Risk of Second Malignancies With Bortezomib-Melphalan-Prednisone Versus Melphalan-Prednisone in Patients With Previously Untreated Multiple Myeloma. <i>Journal of Clinical Oncology</i> , 2013, 31, 448-455.	1.6	250
8	Characterization of haematological parameters with bortezomib-melphalan-prednisone versus melphalan-prednisone in newly diagnosed myeloma, with evaluation of long-term outcomes and risk of thromboembolic events with use of erythropoiesis-stimulating agents: analysis of the VISTA trial. <i>British Journal of Haematology</i> , 2011, 153, 212-221.	2.5	13
9	Superior outcomes associated with complete response in newly diagnosed multiple myeloma patients treated with nonintensive therapy: analysis of the phase 3 VISTA study of bortezomib plus melphalan-prednisone versus melphalan-prednisone. <i>Blood</i> , 2010, 116, 3743-3750.	1.4	101
10	VMP (Bortezomib, Melphalan, and Prednisone) Is Active and Well Tolerated in Newly Diagnosed Patients With Multiple Myeloma With Moderately Impaired Renal Function, and Results in Reversal of Renal Impairment: Cohort Analysis of the Phase III VISTA Study. <i>Journal of Clinical Oncology</i> , 2009, 27, 6086-6093.	1.6	154