

Chung-Jen Lee

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7859836/chung-jen-lee-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

68

papers

766

citations

16

h-index

23

g-index

68

ext. papers

883

ext. citations

3.1

avg, IF

3.92

L-index

#	Paper	IF	Citations
68	Dapagliflozin Ameliorates Lipopolysaccharide Related Acute Kidney Injury in Mice with Streptozotocin-induced Diabetes Mellitus.. <i>International Journal of Medical Sciences</i> , 2022 , 19, 729-739	3.7	1
67	Linagliptin Protects against Endotoxin-Induced Acute Kidney Injury in Rats by Decreasing Inflammatory Cytokines and Reactive Oxygen Species. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
66	High-Serum Angiotensin-Like Protein 3 Levels Associated with Cardiovascular Outcome in Patients with Coronary Artery Disease. <i>International Journal of Hypertension</i> , 2020 , 2020, 2980954	2.4	3
65	Specific drug delivery efficiently induced human breast tumor regression using a lipoplex by non-covalent association with anti-tumor antibodies. <i>Journal of Nanobiotechnology</i> , 2019 , 17, 25	9.4	14
64	Metabolic Syndrome With Aortic Arterial Stiffness And First Hospitalization Or Mortality In Coronary Artery Disease Patients. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2019 , 12, 2065-2073	3.4	2
63	Osteocalcin and carotid-femoral pulse wave velocity in patients on peritoneal dialysis. <i>Tzu Chi Medical Journal</i> , 2019 , 31, 23-28	1.1	3
62	Reply from authors for Serum leptin levels and peripheral arterial stiffness in patients with type 2 diabetes. <i>Tzu Chi Medical Journal</i> , 2019 , 31, 295	1.1	
61	Serum cystatin C is independently associated with aortic arterial stiffness in patients with type 2 diabetes. <i>Clinica Chimica Acta</i> , 2018 , 480, 114-118	6.2	4
60	Serum adipocyte fatty acid-binding protein level is associated with arterial stiffness quantified with cardio-ankle vascular index in kidney transplant patients. <i>Clinical and Experimental Nephrology</i> , 2018 , 22, 188-195	2.5	4
59	Hyperleptinemia positively associated with central arterial stiffness in hemodialysis patients. <i>PLoS ONE</i> , 2018 , 13, e0190694	3.7	8
58	TRPM8 and RAAS-mediated hypertension is critical for cold-induced immunosuppression in mice. <i>Oncotarget</i> , 2018 , 9, 12781-12795	3.3	9
57	Positive correlation of serum leptin levels and peripheral arterial stiffness in patients with type 2 diabetes. <i>Tzu Chi Medical Journal</i> , 2018 , 30, 10-14	1.1	4
56	High serum leptin level is associated with peripheral artery disease in adult peritoneal dialysis patients. <i>Tzu Chi Medical Journal</i> , 2018 , 30, 85-89	1.1	4
55	Hyperleptinemia is associated with the aortic augmentation index in kidney transplant recipients. <i>Tzu Chi Medical Journal</i> , 2018 , 30, 152-157	1.1	2
54	Association between hyperleptinemia and cardiovascular outcomes in patients with coronary artery disease. <i>Therapeutics and Clinical Risk Management</i> , 2018 , 14, 1855-1862	2.9	12
53	Serum levels of sclerostin as a potential biomarker in central arterial stiffness among hypertensive patients. <i>BMC Cardiovascular Disorders</i> , 2018 , 18, 214	2.3	9
52	High Levels of Serum Adipocyte Fatty Acid-binding Protein Predict Cardiovascular Events in Coronary Artery Disease Patients. <i>International Journal of Medical Sciences</i> , 2018 , 15, 1268-1274	3.7	5

51	Positive correlation of the serum angiopoietin-like protein 3 levels with the aortic augmentation index in patients with coronary artery disease. <i>Therapeutics and Clinical Risk Management</i> , 2018 , 14, 231-236	2.9	5
50	Acute Alcohol Intoxication Exacerbates Rhabdomyolysis-Induced Acute Renal Failure in Rats. <i>International Journal of Medical Sciences</i> , 2017 , 14, 680-689	3.7	14
49	Serum resistin as an independent marker of aortic stiffness in patients with coronary artery disease. <i>PLoS ONE</i> , 2017 , 12, e0183123	3.7	12
48	A comparison of Cornell and Sokolow-Lyon electrocardiographic criteria for left ventricular hypertrophy in a military male population in Taiwan: the Cardiorespiratory fitness and Hospitalization Events in armed Forces study. <i>Cardiovascular Diagnosis and Therapy</i> , 2017 , 7, 244-251	2.6	16
47	High serum adipocyte fatty acid binding protein level as a potential biomarker of aortic arterial stiffness in hypertensive patients with metabolic syndrome. <i>Clinica Chimica Acta</i> , 2017 , 473, 166-172	6.2	16
46	Heat adaptation from regular hot water immersion decreases proinflammatory responses, HSP70 expression, and physical heat stress. <i>Journal of Thermal Biology</i> , 2017 , 69, 95-103	2.9	5
45	Low serum adiponectin level is associated with metabolic syndrome and is an independent marker of peripheral arterial stiffness in hypertensive patients. <i>Diabetology and Metabolic Syndrome</i> , 2017 , 9, 49	5.6	21
44	High serum resistin levels are associated with peripheral artery disease in the hypertensive patients. <i>BMC Cardiovascular Disorders</i> , 2017 , 17, 80	2.3	10
43	Serum leptin is a predictor for central arterial stiffness in hypertensive patients. <i>Nephrology</i> , 2017 , 22, 783-789	2.2	13
42	Association of hyperleptinemia with peripheral arterial disease in hypertensive patients. <i>Tzu Chi Medical Journal</i> , 2017 , 29, 148-153	1.1	3
41	Serum leptin level positively correlates with metabolic syndrome among elderly Taiwanese. <i>Tzu Chi Medical Journal</i> , 2017 , 29, 159-164	1.1	9
40	Calcitriol decreases pro-inflammatory cytokines and protects against severe hemorrhagic shock induced-organ damage in rats. <i>Cytokine</i> , 2016 , 83, 262-268	4	4
39	Prophylactic acetylsalicylic acid attenuates the inflammatory response but fails to protect exercise-induced liver damage in exercised rats. <i>European Journal of Pharmacology</i> , 2016 , 786, 204-211	5.3	5
38	Serum Sclerostin as an Independent Marker of Peripheral Arterial Stiffness in Renal Transplantation Recipients: A Cross-Sectional Study. <i>Medicine (United States)</i> , 2016 , 95, e3300	1.8	26
37	Serum osteoprotegerin levels associated with the aortic augmentation index in renal transplant recipients. <i>Tzu Chi Medical Journal</i> , 2016 , 28, 20-23	1.1	2
36	Hypoadiponectinemia correlates with arterial stiffness in kidney transplantation patients. <i>Clinical and Experimental Nephrology</i> , 2015 , 19, 534-41	2.5	7
35	Positive correlation of serum adipocyte fatty acid binding protein levels with carotid-femoral pulse wave velocity in geriatric population. <i>BMC Geriatrics</i> , 2015 , 15, 88	4.1	19
34	Serum osteoprotegerin is associated with arterial stiffness assessed according to the cardio-ankle vascular index in hypertensive patients. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 304-12	4	22

33	N-terminal pro-B-type natriuretic peptide level inversely associates with metabolic syndrome in elderly persons. <i>Diabetology and Metabolic Syndrome</i> , 2014 , 6, 15	5.6	5
32	Serum atrial natriuretic peptide level inversely associates with metabolic syndrome in older adults. <i>Geriatrics and Gerontology International</i> , 2014 , 14, 640-6	2.9	6
31	Calcitriol decreases TGF- β and angiotensin II production and protects against chlorhexide digluconate-induced liver peritoneal fibrosis in rats. <i>Cytokine</i> , 2014 , 65, 105-18	4	18
30	Serum osteopontin level correlates with carotid-femoral pulse wave velocity in geriatric persons. <i>BioMed Research International</i> , 2014 , 2014, 570698	3	14
29	Association of serum osteoprotegerin levels with carotid-femoral pulse wave velocity in hypertensive patients. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 301-8	2.3	27
28	Inverse association of long-acting natriuretic peptide with metabolic syndrome in congestive heart failure patients. <i>Diabetology and Metabolic Syndrome</i> , 2013 , 5, 19	5.6	8
27	Low serum long-acting natriuretic peptide level correlates with metabolic syndrome in hypertensive patients: a cross-sectional study. <i>Archives of Medical Research</i> , 2013 , 44, 215-20	6.6	11
26	Heavy ethanol intoxication increases proinflammatory cytokines and aggravates hemorrhagic shock-induced organ damage in rats. <i>Mediators of Inflammation</i> , 2013 , 2013, 121786	4.3	11
25	N-terminal pro-B-type natriuretic peptide is associated with arterial stiffness measured using the cardio-ankle vascular index in renal transplant recipients. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 646-53	4	14
24	Recombinant human erythropoietin reduces rhabdomyolysis-induced acute renal failure in rats. <i>Injury</i> , 2012 , 43, 367-73	2.5	27
23	Erythropoietin ameliorates severe hemorrhagic shock-induced serum proinflammatory cytokines and biochemical changes in spontaneously hypertensive rats. <i>Tzu Chi Medical Journal</i> , 2012 , 24, 46-50	1.1	2
22	Inverse association of serum long-acting natriuretic peptide and bone mineral density in renal transplant recipients. <i>Clinical Transplantation</i> , 2012 , 26, E105-10	3.8	10
21	A model of chlorhexidine digluconate-induced peritoneal fibrosis in rats. <i>Tzu Chi Medical Journal</i> , 2012 , 24, 108-115	1.1	3
20	Freshwater clam extract decreased hemorrhagic shock-induced liver injury by attenuating TNF- α production. <i>Biological Research for Nursing</i> , 2012 , 14, 286-93	2.6	12
19	Valsartan decreases TGF- β production and protects against chlorhexidine digluconate-induced liver peritoneal fibrosis in rats. <i>Cytokine</i> , 2011 , 53, 223-30	4	26
18	Hyperleptinaemia positively correlated with metabolic syndrome in hemodialysis patients. <i>European Journal of Internal Medicine</i> , 2011 , 22, e105-9	3.9	12
17	Melatonin ameliorates hemorrhagic shock-induced organ damage in rats. <i>Journal of Surgical Research</i> , 2011 , 167, e315-21	2.5	27
16	Hypoadiponectinemia is associated with metabolic syndrome in hemodialysis patients. <i>Tzu Chi Medical Journal</i> , 2011 , 23, 77-81	1.1	1

15	Rosiglitazone ameliorates endotoxin-induced organ damage in conscious rats. <i>Biological Research for Nursing</i> , 2011 , 13, 38-43	2.6	16
14	Beneficial effects of enalapril on chlorhexidine digluconate-induced liver peritoneal fibrosis in rats. <i>Chinese Journal of Physiology</i> , 2011 , 54, 225-34	1.6	5
13	Fasting serum leptin level correlates with mid-arm fat area in peritoneal dialysis patients. <i>Therapeutic Apheresis and Dialysis</i> , 2010 , 14, 583-8	1.9	9
12	Aliskiren ameliorates chlorhexidine digluconate-induced peritoneal fibrosis in rats. <i>European Journal of Clinical Investigation</i> , 2010 , 40, 301-9	4.6	18
11	Hyperleptinemia positively correlated with metabolic syndrome in renal transplant recipients. <i>Clinical Transplantation</i> , 2010 , 24, E124-9	3.8	11
10	Fasting serum adiponectin level inversely correlates with metabolic syndrome in peritoneal dialysis patients. <i>Blood Purification</i> , 2010 , 30, 1-7	3.1	12
9	Fasting adiponectin is inversely correlated with metabolic syndrome in patients with coronary artery disease. <i>Internal Medicine</i> , 2010 , 49, 739-47	1.1	9
8	Fasting serum level of fatty-acid-binding protein 4 positively correlates with metabolic syndrome in patients with coronary artery disease. <i>Circulation Journal</i> , 2010 , 74, 327-31	2.9	47
7	Fasting serum fatty acid-binding protein 4 level positively correlates with metabolic syndrome in hemodialysis patients. <i>Archives of Medical Research</i> , 2010 , 41, 536-40	6.6	12
6	Propofol protects against hemorrhagic shock-induced organ damage in conscious spontaneously hypertensive rats. <i>Biological Research for Nursing</i> , 2009 , 11, 152-62	2.6	9
5	Fluvastatin attenuates severe hemorrhagic shock-induced organ damage in rats. <i>Resuscitation</i> , 2009 , 80, 372-8	4	33
4	Intraocular pressure assessment in both eyes of the same patient after laser in situ keratomileusis. <i>Journal of Cataract and Refractive Surgery</i> , 2009 , 35, 76-82	2.3	11
3	Pentobarbital reduces rhabdomyolysis-induced acute renal failure in conscious rats. <i>Journal of Trauma</i> , 2009 , 67, 132-8		10
2	Low-dose propofol ameliorates haemorrhagic shock-induced organ damage in conscious rats. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2008 , 35, 766-74	3	15
1	Freshwater clam extract ameliorates acute liver injury induced by hemorrhage in rats. <i>The American Journal of Chinese Medicine</i> , 2008 , 36, 1121-33	6	19