

Anne Tailleux

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7865598/publications.pdf>

Version: 2024-02-01

93
papers

6,433
citations

87401

40
h-index

78623

77
g-index

94
all docs

94
docs citations

94
times ranked

10353
citing authors

#	ARTICLE	IF	CITATIONS
1	Apolipoprotein F is reduced in humans with steatosis and controls plasma triglyceride-rich lipoprotein metabolism. <i>Hepatology</i> , 2023, 77, 1287-1302.	3.6	3
2	Bile acids contribute to the development of non-alcoholic steatohepatitis in mice. <i>JHEP Reports</i> , 2022, 4, 100387.	2.6	28
3	Enterohepatic, Gluco-metabolic, and Gut Microbial Characterization of Individuals With Bile Acid Malabsorption. , 2022, 1, 299-312.		5
4	Enterohepatic Takeda G-Protein Coupled Receptor 5 Agonism in Metabolic Dysfunction-Associated Fatty Liver Disease and Related Glucose Dysmetabolism. <i>Nutrients</i> , 2022, 14, 2707.	1.7	8
5	Deletion of fibroblast activation protein provides atheroprotection. <i>Cardiovascular Research</i> , 2021, 117, 1060-1069.	1.8	20
6	NASH-related increases in plasma bile acid levels depend on insulin resistance. <i>JHEP Reports</i> , 2021, 3, 100222.	2.6	24
7	Beyond the Rule of 5: Impact of PEGylation with Various Polymer Sizes on Pharmacokinetic Properties, Structure-Properties Relationships of mPEGylated Small Agonists of TGR5 Receptor. <i>Journal of Medicinal Chemistry</i> , 2021, 64, 1593-1610.	2.9	9
8	Multiple Selection Criteria for Probiotic Strains with High Potential for Obesity Management. <i>Nutrients</i> , 2021, 13, 713.	1.7	19
9	Characterization of one anastomosis gastric bypass and impact of biliary and common limbs on bile acid and postprandial glucose metabolism in a minipig model. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2021, 320, E772-E783.	1.8	8
10	A randomized placebo-controlled trial of elafibranor in patients with primary biliary cholangitis and incomplete response to UDCA. <i>Journal of Hepatology</i> , 2021, 74, 1344-1354.	1.8	77
11	Hypothalamic bile acid-TGR5 signaling protects from obesity. <i>Cell Metabolism</i> , 2021, 33, 1483-1492.e10.	7.2	79
12	The ALGOVUE Clinical Trial: Effects of the Daily Consumption of Eggs Enriched with Lutein and Docosahexaenoic Acid on Plasma Composition and Macular Pigment Optical Density. <i>Nutrients</i> , 2021, 13, 3347.	1.7	9
13	Inflammation-induced cholestasis in cancer cachexia. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2021, 12, 70-90.	2.9	24
14	Farnesoid X Receptor Activation in Brain Alters Brown Adipose Tissue Function via the Sympathetic System. <i>Frontiers in Molecular Neuroscience</i> , 2021, 14, 808603.	1.4	9
15	Bile acids associate with glucose metabolism, but do not predict conversion from impaired fasting glucose to diabetes. <i>Metabolism: Clinical and Experimental</i> , 2020, 103, 154042.	1.5	21
16	Deletion of the nuclear receptor ROR α in macrophages does not modify the development of obesity, insulin resistance and NASH. <i>Scientific Reports</i> , 2020, 10, 21095.	1.6	6
17	Analysis of the association of MPO and MMP-9 with stroke severity and outcome. <i>Neurology</i> , 2020, 95, e97-e108.	1.5	42
18	The nuclear receptor FXR inhibits Glucagon-Like Peptide-1 secretion in response to microbiota-derived Short-Chain Fatty Acids. <i>Scientific Reports</i> , 2020, 10, 174.	1.6	45

#	ARTICLE	IF	CITATIONS
19	Sirt6 deletion in bone marrow-derived cells increases atherosclerosis – Central role of macrophage scavenger receptor 1. <i>Journal of Molecular and Cellular Cardiology</i> , 2020, 139, 24-32.	0.9	26
20	Plasma BCAA Changes in Patients With NAFLD Are Sex Dependent. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, 2311-2321.	1.8	39
21	Transcription profiling in the liver of undernourished male rat offspring reveals altered lipid metabolism pathways and predisposition to hepatic steatosis. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2019, 317, E1094-E1107.	1.8	6
22	Brain insulin response and peripheral metabolic changes in a Tau transgenic mouse model. <i>Neurobiology of Disease</i> , 2019, 125, 14-22.	2.1	16
23	Hepatic PPAR α is critical in the metabolic adaptation to sepsis. <i>Journal of Hepatology</i> , 2019, 70, 963-973.	1.8	53
24	Bile acid alterations in nonalcoholic fatty liver disease, obesity, insulin resistance and type 2 diabetes: what do the human studies tell?. <i>Current Opinion in Lipidology</i> , 2019, 30, 244-254.	1.2	39
25	Targeting the gut microbiota with inulin-type fructans: preclinical demonstration of a novel approach in the management of endothelial dysfunction. <i>Gut</i> , 2018, 67, 271-283.	6.1	150
26	Roux-en-Y gastric bypass increases systemic but not portal bile acid concentrations by decreasing hepatic bile acid uptake in minipigs. <i>International Journal of Obesity</i> , 2017, 41, 664-668.	1.6	21
27	Bile Acid Control of Metabolism and Inflammation in Obesity, Type 2 Diabetes, Dyslipidemia, and Nonalcoholic Fatty Liver Disease. <i>Gastroenterology</i> , 2017, 152, 1679-1694.e3.	0.6	630
28	Anacetrapib, but not evacetrapib, impairs endothelial function in CETP-transgenic mice in spite of marked HDL-C increase. <i>Atherosclerosis</i> , 2017, 257, 186-194.	0.4	17
29	Topical Intestinal Aminoimidazole Agonists of G-Protein-Coupled Bile Acid Receptor 1 Promote Glucagon Like Peptide-1 Secretion and Improve Glucose Tolerance. <i>Journal of Medicinal Chemistry</i> , 2017, 60, 4185-4211.	2.9	48
30	Bile Acid Alterations Are Associated With Insulin Resistance, but Not With NASH, in Obese Subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 3783-3794.	1.8	78
31	Tau deletion promotes brain insulin resistance. <i>Journal of Experimental Medicine</i> , 2017, 214, 2257-2269.	4.2	158
32	Metabolic effects of bile acid sequestration. <i>Current Opinion in Endocrinology, Diabetes and Obesity</i> , 2016, 23, 138-144.	1.2	9
33	Lipid-lowering drugs prevent neurovascular and cognitive consequences of cardiopulmonary bypass. <i>Vascular Pharmacology</i> , 2016, 80, 59-66.	1.0	11
34	Influence of Roux-en-Y gastric bypass on plasma bile acid profiles: a comparative study between rats, pigs and humans. <i>International Journal of Obesity</i> , 2016, 40, 1260-1267.	1.6	61
35	Liver X Receptor Regulates Triglyceride Absorption Through Intestinal Down-regulation of Scavenger Receptor Class B, Type 1. <i>Gastroenterology</i> , 2016, 150, 650-658.	0.6	41
36	Bariatric surgery, lipoprotein metabolism and cardiovascular risk. <i>Current Opinion in Lipidology</i> , 2015, 26, 317-324.	1.2	15

#	ARTICLE	IF	CITATIONS
37	The Sirt1 activator SRT3025 provides atheroprotection in ApoE ^{-/-} mice by reducing hepatic Pcsk9 secretion and enhancing Ldlr expression. <i>European Heart Journal</i> , 2015, 36, 51-59.	1.0	117
38	Cholesterol uptake disruption, in association with chemotherapy, is a promising combined metabolic therapy for pancreatic adenocarcinoma. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 2473-2478.	3.3	310
39	Rapid and Body Weight-Independent Improvement of Endothelial and High-Density Lipoprotein Function After Roux-en-Y Gastric Bypass. <i>Circulation</i> , 2015, 131, 871-881.	1.6	103
40	The Bile Acid Chenodeoxycholic Acid Increases Human Brown Adipose Tissue Activity. <i>Cell Metabolism</i> , 2015, 22, 418-426.	7.2	342
41	Farnesoid X receptor inhibits glucagon-like peptide-1 production by enteroendocrine L cells. <i>Nature Communications</i> , 2015, 6, 7629.	5.8	274
42	Screening strategy to generate cell specific recombination: a case report with the RIP-Cre mice. <i>Transgenic Research</i> , 2015, 24, 803-812.	1.3	8
43	Glucose sensing O-GlcNAcylation pathway regulates the nuclear bile acid receptor farnesoid X receptor (FXR). <i>Hepatology</i> , 2014, 59, 2022-2033.	3.6	55
44	Impact of Endotoxin Challenge in Obese Pigs. <i>Shock</i> , 2014, 41, 546-553.	1.0	10
45	Hepatic trans-Golgi action coordinated by the GTPase ARFRP1 is crucial for lipoprotein lipidation and assembly. <i>Journal of Lipid Research</i> , 2014, 55, 41-52.	2.0	15
46	<i>Cdkn2a</i> / <i>p16</i> / <i>Ink4a</i> Regulates Fasting-Induced Hepatic Gluconeogenesis Through the PKA-CREB-PGC1 β Pathway. <i>Diabetes</i> , 2014, 63, 3199-3209.	0.3	36
47	The transrepressive activity of peroxisome proliferator-activated receptor alpha is necessary and sufficient to prevent liver fibrosis in mice. <i>Hepatology</i> , 2014, 60, 1593-1606.	3.6	116
48	Sox17 Regulates Liver Lipid Metabolism and Adaptation to Fasting. <i>PLoS ONE</i> , 2014, 9, e104925.	1.1	15
49	Hepatoprotective effects of the dual peroxisome proliferator-activated receptor alpha/delta agonist, GFT505, in rodent models of nonalcoholic fatty liver disease/nonalcoholic steatohepatitis. <i>Hepatology</i> , 2013, 58, 1941-1952.	3.6	355
50	Long-term prognostic value of preprocedural adiponectin levels in patients undergoing percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2013, 168, 4921-4924.	0.8	3
51	Normal HDL apo AI turnover and cholesterol enrichment of HDL subclasses in New Zealand rabbits with partial nephrectomy. <i>Metabolism: Clinical and Experimental</i> , 2013, 62, 492-498.	1.5	8
52	Detrimental Effects of Diet-Induced Obesity on β , Pathology Are Independent of Insulin Resistance in β , Transgenic Mice. <i>Diabetes</i> , 2013, 62, 1681-1688.	0.3	88
53	Activation of intestinal peroxisome proliferator-activated receptor- δ increases high-density lipoprotein production. <i>European Heart Journal</i> , 2013, 34, 2566-2574.	1.0	44
54	Endogenous cannabinoid receptor CB1 activation promotes vascular smooth-muscle cell proliferation and neointima formation. <i>Journal of Lipid Research</i> , 2013, 54, 1360-1368.	2.0	23

#	ARTICLE	IF	CITATIONS
55	171. Critical Care Medicine, 2013, 41, A37.	0.4	1
56	Inhibition of hepatic scavenger receptor-class B type I by RNA interference decreases atherosclerosis in rabbits. <i>Atherosclerosis</i> , 2012, 222, 360-366.	0.4	11
57	Roles of PPARs in NAFLD: Potential therapeutic targets. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2012, 1821, 809-818.	1.2	229
58	Cell-derived microparticles in atherosclerosis: biomarkers and targets for pharmacological modulation?. <i>Journal of Cellular and Molecular Medicine</i> , 2012, 16, 1365-1376.	1.6	65
59	Bone Marrow p16INK4a-Deficiency Does Not Modulate Obesity, Glucose Homeostasis or Atherosclerosis Development. <i>PLoS ONE</i> , 2012, 7, e32440.	1.1	14
60	PPAR α activation differently affects microparticle content in atherosclerotic lesions and liver of a mouse model of atherosclerosis and NASH. <i>Atherosclerosis</i> , 2011, 218, 69-76.	0.4	24
61	p16INK4a deficiency promotes IL-4-induced polarization and inhibits proinflammatory signaling in macrophages. <i>Blood</i> , 2011, 118, 2556-2566.	0.6	89
62	Overview of the Measurement of Lipids and Lipoproteins in Mice. <i>Current Protocols in Mouse Biology</i> , 2011, 1, 265-277.	1.2	2
63	Peroxisome Proliferator-Activated Receptor- α Gene Level Differently Affects Lipid Metabolism and Inflammation in Apolipoprotein E2 Knock-In Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 1573-1579.	1.1	66
64	Human Atherosclerotic Plaque Alternative Macrophages Display Low Cholesterol Handling but High Phagocytosis Because of Distinct Activities of the PPAR α and LXR α Pathways. <i>Circulation Research</i> , 2011, 108, 985-995.	2.0	318
65	Rexinoid Bexarotene Modulates Triglyceride but not Cholesterol Metabolism via Gene-Specific Permissivity of the RXR/LXR Heterodimer in the Liver. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009, 29, 1488-1495.	1.1	63
66	Morphologic and Electroretinographic Phenotype of SR-BI Knockout Mice after a Long-Term Atherogenic Diet. , 2009, 50, 3931.		35
67	Novel non-carboxylic acid retinoids: 1,2,4-Oxadiazol-5-one derivatives. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 489-492.	1.0	15
68	Niemann-Pick C1 like 1 gene expression is down-regulated by LXR activators in the intestine. <i>Biochemical and Biophysical Research Communications</i> , 2006, 340, 1259-1263.	1.0	156
69	The RXR Agonist Bexarotene Improves Cholesterol Homeostasis and Inhibits Atherosclerosis Progression in a Mouse Model of Mixed Dyslipidemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006, 26, 2731-2737.	1.1	69
70	Increased Susceptibility of Low-Density Lipoprotein to Ex Vivo Oxidation in Mice Transgenic for Human Apolipoprotein B Treated with 1 Melatonin-Related Compound Is Not Associated with Atherosclerosis Progression. <i>Journal of Cardiovascular Pharmacology</i> , 2005, 46, 241-249.	0.8	18
71	HOST GLUCOSE METABOLISM MEDIATES T4 AND IL-7 ACTION ON SCHISTOSOMA MANSONI DEVELOPMENT. <i>Journal of Parasitology</i> , 2005, 91, 737-744.	0.3	18
72	PPAR α , but not PPAR β , Activators Decrease Macrophage-Laden Atherosclerotic Lesions in a Nondiabetic Mouse Model of Mixed Dyslipidemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005, 25, 1897-1902.	1.1	70

#	ARTICLE	IF	CITATIONS
73	Paradoxical exacerbation of combined hyperlipidemia in human apolipoprotein A-II transgenic mice treated with fenofibrate. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1737, 130-137.	1.2	20
74	Murine models to investigate pharmacological compounds acting as ligands of PPARs in dyslipidemia and atherosclerosis. <i>Trends in Pharmacological Sciences</i> , 2003, 24, 530-534.	4.0	26
75	Lipid Free Apolipoprotein E Binds to the Class B Type I Scavenger Receptor I (SR-BI) and Enhances Cholesteryl Ester Uptake from Lipoproteins. <i>Journal of Biological Chemistry</i> , 2002, 277, 36092-36099.	1.6	50
76	Apolipoprotein A-II, HDL metabolism and atherosclerosis. <i>Atherosclerosis</i> , 2002, 164, 1-13.	0.4	126
77	Daily melatonin supplementation in mice increases atherosclerosis in proximal aorta. <i>Biochemical and Biophysical Research Communications</i> , 2002, 293, 1114-1123.	1.0	31
78	Melatonin related compounds inhibit lipid peroxidation during copper or free radical-induced LDL oxidation. <i>Journal of Pineal Research</i> , 2002, 33, 109-117.	3.4	25
79	Increased levels of high-density lipoprotein cholesterol are ineffective in inhibiting the development of immune responses to oxidized low-density lipoprotein and atherosclerosis in transgenic rabbits expressing human apolipoprotein (apo) A-I with severe hypercholesterolaemia. <i>Clinical Science</i> , 2001, 100, 343-355.	1.8	9
80	Increased levels of high-density lipoprotein cholesterol are ineffective in inhibiting the development of immune responses to oxidized low-density lipoprotein and atherosclerosis in transgenic rabbits expressing human apolipoprotein (apo) A-I with severe hypercholesterolaemia. <i>Clinical Science</i> , 2001, 100, 343.	1.8	8
81	Improved Lipid and Lipoprotein Profile, Hepatic Insulin Sensitivity, and Glucose Tolerance in 11 β -Hydroxysteroid Dehydrogenase Type 1 Null Mice. <i>Journal of Biological Chemistry</i> , 2001, 276, 41293-41300.	1.6	395
82	Peroxisome Proliferator-activated Receptor α Is Not Rate-limiting for the Lipoprotein-lowering Action of Fish Oil. <i>Journal of Biological Chemistry</i> , 2001, 276, 4634-4639.	1.6	70
83	Decreased Susceptibility to Diet-Induced Atherosclerosis in Human Apolipoprotein A-II Transgenic Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2000, 20, 2453-2458.	1.1	51
84	Paraoxonase Activity Is Reduced by a Pro-atherosclerotic Diet in Rabbits. <i>Biochemical and Biophysical Research Communications</i> , 2000, 269, 232-236.	1.0	85
85	Absence of relationship between plasma Lp(a), Lp-AI, anti-oxidized LDL autoantibodies, LDL immune complexes concentrations and restenosis after percutaneous transluminal coronary angioplasty. <i>Clinica Chimica Acta</i> , 2000, 299, 129-140.	0.5	12
86	3-Hydroxy-3-methylglutaryl CoA reductase inhibitors reduce serum triglyceride levels through modulation of apolipoprotein C-III and lipoprotein lipase. <i>FEBS Letters</i> , 1999, 452, 160-164.	1.3	80
87	Cholesterol Efflux, Lecithin â^{c} Cholesterol Acyltransferase Activity, and Pre- β Particle Formation by Serum from Human Apolipoprotein A-I and Apolipoprotein A-I/Apolipoprotein A-II Transgenic Mice Consistent with the Latter Being Less Effective for Reverse Cholesterol Transport â^{c} . <i>Biochemistry</i> , 1997, 36, 2243-2249.	1.2	66
88	Transgenic Rabbits Expressing Human Apolipoprotein A-I in the Liver. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1996, 16, 1424-1429.	1.1	55
89	Inhibition of Atherosclerosis Development in Cholesterol-Fed Human Apolipoprotein A-II Transgenic Rabbits. <i>Circulation</i> , 1996, 94, 713-717.	1.6	184
90	A 700-bp fragment of the human antithrombin III promoter is sufficient to confer high, tissue-specific expression on human apolipoprotein A-II in transgenic mice. <i>Gene</i> , 1995, 156, 199-205.	1.0	17

#	ARTICLE	IF	CITATIONS
91	Prevention of in vitro low-density lipoprotein oxidation by an albumin-containing Lp A-I subfraction. <i>Lipids and Lipid Metabolism</i> , 1995, 1255, 31-38.	2.6	20
92	Tissue-specific Expression of the Human Gene for Lecithin: Cholesterol Acyltransferase in Transgenic Mice Alters Blood Lipids, Lipoproteins and Lipases towards a Less Atherogenic Profile. <i>FEBS Journal</i> , 1995, 230, 567-575.	0.2	53
93	Expression of apolipoprotein B epitopes in low density lipoproteins of hemodialyzed patients. <i>Kidney International</i> , 1993, 44, 1360-1365.	2.6	11