
Ines Bado

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7876944/publications.pdf Version: 2024-02-01



INES RADO

#	Article	IF	CITATIONS
1	Transferable Resistance to Highest Priority Critically Important Antibiotics for Human Health in Escherichia coli Strains Obtained From Livestock Feces in Uruguay. Frontiers in Veterinary Science, 2020, 7, 588919.	2.2	18
2	Detection of qnrVC6, within a new genetic context, in an NDM-1-producing Citrobacter freundii clinical isolate from Uruguay. Journal of Global Antimicrobial Resistance, 2018, 14, 95-98.	2.2	11
3	Zoonotic Potential and Antibiotic Resistance of <i>Escherichia coli</i> in Neonatal Calves in Uruguay. Microbes and Environments, 2017, 32, 275-282.	1.6	32
4	CTX-M-15 in combination with aac(6′)-lb -cr is the most prevalent mechanism of resistance both in Escherichia coli and Klebsiella pneumoniae , including K. pneumoniae ST258, in an ICU in Uruguay. Journal of Global Antimicrobial Resistance, 2016, 6, 5-9.	2.2	32
5	Extended-spectrum β-lactamases, transferable quinolone resistance, and virulotyping in extra-intestinal E. coli in Uruguay. Journal of Infection in Developing Countries, 2016, 10, 43-52.	1.2	23
6	Sepsis caused by New Delhi metallo-β-lactamase (blaNDM-1) and qnrD-producing Morganella morganii, treated successfully with fosfomycin and meropenem: case report and literature review. International Journal of Infectious Diseases, 2015, 30, 20-26.	3.3	55
7	Polyclonal endemicity of Acinetobacter baumannii in ventilated patients in an intensive care unit in Uruguay. International Journal of Infectious Diseases, 2013, 17, e422-e427.	3.3	15
8	Risk factors for the acquisition of extended-spectrum beta-lactamase-producing Enterobacteriaceae in hospitalized children. Journal of Infection in Developing Countries, 2013, 7, 361-364.	1.2	5
9	First Human Isolate of Salmonella enterica Serotype Enteritidis HarboringblaCTX-M-14in South America. Antimicrobial Agents and Chemotherapy, 2012, 56, 2132-2134.	3.2	20
10	Detection of class 1 and 2 integrons, extended-spectrum β-lactamases and qnr alleles in enterobacterial isolates from the digestive tract of Intensive Care Unit inpatients. International Journal of Antimicrobial Agents, 2010, 36, 453-458.	2.5	39
11	Ciprofloxacin-Resistant Enterobacteria Harboring the <i>aac(6</i> ′ <i>)-lb-cr</i> Variant Isolated from Feces of Inpatients in an Intensive Care Unit in Uruguay. Antimicrobial Agents and Chemotherapy, 2008, 52, 806-807.	3.2	28