## Kaori Kitamura

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7889677/publications.pdf

Version: 2024-02-01

	933447		888059	
18	455	10	17	
papers	citations	h-index	g-index	
19	19	19	594	
all docs	docs citations	times ranked	citing authors	

#	Article	IF	CITATIONS
1	Validity of Short and Long Self-Administered Food Frequency Questionnaires in Ranking Dietary Intake in Middle-Aged and Elderly Japanese in the Japan Public Health Center-Based Prospective Study for the Next Generation (JPHC-NEXT) Protocol Area. Journal of Epidemiology, 2016, 26, 420-432.	2.4	180
2	Elevated C-Reactive Protein Is Associated with Cognitive Decline in Outpatients of a General Hospital: The Project in Sado for Total Health (PROST). Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 10-19.	1.3	48
3	Rural–urban differences in the prevalence of cognitive impairment in independent community-dwelling elderly residents of Ojiya city, Niigata Prefecture, Japan. Environmental Health and Preventive Medicine, 2016, 21, 422-429.	3.4	32
4	Modifiable Factors Associated with Cognitive Impairment in 1,143 Japanese Outpatients: The Project in Sado for Total Health (PROST). Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 341-349.	1.3	29
5	The Murakami Cohort Study of vitamin D for the prevention of musculoskeletal and other age-related diseases: a study protocol. Environmental Health and Preventive Medicine, 2018, 23, 28.	3.4	25
6	Molecular Network Analysis of the Urinary Proteome of Alzheimer's Disease Patients. Dementia and Geriatric Cognitive Disorders Extra, 2019, 9, 53-65.	1.3	24
7	Urinary Apolipoprotein C3 Is a Potential Biomarker for Alzheimer's Disease. Dementia and Geriatric Cognitive Disorders Extra, 2021, 10, 94-104.	1.3	19
8	Short daytime napping reduces the risk of cognitive decline in community-dwelling older adults: a 5-year longitudinal study. BMC Geriatrics, 2021, 21, 474.	2.7	18
9	Psychological distress as a risk factor for dementia after the 2004 Niigata–Chuetsu earthquake in Japan. Journal of Affective Disorders, 2019, 259, 121-127.	4.1	14
10	Low serum 25-hydroxyvitamin D increases cognitive impairment in elderly people. Journal of Bone and Mineral Metabolism, 2019, 37, 368-375.	2.7	13
11	Online version of the self-administered food frequency questionnaire for the Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT) protocol: Relative validity, usability, and comparison with a printed questionnaire. Journal of Epidemiology, 2017, 27, 435-446.	2.4	12
12	High serum 25-hydroxyvitamin D levels do not retard postmenopausal bone loss in Japanese women: the Yokogoshi study. Archives of Osteoporosis, 2013, 8, 153.	2.4	9
13	Association between dialysis treatment and cognitive decline: A study from the Project in Sado for Total Health (PROST), Japan. Geriatrics and Gerontology International, 2017, 17, 1584-1587.	1.5	9
14	Weight loss from 20 years of age is associated with cognitive impairment in middle-aged and elderly individuals. PLoS ONE, 2017, 12, e0185960.	2.5	9
15	Alterations in Glycerolipid and Fatty Acid Metabolic Pathways in Alzheimer's Disease Identified by Urinary Metabolic Profiling: A Pilot Study. Frontiers in Neurology, 2021, 12, 719159.	2.4	8
16	Body mass index and risk of recurrent falls in communityâ€dwelling Japanese aged 40–74 years: The Murakami cohort study. Geriatrics and Gerontology International, 2021, 21, 498-505.	1.5	3
17	Leisure-Time and Non–Leisure-Time Physical Activities are Dose-Dependently Associated With a Reduced Risk of Dementia in Community-Dwelling People Aged 40-74ÂYears: The Murakami Cohort Study. Journal of the American Medical Directors Association, 2022, 23, 1197-1204.e4.	2.5	3
18	Secular changes in bone mineral density of adult Japanese women from 1995 to 2013. Fukushima Journal of Medical Sciences, 2021, 67, 128-134.	0.4	0