Waclaw Tworzydlo

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/7889940/waclaw-tworzydlo-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

45	517	14	19
papers	citations	h-index	g-index
47 ext. papers	635 ext. citations	2.6 avg, IF	4.02 L-index

#	Paper	IF	Citations
45	A mixture of persistent organic pollutants detected in human follicular fluid increases progesterone secretion and mitochondrial activity in human granulosa HGrC1 cells. <i>Reproductive Toxicology</i> , 2021 , 104, 114-124	3.4	O
44	Towards understanding leydigioma: do G protein-coupled estrogen receptor and peroxisome proliferator-activated receptor regulate lipid metabolism and steroidogenesis in Leydig cell tumors?. <i>Protoplasma</i> , 2020 , 257, 1149-1163	3.4	7
43	Transmission of Functional, Wild-Type Mitochondria and the Fittest mtDNA to the Next Generation: Bottleneck Phenomenon, Balbiani Body, and Mitophagy. <i>Genes</i> , 2020 , 11,	4.2	5
42	Interstitial Leydig Cell Tumorigenesis-Leptin and Adiponectin Signaling in Relation to Aromatase Expression in the Human Testis. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	7
41	Apelin and apelin receptor in human placenta: Expression, signalling pathway and regulation of trophoblast JEG-3 and BeWo cells proliferation and cell cycle. <i>International Journal of Molecular Medicine</i> , 2020 , 45, 691-702	4.4	3
40	"Real life" polycyclic aromatic hydrocarbon (PAH) mixtures modulate hCG, hPL and hPLGF levels and disrupt the physiological ratio of MMP-2 to MMP-9 and VEGF expression in human placenta cell lines. <i>Reproductive Toxicology</i> , 2020 , 95, 1-10	3.4	3
39	Morphogenesis of the ovarian follicular epithelium during initial stages of embryogenesis of the viviparous earwig, Hemimerus talpoides. <i>Journal of Morphology</i> , 2020 , 281, 47-54	1.6	2
38	Morphogenesis of the Balbiani body in developing oocytes of an orthopteran, Metrioptera brachyptera, and multiplication of female germline mitochondria. <i>Journal of Morphology</i> , 2020 , 281, 114	42:915	51 ^O
37	Do estrogens regulate lipid status in testicular steroidogenic Leydig cell?. <i>Acta Histochemica</i> , 2019 , 121, 611-618	2	5
36	Organelle assemblages implicated in the transfer of oocyte components to the embryo: an insect perspective. <i>Current Opinion in Insect Science</i> , 2019 , 31, 1-7	5.1	8
35	Viviparity in the dermapteran Arixenia esau: respiration inside mother body requires both maternal and larval contribution. <i>Protoplasma</i> , 2019 , 256, 1573-1584	3.4	2
34	Do G-protein coupled estrogen receptor and bisphenol A analogs influence on Leydig cell epigenetic regulation in immature boar testis ex vivo?. <i>Animal Reproduction Science</i> , 2019 , 207, 21-35	2.1	14
33	Evolutionary origin and functioning of pregenital abdominal outgrowths in a viviparous insect, Arixenia esau. <i>Scientific Reports</i> , 2019 , 9, 16090	4.9	1
32	Viviparity in Two Closely Related Epizoic Dermapterans Relies on Disparate Modifications of Reproductive Systems and Embryogenesis. <i>Results and Problems in Cell Differentiation</i> , 2019 , 68, 455-47	, 5 1.4	1
31	Telocytes in the mouse testicular interstitium: implications of G-protein-coupled estrogen receptor (GPER) and estrogen-related receptor (ERR) in the regulation of mouse testicular interstitial cells. Protoplasma, 2019, 256, 393-408	3.4	16
30	Telocytes are localized to testis of the bank vole (Myodes glareolus) and are affected by lighting conditions and G-coupled membrane estrogen receptor (GPER) signaling. <i>General and Comparative Endocrinology</i> , 2019 , 271, 39-48	3	13
29	Morphogenesis of serial abdominal outgrowths during development of the viviparous dermapteran, Arixenia esau (Insecta, Dermaptera). <i>Arthropod Structure and Development</i> , 2019 , 49, 62-6	5 ∮ .8	4

(2014-2018)

28	Apelin and apelin receptor at different stages of corpus luteum development and effect of apelin on progesterone secretion and 3Ehydroxysteroid dehydrogenase (3EHSD) in pigs. <i>Animal Reproduction Science</i> , 2018 , 192, 251-260	2.1	14
27	The role of G-protein-coupled membrane estrogen receptor in mouse Leydig cell function-in vivo and in vitro evaluation. <i>Cell and Tissue Research</i> , 2018 , 374, 389-412	4.2	19
26	Excretion in the mother body: modifications of the larval excretory system in the viviparous dermapteran, Arixenia esau. <i>Protoplasma</i> , 2018 , 255, 1799-1809	3.4	4
25	Regulation of steroidogenic function of mouse Leydig cells: G-coupled membrane estrogen receptor and peroxisome proliferator-activated receptor partnership. <i>Journal of Physiology and Pharmacology</i> , 2018 , 69,	2.1	6
24	Unusual morphological adaptations and processes associated with viviparity in an epizoic dermapteran. <i>PLoS ONE</i> , 2018 , 13, e0195647	3.7	6
23	Insights into the role of estrogen-related receptors pland on tumor Leydig cells. <i>Tissue and Cell</i> , 2018 , 52, 78-91	2.7	16
22	Meiosis, Balbiani body and early asymmetry of Thermobia oocyte. <i>Protoplasma</i> , 2017 , 254, 649-655	3.4	9
21	Chlorinated biphenyls effect on estrogen-related receptor expression, steroid secretion, mitochondria ultrastructure but not on mitochondrial membrane potential in Leydig cells. <i>Cell and Tissue Research</i> , 2017 , 369, 429-444	4.2	16
20	Selection of mitochondria in female germline cells: is Balbiani body implicated in this process?. Journal of Assisted Reproduction and Genetics, 2017, 34, 1405-1412	3.4	30
19	Ovaries and oogenesis in an epizoic dermapteran, Hemimerus talpoides (Dermaptera, Hemimeridae): Structural and functional adaptations to viviparity and matrotrophy. <i>Zoology</i> , 2017 , 125, 32-40	1.7	7
18	The Pole (Germ) Plasm in Insect Oocytes. Results and Problems in Cell Differentiation, 2017, 63, 103-126	1.4	3
17	Are aryl hydrocarbon receptor and G-protein-coupled receptor 30 involved in the regulation of seasonal testis activity in photosensitive rodent-the bank vole (Myodes glareolus)?. <i>Theriogenology</i> , 2016 , 86, 674-686.e1	2.8	14
16	Flutamide induces alterations in the cell-cell junction ultrastructure and reduces the expression of Cx43 at the blood-testis barrier with no disturbance in the rat seminiferous tubule morphology. <i>Reproductive Biology and Endocrinology</i> , 2016 , 14, 14	5	12
15	Exclusion of dysfunctional mitochondria from Balbiani body during early oogenesis of Thermobia. <i>Cell and Tissue Research</i> , 2016 , 366, 191-201	4.2	24
14	Relationship between lateral oviduct morphology and reproductive strategy in earwigs. <i>Zoologischer Anzeiger</i> , 2015 , 254, 41-47	1.1	7
13	Octylphenol induces changes in glycosylation pattern, calcium level and ultrastructure of bank vole spermatozoa in vitro. <i>Toxicology in Vitro</i> , 2015 , 29, 529-37	3.6	5
12	Balbiani body, nuage and sponge bodiesterm plasm pathway players. <i>Arthropod Structure and Development</i> , 2014 , 43, 341-8	1.8	57
11	Ovaries and phylogeny of dermapterans once more: Ovarian characters support paraphyly of Spongiphoridae. <i>Zoologischer Anzeiger</i> , 2014 , 253, 321-326	1.1	7

10	Morphology and ultrastructure of the germarium in panoistic ovarioles of a basal "apterygotous" insect, Thermobia domestica. <i>Zoology</i> , 2014 , 117, 200-6	1.7	22
9	Morphology of the ovarioles and the mode of oogenesis of Arixenia esau support the inclusion of Arixeniina to the Eudermaptera. <i>Zoologischer Anzeiger</i> , 2013 , 252, 410-416	1.1	18
8	Embryos of the viviparous dermapteran, Arixenia esau develop sequentially in two compartments: terminal ovarian follicles and the uterus. <i>PLoS ONE</i> , 2013 , 8, e64087	3.7	16
7	A very simple mode of follicular cell diversification in Euborellia fulviceps (Dermaptera, Anisolabididae) involves actively migrating cells. <i>Zoological Science</i> , 2011 , 28, 802-8	0.8	7
6	Female germline stem cell niches of earwigs are structurally simple and different from those of Drosophila melanogaster. <i>Journal of Morphology</i> , 2010 , 271, 634-40	1.6	17
5	Structure of ovaries and oogenesis in dermapterans. II. The nurse cells, nuage aggregates and sponge bodies. <i>Folia Biologica</i> , 2010 , 58, 67-72	0.7	8
4	Ovaries and germline cysts and their evolution in Dermaptera (Insecta). <i>Arthropod Structure and Development</i> , 2010 , 39, 360-8	1.8	27
3	The Balbiani body in the female germline cells of an earwig, Opisthocosmia silvestris. <i>Zoological Science</i> , 2009 , 26, 754-7	0.8	13
2	Structure of ovaries and oogenesis in dermapterans. I. Origin and functioning of the ovarian follicles. <i>Arthropod Structure and Development</i> , 2008 , 37, 310-20	1.8	23
1	Differing strategies of patterning of follicular cells in higher and lower brachycerans (Diptera: Brachycera). <i>Genesis</i> , 2005 , 43, 49-58	1.9	18