

Jennifer M Specht

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7891428/publications.pdf>

Version: 2024-02-01

35
papers

1,862
citations

361413

20
h-index

414414

32
g-index

35
all docs

35
docs citations

35
times ranked

2582
citing authors

#	ARTICLE	IF	CITATIONS
1	Dendritic Cells Retrovirally Transduced with a Model Antigen Gene Are Therapeutically Effective against Established Pulmonary Metastases. <i>Journal of Experimental Medicine</i> , 1997, 186, 1213-1221.	8.5	297
2	Phase III Study of Iniparib Plus Gemcitabine and Carboplatin Versus Gemcitabine and Carboplatin in Patients With Metastatic Triple-Negative Breast Cancer. <i>Journal of Clinical Oncology</i> , 2014, 32, 3840-3847.	1.6	253
3	Immunogenic Chemotherapy Enhances Recruitment of CAR-T Cells to Lung Tumors and Improves Antitumor Efficacy when Combined with Checkpoint Blockade. <i>Cancer Cell</i> , 2021, 39, 193-208.e10.	16.8	157
4	Tumor Metabolism and Blood Flow Changes by Positron Emission Tomography: Relation to Survival in Patients Treated With Neoadjuvant Chemotherapy for Locally Advanced Breast Cancer. <i>Journal of Clinical Oncology</i> , 2008, 26, 4449-4457.	1.6	142
5	Fluoroestradiol Positron Emission Tomography Reveals Differences in Pharmacodynamics of Aromatase Inhibitors, Tamoxifen, and Fulvestrant in Patients with Metastatic Breast Cancer. <i>Clinical Cancer Research</i> , 2011, 17, 4799-4805.	7.0	120
6	Serial 2-[18F] fluoro-2-deoxy-d-glucose positron emission tomography (FDG-PET) to monitor treatment of bone-dominant metastatic breast cancer predicts time to progression (TTP). <i>Breast Cancer Research and Treatment</i> , 2007, 105, 87-94.	2.5	97
7	PET Tumor Metabolism in Locally Advanced Breast Cancer Patients Undergoing Neoadjuvant Chemotherapy: Value of Static versus Kinetic Measures of Fluorodeoxyglucose Uptake. <i>Clinical Cancer Research</i> , 2011, 17, 2400-2409.	7.0	94
8	A Phase 2 Study of ¹⁶ Î±-[18F]-fluoro-17Î²-estradiol Positron Emission Tomography (FES-PET) as a Marker of Hormone Sensitivity in Metastatic Breast Cancer (MBC). <i>Molecular Imaging and Biology</i> , 2014, 16, 431-440.	2.6	80
9	Tumor Metabolism and Blood Flow as Assessed by Positron Emission Tomography Varies by Tumor Subtype in Locally Advanced Breast Cancer. <i>Clinical Cancer Research</i> , 2010, 16, 2803-2810.	7.0	72
10	Kinetic Analysis of ¹⁸ F-Fluoride PET Images of Breast Cancer Bone Metastases. <i>Journal of Nuclear Medicine</i> , 2010, 51, 521-527.	5.0	65
11	Neoadjuvant Chemotherapy for Locally Advanced Breast Cancer. <i>Seminars in Radiation Oncology</i> , 2009, 19, 222-228.	2.2	51
12	Blood Flow-Metabolism Mismatch: Good for the Tumor, Bad for the Patient. <i>Clinical Cancer Research</i> , 2009, 15, 5294-5296.	7.0	43
13	Association between serial dynamic contrast-enhanced MRI and dynamic ¹⁸ F-FDG PET measures in patients undergoing neoadjuvant chemotherapy for locally advanced breast cancer. <i>Journal of Magnetic Resonance Imaging</i> , 2010, 32, 1124-1131.	3.4	41
14	Prospective Study of Serial ¹⁸ F-FDG PET and ¹⁸ F-Fluoride PET to Predict Time to Skeletal-Related Events, Time to Progression, and Survival in Patients with Bone-Dominant Metastatic Breast Cancer. <i>Journal of Nuclear Medicine</i> , 2018, 59, 1823-1830.	5.0	41
15	Dasatinib plus Capecitabine for Advanced Breast Cancer: Safety and Efficacy in Phase I Study CA180004. <i>Clinical Cancer Research</i> , 2013, 19, 1884-1893.	7.0	38
16	TBCRC026: Phase II Trial Correlating Standardized Uptake Value With Pathologic Complete Response to Pertuzumab and Trastuzumab in Breast Cancer. <i>Journal of Clinical Oncology</i> , 2019, 37, 714-722.	1.6	36
17	Advances in molecular imaging for breast cancer detection and characterization. <i>Breast Cancer Research</i> , 2012, 14, 206.	5.0	32
18	Feasibility study of FDG PET as an indicator of early response to aromatase inhibitors and trastuzumab in a heterogeneous group of breast cancer patients. <i>EJNMMI Research</i> , 2012, 2, 34.	2.5	27

#	ARTICLE	IF	CITATIONS
19	Combined Targeted Therapies for First-line Treatment of Metastatic Triple Negative Breast Cancer—A Phase II Trial of Weekly Nab-Paclitaxel and Bevacizumab Followed by Maintenance Targeted Therapy With Bevacizumab and Erlotinib. <i>Clinical Breast Cancer</i> , 2019, 19, e283-e296.	2.4	24
20	Updated Results of TBCRC026: Phase II Trial Correlating Standardized Uptake Value With Pathological Complete Response to Pertuzumab and Trastuzumab in Breast Cancer. <i>Journal of Clinical Oncology</i> , 2021, 39, 2247-2256.	1.6	22
21	Test-Retest Reproducibility of ¹⁸ F-FDG PET/CT Uptake in Cancer Patients Within a Qualified and Calibrated Local Network. <i>Journal of Nuclear Medicine</i> , 2019, 60, 608-614.	5.0	21
22	¹⁸ F-Fluoroestradiol PET Imaging in a Phase II Trial of Vorinostat to Restore Endocrine Sensitivity in ER+/HER2- Metastatic Breast Cancer. <i>Journal of Nuclear Medicine</i> , 2021, 62, 184-190.	5.0	20
23	Phase I/II Trial of Combined Pegylated Liposomal Doxorubicin and Cyclophosphamide in Metastatic Breast Cancer. <i>Clinical Breast Cancer</i> , 2018, 18, e143-e149.	2.4	15
24	Digital Mammography and Breast Tomosynthesis Performance in Women with a Personal History of Breast Cancer, 2007–2016. <i>Radiology</i> , 2021, 300, 290-300.	7.3	13
25	[¹⁸ F]Fluorodeoxyglucose Positron Emission Tomography–Computed Tomography in Breast Cancer: When and When Not?. <i>Journal of Clinical Oncology</i> , 2012, 30, 1252-1254.	1.6	11
26	Surveillance for second breast cancer events in women with a personal history of breast cancer using breast MRI: a systematic review and meta-analysis. <i>Breast Cancer Research and Treatment</i> , 2020, 181, 255-268.	2.5	11
27	¹⁸ F-fluorodeoxyglucose (FDG) PET or ¹⁸ F-fluorothymidine (FLT) PET to assess early response to aromatase inhibitors (AI) in women with ER+ operable breast cancer in a window-of-opportunity study. <i>Breast Cancer Research</i> , 2021, 23, 88.	5.0	11
28	Facility Variability in Examination Indication Among Women With Prior Breast Cancer: Implications and the Need for Standardization. <i>Journal of the American College of Radiology</i> , 2020, 17, 755-764.	1.8	9
29	Circulating biomarkers in patients receiving neoadjuvant chemotherapy combined with sunitinib for locally advanced breast cancer.. <i>Journal of Clinical Oncology</i> , 2013, 31, 1089-1089.	1.6	8
30	Adjuvant Metronomic CMF in a Contemporary Breast Cancer Cohort: What's Old Is New. <i>Clinical Breast Cancer</i> , 2015, 15, e277-e285.	2.4	6
31	A Phase II Study Evaluating the Safety and Efficacy of Sunitinib Malate in Combination With Weekly Paclitaxel Followed by Doxorubicin and Daily Oral Cyclophosphamide Plus G-CSF as Neoadjuvant Chemotherapy for Locally Advanced or Inflammatory Breast Cancer. <i>Clinical Breast Cancer</i> , 2022, 22, 32-42.	2.4	4
32	Optimal duration of trastuzumab for early HER2-positive breast cancer. <i>Lancet, The</i> , 2017, 389, 1167-1168.	13.7	1
33	Vorinostat to restore sensitivity to aromatase inhibitor therapy in metastatic breast cancer: A phase II clinical trial with ER imaging correlates.. <i>Journal of Clinical Oncology</i> , 2012, 30, TPS3109-TPS3109.	1.6	0
34	Quantitative measures of FDG PET after neoadjuvant chemotherapy to predict breast cancer patient survival.. <i>Journal of Clinical Oncology</i> , 2012, 30, 1088-1088.	1.6	0
35	A phase II study evaluating the safety and efficacy of sunitinib with weekly paclitaxel followed by doxorubicin and daily oral cyclophosphamide plus G-CSF as neoadjuvant chemotherapy (NC) for locally advanced (LABC) or inflammatory breast cancer (IBC).. <i>Journal of Clinical Oncology</i> , 2013, 31, 1090-1090.	1.6	0