
Walid Medhioub

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7894963/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Characterization of Coolia spp. (Gonyaucales, Dinophyceae) from Southern Tunisia: first record of Coolia malayensis in the Mediterranean Sea. Algae, 2021, 36, 175-193.	2.3	5
2	Ultraphytoplankton community composition in Southwestern and Eastern Mediterranean Basin: Relationships to water mass properties and nutrients. Journal of Sea Research, 2020, 158, 101875.	1.6	11
3	Strain selection of microalgae isolated from Tunisian coast: characterization of the lipid profile for potential biodiesel production. Bioprocess and Biosystems Engineering, 2018, 41, 1449-1459.	3.4	12
4	A Feedback Mechanism to Control Apoptosis Occurs in the Digestive Gland of the Oyster Crassostrea gigas Exposed to the Paralytic Shellfish Toxins Producer Alexandrium catenella. Marine Drugs, 2014, 12, 5035-5054.	4.6	12
5	Physiological responses of Manila clams Venerupis (=Ruditapes) philippinarum with varying parasite Perkinsus olseni burden to toxic algal Alexandrium ostenfeldii exposure. Aquatic Toxicology, 2014, 154, 27-38.	4.0	34
6	Exposure to the Neurotoxic Dinoflagellate, Alexandrium catenella, Induces Apoptosis of the Hemocytes of the Oyster, Crassostrea gigas. Marine Drugs, 2013, 11, 4799-4814.	4.6	30
7	Spirolide uptake and detoxification by Crassostrea gigas exposed to the toxic dinoflagellate Alexandrium ostenfeldii. Aquaculture, 2012, 358-359, 108-115.	3.5	17
8	Alexandrium ostenfeldii growth and spirolide production in batch culture and photobioreactor. Harmful Algae, 2011, 10, 794-803.	4.8	17
9	Detoxification enhancement in the gymnodimine-contaminated grooved carpet shell, Ruditapes decussatus (Linné). Harmful Algae, 2010, 9, 200-207.	4.8	24