
Daniel Gordin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7896568/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Glycemic control is not related to cerebral small vessel disease in neurologically asymptomatic individuals with type 1 diabetes. Acta Diabetologica, 2022, 59, 481-490.	1.2	2
2	Lipids in CKD: What do we actually know?. Atherosclerosis, 2022, , .	0.4	0
3	Arterial function, biomarkers, carcinoid syndrome and carcinoid heart disease in patients with small intestinal neuroendocrine tumours. Endocrine, 2022, 77, 177-187.	1.1	5
4	The European/International Fibromuscular Dysplasia Registry and Initiative (FEIRI)—clinical phenotypes and their predictors based on a cohort of 1000 patients. Cardiovascular Research, 2021, 117, 950-959.	1.8	33
5	Genome-wide association study on coronary artery disease in type 1 diabetes suggests beta-defensin 127 as a risk locus. Cardiovascular Research, 2021, 117, 600-612.	1.8	12
6	The Long-Term Incidence of Hospitalization for Ketoacidosis in Adults with Established T1D—A Prospective Cohort Study. Journal of Clinical Endocrinology and Metabolism, 2020, 105, 231-241.	1.8	14
7	Good longâ€ŧerm renal graft survival and low incidence of cardiac pathology in adults after short dialysis period and renal transplantation in early childhood – a cohort study. Transplant International, 2020, 33, 89-97.	0.8	4
8	Pregnancy-Related Complications in Patients With Fibromuscular Dysplasia. Hypertension, 2020, 76, 545-553.	1.3	10
9	Blood pressure response to renal denervation is correlated with baseline blood pressure variability. Journal of Hypertension, 2018, 36, 221-229.	0.3	20
10	Ambulatory blood pressure and arterial stiffness in individuals with type 1 diabetes. Diabetologia, 2018, 61, 1935-1945.	2.9	21
11	Urinary liver-type fatty acid binding protein is an independent predictor of stroke and mortality in individuals with type 1 diabetes. Diabetologia, 2017, 60, 1782-1790.	2.9	9
12	The effects of baroreflex activation therapy on blood pressure and sympathetic function in patients with refractory hypertension: the rationale and design of the Nordic BAT study*. Blood Pressure, 2017, 26, 294-302.	0.7	13
13	Oxygen-induced impairment in arterial function is corrected by slow breathing in patients with type 1 diabetes. Scientific Reports, 2017, 7, 6001.	1.6	14
14	Searching for Explanations for Cryptogenic Stroke in the Young: Revealing the Triggers, Causes, and Outcome (SECRETO): Rationale and design. European Stroke Journal, 2017, 2, 116-125.	2.7	30
15	Prognosis and Its Predictors After Incident Stroke in Patients With Type 1 Diabetes. Diabetes Care, 2017, 40, 1394-1400.	4.3	9
16	Acute-Phase Blood Pressure Levels Correlate With a High Risk of Recurrent Strokes in Young-Onset Ischemic Stroke. Stroke, 2016, 47, 1593-1598.	1.0	14
17	Influence of Postprandial Hyperglycemic Conditions on Arterial Stiffness in Patients With Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2016, 101, 1134-1143.	1.8	28
18	Aspects of Hyperglycemia Contribution to Arterial Stiffness and Cardiovascular Complications in Patients With Type 1 Diabetes. Journal of Diabetes Science and Technology, 2016, 10, 1059-1064.	1.3	22

DANIEL GORDIN

#	Article	IF	CITATIONS
19	Oxygen deteriorates arterial function in type 1 diabetes. Acta Diabetologica, 2016, 53, 349-357.	1.2	3
20	Urinary Adiponectin Is an Independent Predictor of Progression to End-Stage Renal Disease in Patients With Type 1 Diabetes and Diabetic Nephropathy. Diabetes Care, 2015, 38, 883-890.	4.3	32
21	Kidney Injury Molecule-1 and the Loss of Kidney Function in Diabetic Nephropathy: A Likely Causal Link in Patients With Type 1 Diabetes. Diabetes Care, 2015, 38, 1130-1137.	4.3	61
22	Health after Pregnancy in the Mother with Diabetes. Women's Health, 2015, 11, 471-476.	0.7	3
23	The Presence and Consequence of Nonalbuminuric Chronic Kidney Disease in Patients With Type 1 Diabetes. Diabetes Care, 2015, 38, 2128-2133.	4.3	56
24	Added Value of Soluble Tumor Necrosis Factor-α Receptor 1 as a Biomarker of ESRD Risk in Patients With Type 1 Diabetes. Diabetes Care, 2014, 37, 2334-2342.	4.3	45
25	Different Risk Factor Profiles for Ischemic and Hemorrhagic Stroke in Type 1 Diabetes Mellitus. Stroke, 2014, 45, 2558-2562.	1.0	39
26	Risk factors of hypertensive pregnancies in women with diabetes and the influence on their future life. Annals of Medicine, 2014, 46, 498-502.	1.5	11
27	Patients with type 1 diabetes show signs of vascular dysfunction in response to multiple high-fat meals. Nutrition and Metabolism, 2014, 11, 28.	1.3	17
28	Different Lipid Variables Predict Incident Coronary Artery Disease in Patients With Type 1 Diabetes With or Without Diabetic Nephropathy: The FinnDiane Study. Diabetes Care, 2014, 37, 2374-2382.	4.3	24
29	Osteopontin Is a Strong Predictor of Incipient Diabetic Nephropathy, Cardiovascular Disease, and All-Cause Mortality in Patients With Type 1 Diabetes. Diabetes Care, 2014, 37, 2593-2600.	4.3	63
30	Arterial function can be obtained by noninvasive finger pressure waveform. International Journal of Cardiology, 2014, 175, 169-171.	0.8	7
31	Pre-eclampsia and pregnancy-induced hypertension are associated with severe diabetic retinopathy in type 1 diabetes later in life. Acta Diabetologica, 2013, 50, 781-787.	1.2	34
32	Osteoprotegerin Is an Independent Predictor of Vascular Events in Finnish Adults With Type 1 Diabetes. Diabetes Care, 2013, 36, 1827-1833.	4.3	36
33	Arterial stiffness and vascular complications in patients with type 1 diabetes: The Finnish Diabetic Nephropathy (FinnDiane) Study. Annals of Medicine, 2012, 44, 196-204.	1.5	35
34	Pulse Pressure Predicts Incident Cardiovascular Disease but Not Diabetic Nephropathy in Patients With Type 1 Diabetes (The FinnDiane Study). Diabetes Care, 2011, 34, 886-891.	4.3	38
35	Acute hyperglycaemia induces an inflammatory response in young patients with type 1 diabetes. Annals of Medicine, 2008, 40, 627-633.	1.5	50