

Jos A Paramo

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7897874/jose-a-paramo-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40
papers

729
citations

15
h-index

26
g-index

44
ext. papers

879
ext. citations

4.4
avg, IF

3.72
L-index

#	Paper	IF	Citations
40	Different expression of MMPs/TIMP-1 in human atherosclerotic lesions. Relation to plaque features and vascular bed. <i>Atherosclerosis</i> , 2003 , 170, 269-76	3.1	85
39	Spanish Consensus Statement on alternatives to allogeneic blood transfusion: the 2013 update of the "Seville Document". <i>Blood Transfusion</i> , 2013 , 11, 585-610	3.6	75
38	Vitamins C and E downregulate vascular VEGF and VEGFR-2 expression in apolipoprotein-E-deficient mice. <i>Atherosclerosis</i> , 2003 , 171, 67-73	3.1	56
37	Arterial spin labeling MRI is able to detect early hemodynamic changes in diabetic nephropathy. <i>Journal of Magnetic Resonance Imaging</i> , 2017 , 46, 1810-1817	5.6	47
36	Matrix metalloproteinase 10 contributes to hepatocarcinogenesis in a novel crosstalk with the stromal derived factor 1/C-X-C chemokine receptor 4 axis. <i>Hepatology</i> , 2015 , 62, 166-78	11.2	44
35	Radioembolization of hepatocellular carcinoma activates liver regeneration, induces inflammation and endothelial stress and activates coagulation. <i>Liver International</i> , 2015 , 35, 1590-6	7.9	37
34	Matrix metalloproteinase-10 expression is induced during hepatic injury and plays a fundamental role in liver tissue repair. <i>Liver International</i> , 2014 , 34, e257-70	7.9	37
33	Matrix metalloproteinase 10 is associated with disease severity and mortality in patients with peripheral arterial disease. <i>Journal of Vascular Surgery</i> , 2015 , 61, 428-35	3.5	31
32	Serum levels of matrix metalloproteinase-10 are associated with the severity of atherosclerosis in patients with chronic kidney disease. <i>Kidney International</i> , 2010 , 78, 1275-80	9.9	31
31	Association between serum tissue inhibitor of matrix metalloproteinase-1 levels and mortality in patients with severe brain trauma injury. <i>PLoS ONE</i> , 2014 , 9, e94370	3.7	27
30	Randomized clinical trial on acute effects of i.v. iron sucrose during haemodialysis. <i>Nephrology</i> , 2010 , 15, 178-83	2.2	26
29	The CXCR4/SDF1 axis improves muscle regeneration through MMP-10 activity. <i>Stem Cells and Development</i> , 2014 , 23, 1417-27	4.4	24
28	Matrix metalloproteinase-10 deficiency delays atherosclerosis progression and plaque calcification. <i>Atherosclerosis</i> , 2018 , 278, 124-134	3.1	20
27	Functional and transcriptomic analysis of extracellular vesicles identifies calprotectin as a new prognostic marker in peripheral arterial disease (PAD). <i>Journal of Extracellular Vesicles</i> , 2020 , 9, 1729646	16.4	17
26	Design, synthesis, and biological evaluation of novel matrix metalloproteinase inhibitors as potent antihemorrhagic agents: from hit identification to an optimized lead. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 2465-88	8.3	17
25	CM352 Reduces Brain Damage and Improves Functional Recovery in a Rat Model of Intracerebral Hemorrhage. <i>Journal of the American Heart Association</i> , 2017 , 6,	6	14
24	Functional MMP-10 is required for efficient tissue repair after experimental hind limb ischemia. <i>FASEB Journal</i> , 2015 , 29, 960-72	0.9	14

23	Impact of surgery and chemotherapy on von Willebrand factor and vascular endothelial growth factor levels in colorectal cancer. <i>Clinical and Translational Oncology</i> , 2005 , 7, 150-5	3.6	12
22	Reduced high-density lipoprotein cholesterol: A valuable, independent prognostic marker in peripheral arterial disease. <i>Journal of Vascular Surgery</i> , 2017 , 66, 1527-1533.e1	3.5	11
21	Topical issues in venous thromboembolism. <i>Drugs</i> , 2010 , 70 Suppl 2, 11-8	12.1	11
20	Serum tissue inhibitor of matrix metalloproteinase-1 levels are associated with mortality in patients with malignant middle cerebral artery infarction. <i>BMC Neurology</i> , 2015 , 15, 111	3.1	10
19	Discovery and safety profiling of a potent preclinical candidate, (4-[4-[[[(3R)-3-(hydroxycarbamoyl)-8-azaspiro[4.5]decan-3-yl]sulfonyl]phenoxy]-N-methylbenzamide) (CM-352), for the prevention and treatment of hemorrhage. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 2941-57	8.3	10
18	Selective increase of cardiomyocyte derived extracellular vesicles after experimental myocardial infarction and functional effects on the endothelium. <i>Thrombosis Research</i> , 2018 , 170, 1-9	8.2	9
17	Purification and characterization of a variant of human prothrombin: prothrombin Segovia. <i>Thrombosis Research</i> , 1997 , 85, 465-77	8.2	7
16	The Role of Circulating Biomarkers in Peripheral Arterial Disease. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	7
15	Pulmonary Embolism, Pulmonary Microvascular Thrombosis, or Both in COVID-19?. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2020 , 26, 1076029620933953	3.3	6
14	Rivaroxaban in the treatment of venous thromboembolism and the prevention of recurrences: a practical approach. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2015 , 21, 297-308	3.3	4
13	Persistently high circulating tissue inhibitor of matrix metalloproteinase-1 levels in non-survivor brain trauma injury patients. <i>Journal of Critical Care</i> , 2019 , 51, 117-121	4	3
12	Integrating soluble biomarkers and imaging technologies in the identification of vulnerable atherosclerotic patients. <i>Biomarker Insights</i> , 2007 , 1, 165-73	3.5	3
11	[Microvascular thrombosis and clinical implications]. <i>Medicina Clínica</i> , 2021 , 156, 609-614	1	3
10	Phenotypic Screening To Discover Novel Chemical Series as Efficient Antihemorrhagic Agents. <i>ACS Medicinal Chemistry Letters</i> , 2018 , 9, 428-433	4.3	2
9	Identification of new markers of recurrence in patients with unprovoked deep vein thrombosis by gene expression profiling: the retro study. <i>European Journal of Haematology</i> , 2016 , 97, 128-36	3.8	2
8	Circulating TIMP-1 is associated with hematoma volume in patients with spontaneous intracranial hemorrhage. <i>Scientific Reports</i> , 2020 , 10, 10329	4.9	1
7	Integrating Soluble Biomarkers and Imaging Technologies in the Identification of Vulnerable Atherosclerotic Patients. <i>Biomarker Insights</i> , 2006 , 1, 117727190600100	3.5	1
6	Association of SDF1 and MMP12 with Atherosclerosis and Inflammation: Clinical and Experimental Study. <i>Life</i> , 2021 , 11,	3	1

- 5 Differences in Venous Thromboembolism Prevention and Outcome between Hospitalized Patients with Solid and Hematologic Malignancies. *TH Open*, **2019**, 3, e153-e156 2.7
- 4 Corrigendum to Preliminary characterisation of the promoter of the human p22phox gene: Identification of a new polymorphism associated with hypertension [FEBS Lett. 542 (2003) 27B1]. *FEBS Letters*, **2010**, 584, 4709-4709 3.8
- 3 Microvascular thrombosis and clinical implications. *Medicina Clínica (English Edition)*, **2021**, 156, 609-614 0.3
- 2 SP453MATRIX METALLOPROTEINASE-10 AND TISSUE INHIBITOR OF METALLOPROTEINASE-1 (TIMP-1) AS EARLY PREDICTORS OF NEPHROPATHY IN PATIENTS WITH TYPE 2 DIABETES MELLITUS. *Nephrology Dialysis Transplantation*, **2018**, 33, i500-i500 4.3
- 1 Hemostatic Biomarkers and Volumetry Help to Identify High-Risk Abdominal Aortic Aneurysms. *Life*, **2022**, 12, 823 3