
Jorge P Pinto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7899705/publications.pdf Version: 2024-02-01



LODGE D DINITO

#	Article	IF	CITATIONS
1	Physiological implications of NTBI uptake by T lymphocytes. Frontiers in Pharmacology, 2014, 5, 24.	1.6	36
2	Non-Transferrin-Bound Iron (NTBI) Uptake by T Lymphocytes: Evidence for the Selective Acquisition of Oligomeric Ferric Citrate Species. PLoS ONE, 2013, 8, e79870.	1.1	42
3	Two novel mutations in the <i>tmprss6</i> gene associated with ironâ€refractory ironâ€deficiency anaemia (irida) and partial expression in the heterozygous form. British Journal of Haematology, 2012, 158, 668-672.	1.2	24
4	ER Stress and Iron Homeostasis: A New Frontier for the UPR. Biochemistry Research International, 2011, 2011, 1-10.	1.5	18
5	A High Through-Put Screen Identifies MCP-1 (CCL2) As a Novel Regulator of Iron Homeostasis and a Modifier of Hereditary Hemochromatosis Disease Severity. Blood, 2011, 118, 685-685.	0.6	0
6	Hepcidin messenger RNA expression in human lymphocytes. Immunology, 2010, 130, 217-230.	2.0	59
7	ER Stress-Inducible Factor CHOP Affects the Expression of Hepcidin by Modulating C/EBPalpha Activity. PLoS ONE, 2009, 4, e6618.	1.1	88
8	CAT53 and HFE alleles in Alzheimer's disease: A putative protective role of the C282Y HFE mutation. Neuroscience Letters, 2009, 457, 129-132.	1.0	15
9	Protective role of calreticulin in HFE hemochromatosis. Free Radical Biology and Medicine, 2008, 44, 99-108.	1.3	17
10	Erythropoietin mediates hepcidin expression in hepatocytes through EPOR signaling and regulation of C/EBPα. Blood, 2008, 111, 5727-5733.	0.6	212
11	Overexpression of HFE in HepG2 cells reveals differences in intracellular distribution and co-localization of wt- and mutated forms. Blood Cells, Molecules, and Diseases, 2007, 39, 75-81.	0.6	10
12	A putative gene located at the MHC class I region around the D6S105 marker contributes to the setting of CD8+ T-lymphocyte numbers in humans. International Journal of Immunogenetics, 2007, 34, 359-367.	0.8	7
13	ldentification of a Promoter Element within the Zebrafish colXα1 Gene Responsive to Runx2 Isoforms Osf2/Cbfa1 and til-1 but not to pebp2αA2. Calcified Tissue International, 2006, 79, 230-244.	1.5	20
14	Osteocalcin and matrix Gla protein in zebrafish (Danio rerio) and Senegal sole (Solea senegalensis): Comparative gene and protein expression during larval development through adulthood. Gene Expression Patterns, 2006, 6, 637-652.	0.3	84
15	Efficient Screening of the Cystinuria-Related C663T Slc3a1 Nonsense Mutation in Newfoundland Dogs by Denaturing High-Performance Liquid Chromatography. Journal of Veterinary Diagnostic Investigation, 2006, 18, 102-105.	O.5	1
16	A Portuguese patient homozygous for the -25G>A mutation of the HAMP promoter shows evidence of steady-state transcription but fails to up-regulate hepcidin levels by iron. Blood, 2005, 106, 2922-2923.	0.6	30
17	Identification of a New pebp2αA2 Isoform From Zebrafishrunx2Capable of Inducing Osteocalcin Gene Expression In Vitro. Journal of Bone and Mineral Research, 2005, 20, 1440-1453.	3.1	16
18	Phox2b function in the enteric nervous system is conserved in zebrafish and is sox10-dependent. Mechanisms of Development, 2005, 122, 659-669.	1.7	126

Jorge P Pinto

#	Article	IF	CITATIONS
19	Matrix Gla protein gene expression and protein accumulation colocalize with cartilage distribution during development of the teleost fish Sparus aurata. Bone, 2003, 32, 201-210.	1.4	36
20	Cloning and characterization of the cDNA and gene encoding Xenopus laevis osteocalcin. Gene, 2002, 289, 97-107.	1.0	15
21	Cloning of the bone Gla protein gene from the teleost fish Sparus aurata. Evidence for overall conservation in gene organization and bone-specific expression from fish to man. Gene, 2001, 270, 77-91.	1.0	49