

Diego dos Santos Ferreira

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7901789/publications.pdf>

Version: 2024-02-01

19
papers

566
citations

687220

13
h-index

794469

19
g-index

19
all docs

19
docs citations

19
times ranked

1084
citing authors

#	ARTICLE	IF	CITATIONS
1	Molecular Magnetic Resonance Imaging of Liver Fibrosis and Fibrogenesis Is Not Altered by Inflammation. <i>Investigative Radiology</i> , 2021, 56, 244-251.	3.5	6
2	Virtual screening of antibacterial compounds by similarity search of Enoyl-ACP reductase (FabI) inhibitors. <i>Future Medicinal Chemistry</i> , 2020, 12, 51-68.	1.1	12
3	Targeting acid ceramidase inhibits YAP/TAZ signaling to reduce fibrosis in mice. <i>Science Translational Medicine</i> , 2020, 12, .	5.8	71
4	Peroxidase Sensitive Amplifiable Probe for Molecular Magnetic Resonance Imaging of Pulmonary Inflammation. <i>ACS Sensors</i> , 2019, 4, 2412-2419.	4.0	17
5	⁶⁸ Ga-NODAGA-Indole: An Allysine-Reactive Positron Emission Tomography Probe for Molecular Imaging of Pulmonary Fibrogenesis. <i>Journal of the American Chemical Society</i> , 2019, 141, 5593-5596.	6.6	23
6	Association of casein micelle size and enzymatic curd strength and dry matter curd yield. <i>Ciencia Rural</i> , 2019, 49, .	0.3	2
7	Encapsulation of trans -aconitic acid in mucoadhesive microspheres prolongs the anti-inflammatory effect in LPS-induced acute arthritis. <i>European Journal of Pharmaceutical Sciences</i> , 2018, 119, 112-120.	1.9	15
8	Prolonged cenicriviroc therapy reduces hepatic fibrosis despite steatohepatitis in a diet-induced mouse model of nonalcoholic steatohepatitis. <i>Hepatology Communications</i> , 2018, 2, 529-545.	2.0	43
9	Molecular magnetic resonance imaging accurately measures the antifibrotic effect of EDP-305, a novel farnesoid X receptor agonist. <i>Hepatology Communications</i> , 2018, 2, 821-835.	2.0	46
10	Evaluation of antitumor activity and cardiac toxicity of a bone-targeted pH-sensitive liposomal formulation in a bone metastasis tumor model in mice. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2017, 13, 1693-1701.	1.7	19
11	Molecular Magnetic Resonance Imaging of Lung Fibrogenesis with an Oxyamine-Based Probe. <i>Angewandte Chemie - International Edition</i> , 2017, 56, 9825-9828.	7.2	41
12	Development of a bone-targeted pH-sensitive liposomal formulation containing doxorubicin: physicochemical characterization, cytotoxicity, and biodistribution evaluation in a mouse model of bone metastasis. <i>International Journal of Nanomedicine</i> , 2016, Volume 11, 3737-3751.	3.3	31
13	Development of imaging probes for bone cancer in animal models. A systematic review. <i>Biomedicine and Pharmacotherapy</i> , 2016, 83, 1253-1264.	2.5	14
14	Noninvasive Assessment of Losartan-Induced Increase in Functional Microvasculature and Drug Delivery in Pancreatic Ductal Adenocarcinoma. <i>Translational Oncology</i> , 2016, 9, 431-437.	1.7	42
15	Alendronate-coated long-circulating liposomes containing ^{99m} technetium-ceftizoxime used to identify osteomyelitis. <i>International Journal of Nanomedicine</i> , 2015, 10, 2441.	3.3	27
16	Influence of race and crossbreeding on casein micelles size. <i>Animal Science Journal</i> , 2015, 86, 553-556.	0.6	3
17	Ursolic Acid Incorporation Does Not Prevent the Formation of a Non-lamellar Phase in pH-Sensitive and Long-Circulating Liposomes. <i>Langmuir</i> , 2014, 30, 15083-15090.	1.6	13
18	pH-sensitive liposomes for drug delivery in cancer treatment. <i>Therapeutic Delivery</i> , 2013, 4, 1099-1123.	1.2	119

#	ARTICLE	IF	CITATIONS
19	Technetium-99m-labeled ceftizoxime loaded long-circulating and pH-sensitive liposomes used to identify osteomyelitis. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2012, 22, 4605-4608.	1.0	22