

Chaowei Hu

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7907872/chaowei-hu-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

13
papers

122
citations

6
h-index

11
g-index

13
ext. papers

229
ext. citations

4.1
avg, IF

2.39
L-index

#	Paper	IF	Citations
13	Extracellular Vesicles Derived from Intermittent Hypoxia-Treated Red Blood Cells Impair Endothelial Function Through Regulating eNOS Phosphorylation and ET-1 Expression. <i>Cardiovascular Drugs and Therapy</i> , 2021 , 35, 901-913	3.9	6
12	Chronic Intermittent Hypoxia Participates in the Pathogenesis of Atherosclerosis and Perturbs the Formation of Intestinal Microbiota. <i>Frontiers in Cellular and Infection Microbiology</i> , 2021 , 11, 560201	5.9	4
11	Impact of chronic intermittent hypoxia on the long non-coding RNA and mRNA expression profiles in myocardial infarction. <i>Journal of Cellular and Molecular Medicine</i> , 2021 , 25, 421-433	5.6	2
10	Increased levels of VCAM-1 is associated with higher occurrence of coronary artery disease in adults with moderate to severe obstructive sleep apnea. <i>Sleep Medicine</i> , 2021 , 85, 131-137	4.6	1
9	p38/JNK Is Required for the Proliferation and Phenotype Changes of Vascular Smooth Muscle Cells Induced by in Essential Hypertension. <i>International Journal of Hypertension</i> , 2020 , 2020, 3123968	2.4	1
8	Increased Circulating Angiotensin-Like Protein 8 Levels Are Associated with Thoracic Aortic Dissection and Higher Inflammatory Conditions. <i>Cardiovascular Drugs and Therapy</i> , 2020 , 34, 65-77	3.9	9
7	Potential Role of mRNAs and LncRNAs in Chronic Intermittent Hypoxia Exposure-Aggravated Atherosclerosis. <i>Frontiers in Genetics</i> , 2020 , 11, 290	4.5	1
6	The Clinical Role of Angiotensin-Like Protein 3 in Evaluating Coronary Artery Disease in Patients with Obstructive Sleep Apnea. <i>Cardiovascular Drugs and Therapy</i> , 2020 , 34, 773-780	3.9	4
5	Shifts in gut microbiome and metabolome are associated with risk of recurrent atrial fibrillation. <i>Journal of Cellular and Molecular Medicine</i> , 2020 , 24, 13356-13369	5.6	11
4	AuthorsResponse to the Letter to the Editor: Increased Circulating Angiotensin-Like Protein 8 Levels Are Associated with Thoracic Aortic Dissection and Higher Inflammatory Conditions. <i>Cardiovascular Drugs and Therapy</i> , 2020 , 34, 881	3.9	
3	Angiotensin-like protein 8 accelerates atherosclerosis in ApoE mice. <i>Atherosclerosis</i> , 2020 , 307, 63-71	3.1	8
2	Disordered gut microbiota and alterations in metabolic patterns are associated with atrial fibrillation. <i>GigaScience</i> , 2019 , 8,	7.6	47
1	Dysbiotic gut microbes may contribute to hypertension by limiting vitamin D production. <i>Clinical Cardiology</i> , 2019 , 42, 710-719	3.3	28