

Kohjiro Ueki

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7910460/kohjiro-ueki-publications-by-year.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

84
papers

8,705
citations

33
h-index

93
g-index

93
ext. papers

9,775
ext. citations

9
avg, IF

5.22
L-index

#	Paper	IF	Citations
84	An antisense transcript transcribed from <i>Irs2</i> locus contributes to the pathogenesis of hepatic steatosis in insulin resistance.. <i>Cell Chemical Biology</i> , 2021 ,	8.2	2
83	Preparation and culture of bone marrow-derived macrophages from mice for functional analysis. <i>STAR Protocols</i> , 2021 , 2, 100246	1.4	15
82	MEK/ERK Signaling in β Cells Bifunctionally Regulates β Cell Mass and Glucose-Stimulated Insulin Secretion Response to Maintain Glucose Homeostasis. <i>Diabetes</i> , 2021 , 70, 1519-1535	0.9	2
81	Effect of empagliflozin on cardiorenal outcomes and mortality according to body mass index: A subgroup analysis of the EMPA-REG OUTCOME trial with a focus on Asia. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 1886-1891	6.7	3
80	Diagnosis, prevention, and treatment of cardiovascular diseases in people with type 2 diabetes and prediabetes: a consensus statement jointly from the Japanese Circulation Society and the Japan Diabetes Society. <i>Diabetology International</i> , 2021 , 12, 1-51	2.3	5
79	Hepatocellular carcinoma development in diabetic patients: a nationwide survey in Japan. <i>Journal of Gastroenterology</i> , 2021 , 56, 261-273	6.9	10
78	SLC15A4 mediates M1-prone metabolic shifts in macrophages and guards immune cells from metabolic stress. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	4
77	Comorbidities and complications in Japanese patients with type 2 diabetes mellitus: Retrospective analyses of J-DREAMS, an advanced electronic medical records database. <i>Diabetes Research and Clinical Practice</i> , 2021 , 178, 108845	7.4	0
76	Prevention of Worsening Diabetes through Behavioral Changes by an IoT-based Self-Monitoring System in Japan (PRISM-J): Study design and rationale for a multicenter, open-label, randomized parallel-group trial. <i>GHM Open</i> , 2021 , 1, 3-11		
75	Kidney Outcomes Associated With SGLT2 Inhibitors Versus Other Glucose-Lowering Drugs in Real-world Clinical Practice: The Japan Chronic Kidney Disease Database. <i>Diabetes Care</i> , 2021 , 44, 2542-2551	14.6	4
74	Long-term safety and efficacy of alogliptin, a DPP-4 inhibitor, in patients with type 2 diabetes: a 3-year prospective, controlled, observational study (J-BRAND Registry). <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	5
73	Associations between diabetes duration and self-stigma development in Japanese people with type 2 diabetes: a secondary analysis of cross-sectional data.. <i>BMJ Open</i> , 2021 , 11, e055013	3	0
72	Insulin- and Lipopolysaccharide-Mediated Signaling in Adipose Tissue Macrophages Regulates Postprandial Glycemia through Akt-mTOR Activation. <i>Molecular Cell</i> , 2020 , 79, 43-53.e4	17.6	12
71	How self-stigma affects patient activation in persons with type 2 diabetes: a cross-sectional study. <i>BMJ Open</i> , 2020 , 10, e034757	3	3
70	IL-7R-Dependent Phosphatidylinositol 3-Kinase Competes with the STAT5 Signal to Modulate T Cell Development and Homeostasis. <i>Journal of Immunology</i> , 2020 , 204, 844-857	5.3	2
69	Diagnosis, Prevention, and Treatment of Cardiovascular Diseases in People With Type 2 Diabetes and Prediabetes - A Consensus Statement Jointly From the Japanese Circulation Society and the Japan Diabetes Society. <i>Circulation Journal</i> , 2020 , 85, 82-125	2.9	7
68	8. Perspective of the Treatment for Diabetes. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2020 , 109, 1912-1918	0	

67	Comparison of effectiveness and drug cost between dipeptidyl peptidase-4 inhibitor and biguanide as the first-line anti-hyperglycaemic medication among Japanese working generation with type 2 diabetes. <i>Journal of Evaluation in Clinical Practice</i> , 2020 , 26, 299-307	2.5	4
66	Changes in the quality of diabetes care in Japan between 2007 and 2015: A repeated cross-sectional study using claims data. <i>Diabetes Research and Clinical Practice</i> , 2019 , 149, 188-199	7.4	6
65	Hepatocellular carcinoma as a leading cause of cancer-related deaths in Japanese type 2 diabetes mellitus patients. <i>Journal of Gastroenterology</i> , 2019 , 54, 64-77	6.9	14
64	Variation in process quality measures of diabetes care by region and institution in Japan during 2015-2016: An observational study of nationwide claims data. <i>Diabetes Research and Clinical Practice</i> , 2019 , 155, 107750	7.4	11
63	3. Recent Progress in the Treatment of Type 2 Diabetes. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2019 , 108, 460-467	0	
62	Hepatic Sdf2l1 controls feeding-induced ER stress and regulates metabolism. <i>Nature Communications</i> , 2019 , 10, 947	17.4	28
61	Hepatic IRS1 and Ectenin expression is associated with histological progression and overt diabetes emergence in NAFLD patients. <i>Journal of Gastroenterology</i> , 2018 , 53, 1261-1275	6.9	16
60	The RNA Methyltransferase Complex of WTAP, METTL3, and METTL14 Regulates Mitotic Clonal Expansion in Adipogenesis. <i>Molecular and Cellular Biology</i> , 2018 , 38,	4.8	65
59	Activin B: A potential target to cure diabetes. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society</i> , 2018 , WCP2018, SY62-4	0	
58	Downregulation of macrophage Irs2 by hyperinsulinemia impairs IL-4-induced M2a-subtype macrophage activation in obesity. <i>Nature Communications</i> , 2018 , 9, 4863	17.4	27
57	Thermographic findings in a case of type 2 diabetes with foot ulcer due to callus deterioration. <i>Diabetology International</i> , 2017 , 8, 328-333	2.3	5
56	Causes of death in Japanese patients with diabetes based on the results of a survey of 45,708 cases during 2001-2010: Report of the Committee on Causes of Death in Diabetes Mellitus. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 397-410	3.9	76
55	Dual Regulation of Gluconeogenesis by Insulin and Glucose in the Proximal Tubules of the Kidney. <i>Diabetes</i> , 2017 , 66, 2339-2350	0.9	44
54	Psychological and behavioural patterns of stigma among patients with type 2 diabetes: a cross-sectional study. <i>BMJ Open</i> , 2017 , 7, e013425	3	19
53	New glycemic targets for patients with diabetes from the Japan Diabetes Society. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 123-125	3.9	33
52	Effect of an intensified multifactorial intervention on cardiovascular outcomes and mortality in type 2 diabetes (J-DOIT3): an open-label, randomised controlled trial. <i>Lancet Diabetes and Endocrinology</i> , 2017 , 5, 951-964	18.1	141
51	CD206 M2-like macrophages regulate systemic glucose metabolism by inhibiting proliferation of adipocyte progenitors. <i>Nature Communications</i> , 2017 , 8, 286	17.4	116
50	Design of and rationale for the Japan Diabetes comprehensive database project based on an Advanced electronic Medical record System (J-DREAMS). <i>Diabetology International</i> , 2017 , 8, 375-382	2.3	19

49	Causes of death in Japanese patients with diabetes based on the results of a survey of 45,708 cases during 2001-2010: report of Committee on Causes of Death in Diabetes Mellitus. <i>Diabetology International</i> , 2017 , 8, 117-136	2.3	6
48	New glyceic targets for patients with diabetes from the Japan Diabetes Society. <i>Diabetology International</i> , 2016 , 7, 327-330	2.3	3
47	Differential hepatic distribution of insulin receptor substrates causes selective insulin resistance in diabetes and obesity. <i>Nature Communications</i> , 2016 , 7, 12977	17.4	51
46	Pioglitazone Ameliorates Smooth Muscle Cell Proliferation in Cuff-Induced Neointimal Formation by Both Adiponectin-Dependent and -Independent Pathways. <i>Scientific Reports</i> , 2016 , 6, 34707	4.9	5
45	Design of and rationale for the Japan Diabetes Optimal Integrated Treatment study for 3 major risk factors of cardiovascular diseases (J-DOIT3): a multicenter, open-label, randomized, parallel-group trial. <i>BMJ Open Diabetes Research and Care</i> , 2016 , 4, e000123	4.5	19
44	Report of the Japan Diabetes Society (JDS)/Japanese Cancer Association (JCA) Joint Committee on Diabetes and Cancer, Second Report. <i>Diabetology International</i> , 2016 , 7, 12-15	2.3	7
43	A qualitative study on the impact of internalized stigma on type 2 diabetes self-management. <i>Patient Education and Counseling</i> , 2016 , 99, 1233-1239	3.1	28
42	Association between self-stigma and self-care behaviors in patients with type 2 diabetes: a cross-sectional study. <i>BMJ Open Diabetes Research and Care</i> , 2016 , 4, e000156	4.5	21
41	Factors Associated With Callus in Patients with Diabetes, Focused on Plantar Shear Stress During Gait. <i>Journal of Diabetes Science and Technology</i> , 2016 , 10, 1353-1359	4.1	17
40	Tofogliflozin Improves Insulin Resistance in Skeletal Muscle and Accelerates Lipolysis in Adipose Tissue in Male Mice. <i>Endocrinology</i> , 2016 , 157, 1029-42	4.8	90
39	Report of the Japan diabetes society/Japanese cancer association joint committee on diabetes and cancer, Second report. <i>Cancer Science</i> , 2016 , 107, 369-71	6.9	6
38	A large-scale, observational study to investigate the current status of diabetes complications and their prevention in Japan: research outline and baseline data for type 2 diabetes JDCP study 1. <i>Diabetology International</i> , 2015 , 6, 243-251	2.3	12
37	Insulin receptor substrate-2 (Irs2) in endothelial cells plays a crucial role in insulin secretion. <i>Diabetes</i> , 2015 , 64, 876-86	0.9	28
36	Effects of beraprost sodium, an oral prostacyclin analog, on insulin resistance in patients with type 2 diabetes. <i>Diabetology International</i> , 2015 , 6, 39-45	2.3	1
35	Perspective of Small-Molecule AdipoR Agonist for Type 2 Diabetes and Short Life in Obesity. <i>Diabetes and Metabolism Journal</i> , 2015 , 39, 363-72	5	43
34	Association between Washing Residue on the Feet and Tinea Pedis in Diabetic Patients. <i>Nursing Research and Practice</i> , 2015 , 2015, 872678	1.9	3
33	Lung abscess without sepsis in a patient with diabetes with refractory episodes of spontaneous hypoglycemia: a case report and review of the literature. <i>Journal of Medical Case Reports</i> , 2014 , 8, 51	1.2	2
32	Protocol for a large-scale prospective observational study with alogliptin in patients with type 2 diabetes: J-BRAND Registry. <i>BMJ Open</i> , 2014 , 4, e004760	3	4

31	Effect of renal impairment on the pharmacokinetics, pharmacodynamics, and safety of empagliflozin, a sodium glucose cotransporter 2 inhibitor, in Japanese patients with type 2 diabetes mellitus. <i>Clinical Therapeutics</i> , 2014 , 36, 1606-15	3.5	22
30	Report of the JDS/JCA Joint Committee on Diabetes and Cancer. <i>Diabetology International</i> , 2013 , 4, 81-96	3	26
29	Insulin regulates liver metabolism in vivo in the absence of hepatic Akt and Foxo1. <i>Nature Medicine</i> , 2012 , 18, 388-95	50.5	260
28	The PREDICTIVETM Study: a multinational, prospective observational study to evaluate the safety and efficacy of insulin detemir treatment in patients with type 1 and 2 diabetes—data from the Japan cohort. <i>Diabetology International</i> , 2012 , 3, 11-20	2.3	1
27	International clinical harmonization of glycated hemoglobin in Japan: From Japan Diabetes Society to National Glycohemoglobin Standardization Program values. <i>Diabetology International</i> , 2012 , 3, 8-10	2.3	192
26	Loss of Akt1 in mice increases energy expenditure and protects against diet-induced obesity. <i>Molecular and Cellular Biology</i> , 2012 , 32, 96-106	4.8	44
25	Adiponectin enhances insulin sensitivity by increasing hepatic IRS-2 expression via a macrophage-derived IL-6-dependent pathway. <i>Cell Metabolism</i> , 2011 , 13, 401-412	24.6	197
24	Long-term safety and efficacy of exenatide twice daily in Japanese patients with suboptimally controlled type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2011 , 2, 448-56	3.9	12
23	Blockade of class IB phosphoinositide-3 kinase ameliorates obesity-induced inflammation and insulin resistance. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 5753-8	11.5	36
22	Class IA phosphatidylinositol 3-kinase in pancreatic β cells controls insulin secretion by multiple mechanisms. <i>Cell Metabolism</i> , 2010 , 12, 619-32	24.6	84
21	Report of the committee on the classification and diagnostic criteria of diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2010 , 1, 212-28	3.9	953
20	Report of the Committee on the classification and diagnostic criteria of diabetes mellitus. <i>Diabetology International</i> , 2010 , 1, 2-20	2.3	243
19	Glucose effects on beta-cell growth and survival require activation of insulin receptors and insulin receptor substrate 2. <i>Molecular and Cellular Biology</i> , 2009 , 29, 3219-28	4.8	116
18	CD8+ effector T cells contribute to macrophage recruitment and adipose tissue inflammation in obesity. <i>Nature Medicine</i> , 2009 , 15, 914-20	50.5	1567
17	Type 1 Diabetes Mellitus Associated with Vogt-Koyanagi-Harada Syndrome, Palmoplantar Pustulosis, and Hashimoto's Thyroiditis. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2009 , 98, 1369-1371	0	
16	Dynamic functional relay between insulin receptor substrate 1 and 2 in hepatic insulin signaling during fasting and feeding. <i>Cell Metabolism</i> , 2008 , 8, 49-64	24.6	172
15	Overexpression of monocyte chemoattractant protein-1 in adipose tissues causes macrophage recruitment and insulin resistance. <i>Journal of Biological Chemistry</i> , 2006 , 281, 26602-14	5.4	638
14	Total insulin and IGF-I resistance in pancreatic beta cells causes overt diabetes. <i>Nature Genetics</i> , 2006 , 38, 583-8	36.3	217

13	Role of suppressors of cytokine signaling SOCS-1 and SOCS-3 in hepatic steatosis and the metabolic syndrome. <i>Hepatology Research</i> , 2005 , 33, 185-92	5.1	74
12	Central role of suppressors of cytokine signaling proteins in hepatic steatosis, insulin resistance, and the metabolic syndrome in the mouse. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 10422-7	11.5	310
11	Suppressor of cytokine signaling 1 (SOCS-1) and SOCS-3 cause insulin resistance through inhibition of tyrosine phosphorylation of insulin receptor substrate proteins by discrete mechanisms. <i>Molecular and Cellular Biology</i> , 2004 , 24, 5434-46	4.8	517
10	Increased insulin sensitivity in mice lacking p85beta subunit of phosphoinositide 3-kinase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 419-24	11.5	209
9	Molecular balance between the regulatory and catalytic subunits of phosphoinositide 3-kinase regulates cell signaling and survival. <i>Molecular and Cellular Biology</i> , 2002 , 22, 965-77	4.8	230
8	Restored insulin-sensitivity in IRS-1-deficient mice treated by adenovirus-mediated gene therapy. <i>Journal of Clinical Investigation</i> , 2000 , 105, 1437-45	15.9	44
7	The mechanism of insulin-induced signal transduction mediated by the insulin receptor substrate family. <i>Endocrine Journal</i> , 1999 , 46, S25-34	2.9	35
6	Vascular endothelial growth factor (VEGF) activates Raf-1, mitogen-activated protein (MAP) kinases, and S6 kinase (p90rsk) in cultured rat cardiac myocytes. <i>Journal of Cellular Physiology</i> , 1998 , 175, 239-46	7	39
5	Growth hormone-induced tyrosine phosphorylation of EGF receptor as an essential element leading to MAP kinase activation and gene expression. <i>Endocrine Journal</i> , 1998 , 45 Suppl, S27-31	2.9	47
4	Vascular endothelial growth factor (VEGF) activates Raf-1, mitogen-activated protein (MAP) kinases, and S6 kinase (p90rsk) in cultured rat cardiac myocytes 1998 , 175, 239		1
3	Tyrosine phosphorylation of the EGF receptor by the kinase Jak2 is induced by growth hormone. <i>Nature</i> , 1997 , 390, 91-6	50.4	252
2	Angiotensin II partly mediates mechanical stress-induced cardiac hypertrophy. <i>Circulation Research</i> , 1995 , 77, 258-65	15.7	204
1	Insulin resistance and growth retardation in mice lacking insulin receptor substrate-1. <i>Nature</i> , 1994 , 372, 182-6	50.4	914