

Angeles Ramos-BarrÃ³n

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7917186/publications.pdf>

Version: 2024-02-01

16
papers

192
citations

1163117

8
h-index

1058476

14
g-index

18
all docs

18
docs citations

18
times ranked

397
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevention of murine lupus disease in (NZB \times NZW)F1 mice by sirolimus treatment. <i>Lupus</i> , 2007, 16, 775-781.	1.6	48
2	Urine metabolomics insight into acute kidney injury point to oxidative stress disruptions in energy generation and H ₂ S availability. <i>Journal of Molecular Medicine</i> , 2017, 95, 1399-1409.	3.9	30
3	Urinary Kininogen-1 and Retinol binding protein-4 respond to Acute Kidney Injury: predictors of patient prognosis?. <i>Scientific Reports</i> , 2016, 6, 19667.	3.3	20
4	Long-Term Mortality among Hospitalized Non-ICU Patients with Acute Kidney Injury Referred to Nephrology. <i>Nephron</i> , 2015, 131, 23-33.	1.8	14
5	B-Cell α Activating Factor Levels Are Associated With Antibody-Mediated Histological Damage in Kidney Transplantation. <i>Transplantation Proceedings</i> , 2016, 48, 2910-2912.	0.6	11
6	Combined use of GM2AP and TCP1-eta urinary levels predicts recovery from intrinsic acute kidney injury. <i>Scientific Reports</i> , 2020, 10, 11599.	3.3	11
7	Analysis of Urinary Gene Expression of Epithelial-Mesenchymal Transition Markers in Kidney Transplant Recipients. <i>Transplantation Proceedings</i> , 2010, 42, 2886-2888.	0.6	10
8	Autoimmune syndrome after induction of neonatal tolerance to I-E antigens. <i>European Journal of Immunology</i> , 1993, 23, 2353-2357.	2.9	8
9	Urinary TCP1-eta: A Cortical Damage Marker for the Pathophysiological Diagnosis and Prognosis of Acute Kidney Injury. <i>Toxicological Sciences</i> , 2020, 174, 3-15.	3.1	8
10	Relation of Urinary Gene Expression of Epithelial-Mesenchymal Transition Markers With Initial Events and 1-Year Kidney Graft Function. <i>Transplantation Proceedings</i> , 2012, 44, 2573-2576.	0.6	6
11	Assessment of Kidney Graft Function Evolution Measured by Creatinine and Cystatin C. <i>Transplantation Proceedings</i> , 2016, 48, 2913-2916.	0.6	5
12	Urinary Spermidine Predicts and Associates with In-Hospital Acute Kidney Injury after Cardiac Surgery. <i>Antioxidants</i> , 2021, 10, 896.	5.1	4
13	Urinary KIM-1 Correlates with the Subclinical Sequelae of Tubular Damage Persisting after the Apparent Functional Recovery from Intrinsic Acute Kidney Injury. <i>Biomedicines</i> , 2022, 10, 1106.	3.2	4
14	Urinary Plasminogen Activator Inhibitor-1: A Biomarker of Acute Tubular Injury. <i>American Journal of Nephrology</i> , 2021, 52, 714-724.	3.1	2
15	Anti-LFA-1 (CD11a) monoclonal antibody interferes with neonatal induction of tolerance to alloantigens. <i>European Journal of Immunology</i> , 1994, 24, 985-990.	2.9	1
16	CD4 ⁺ T cells determine the ability of spleen cells from F1 hybrid mice to induce neonatal tolerance to alloantigens and autoimmunity in parental mice. <i>European Journal of Immunology</i> , 1995, 25, 1760-1764.	2.9	1