

Moufida Abdennadher

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7919120/publications.pdf>

Version: 2024-02-01

7
papers

105
citations

1937685

4
h-index

1720034

7
g-index

7
all docs

7
docs citations

7
times ranked

119
citing authors

#	ARTICLE	IF	CITATIONS
1	Characterization of <i>Coolia</i> spp. (Gonyaucales, Dinophyceae) from Southern Tunisia: first record of <i>Coolia malayensis</i> in the Mediterranean Sea. <i>Algae</i> , 2021, 36, 175-193.	2.3	5
2	A long-term study on <i>Coolia monotis</i> distribution from the south-east Mediterranean Sea. <i>Continental Shelf Research</i> , 2020, 211, 104267.	1.8	3
3	Dinoflagellates encystment with emphasis on blooms in Boughrara Lagoon (South-Western) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Estuarine, Coastal and Shelf Science, 2020, 237, 106648.	2.1	8
4	A Naïve Bayesian network approach to determine the potential drivers of the toxic dinoflagellate <i>Coolia monotis</i> (Meunier, 1919) in the Gulf of Gabès, Tunisia. <i>Euro-Mediterranean Journal for Environmental Integration</i> , 2019, 4, 1.	1.3	2
5	Controlling factors of harmful microalgae distribution in water column, biofilm and sediment in shellfish production area (South of Sfax, Gulf of Gabes) from southern Tunisia. <i>Continental Shelf Research</i> , 2018, 152, 61-70.	1.8	10
6	<i>Ostreopsis cf. ovata</i> in the Gulf of Gabès (south-eastern Mediterranean Sea): morphological, molecular and ecological characterization. <i>Harmful Algae</i> , 2017, 63, 56-67.	4.8	18
7	What are the potential drivers of blooms of the toxic dinoflagellate <i>Karenia selliformis</i> ? A 10-year study in the Gulf of Gabes, Tunisia, southwestern Mediterranean Sea. <i>Harmful Algae</i> , 2013, 23, 8-18.	4.8	59