

Mark Barbour

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7921706/publications.pdf>

Version: 2024-02-01

10
papers

264
citations

1307594

7
h-index

1474206

9
g-index

10
all docs

10
docs citations

10
times ranked

569
citing authors

#	ARTICLE	IF	CITATIONS
1	Interleukin-33 predicts poor prognosis and promotes ovarian cancer cell growth and metastasis through regulating ERK and JNK signaling pathways. <i>Molecular Oncology</i> , 2016, 10, 113-125.	4.6	81
2	IL-33 attenuates the development of experimental autoimmune uveitis. <i>European Journal of Immunology</i> , 2014, 44, 3320-3329.	2.9	64
3	Expression and Function of IL-33/ST2 Axis in the Central Nervous System Under Normal and Diseased Conditions. <i>Frontiers in Immunology</i> , 2018, 9, 2596.	4.8	62
4	Increased Levels of IL-16 in the Central Nervous System during Neuroinflammation Are Associated with Infiltrating Immune Cells and Resident Glial Cells. <i>Biology</i> , 2021, 10, 472.	2.8	19
5	MAP kinase phosphatase 2 deficient mice develop attenuated experimental autoimmune encephalomyelitis through regulating dendritic cells and T cells. <i>Scientific Reports</i> , 2016, 6, 38999.	3.3	9
6	The therapeutic effect of anti-CD52 treatment in murine experimental autoimmune encephalomyelitis is associated with altered IL-33 and ST2 expression levels. <i>Journal of Neuroimmunology</i> , 2018, 318, 87-96.	2.3	9
7	Effect of sphingosine kinase modulators on interleukin-1 β release, sphingosine 1-phosphate receptor 1 expression and experimental autoimmune encephalomyelitis. <i>British Journal of Pharmacology</i> , 2017, 174, 210-222.	5.4	8
8	Role of IL-33/ST2 signaling pathway in systemic sclerosis and other fibrotic diseases. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 119, 141-146.	0.8	8
9	Anti-CD52 antibody treatment in murine experimental autoimmune encephalomyelitis induces dynamic and differential modulation of innate immune cells in peripheral immune and central nervous systems. <i>Immunology</i> , 2021, , .	4.4	3
10	Protease-activated receptor 2 activation induces behavioural changes associated with depression-like behaviour through microglial-independent modulation of inflammatory cytokines. <i>Psychopharmacology</i> , 2022, 239, 229-242.	3.1	1