

Minako Wakasugi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7925924/publications.pdf>

Version: 2024-02-01

53
papers

887
citations

516710

16
h-index

477307

29
g-index

53
all docs

53
docs citations

53
times ranked

1202
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Polypharmacy, chronic kidney disease, and incident fragility fracture: a prospective cohort study. <i>Journal of Bone and Mineral Metabolism</i> , 2022, 40, 157-166. | 2.7 | 11 |
| 2 | Sex Differences in Cause-specific Mortality in Japanese Dialysis Patients. <i>Internal Medicine</i> , 2022, 61, 2831-2839. | 0.7 | 4 |
| 3 | Healthy Lifestyle and Incident Hypertension and Diabetes in Participants with and without Chronic Kidney Disease: The Japan Specific Health Checkups (J-SHC) Study. <i>Internal Medicine</i> , 2022, 61, 2841-2851. | 0.7 | 6 |
| 4 | Authors'™ reply. <i>Journal of Bone and Mineral Metabolism</i> , 2022, , 1. | 2.7 | 0 |
| 5 | Stroke incidence and chronic kidney disease: A hospital-based prospective cohort study. <i>Nephrology</i> , 2022, 27, 577-587. | 1.6 | 3 |
| 6 | Hematuria, bone health, and cardiovascular mortality. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 562-563. | 1.6 | 0 |
| 7 | The Effect of CKD on Associations between Lifestyle Factors and All-cause, Cancer, and Cardiovascular Mortality: A Population-based Cohort Study. <i>Internal Medicine</i> , 2021, 60, 2189-2200. | 0.7 | 6 |
| 8 | Smoking and risk of fractures requiring hospitalization in hemodialysis patients: a nationwide cohort study in Japan. <i>Nephrology Dialysis Transplantation</i> , 2021, , . | 0.7 | 1 |
| 9 | Secular trends in end-stage kidney disease requiring renal replacement therapy in Japan: Japanese Society of Dialysis Therapy Registry data from 1983 to 2016. <i>Nephrology</i> , 2020, 25, 172-178. | 1.6 | 10 |
| 10 | Association between serum IgG antibody titers against <i>Porphyromonas gingivalis</i> and liver enzyme levels: A cross-sectional study in Sado Island. <i>Heliyon</i> , 2020, 6, e05531. | 3.2 | 3 |
| 11 | Long-term excess mortality after hip fracture in hemodialysis patients: a nationwide cohort study in Japan. <i>Journal of Bone and Mineral Metabolism</i> , 2020, 38, 718-729. | 2.7 | 9 |
| 12 | Lifetime and age-conditional risk estimates of end-stage kidney disease requiring maintenance dialysis in Japan. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 518-525. | 1.6 | 3 |
| 13 | The Association between Earlobe Creases and Cardiovascular Events in Japanese Hemodialysis Patients: A Prospective Cohort Study. <i>Internal Medicine</i> , 2020, 59, 927-932. | 0.7 | 2 |
| 14 | Functional impairment attenuates the association between high serum phosphate and mortality in dialysis patients: a nationwide cohort study. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 1207-1216. | 0.7 | 14 |
| 15 | Is the population of Sado Island genetically close to the population of western Japan?. <i>Human Genome Variation</i> , 2019, 6, 26. | 0.7 | 0 |
| 16 | Hemodialysis Product and Hip Fracture in Hemodialysis Patients: A Nationwide Cohort Study in Japan. <i>Therapeutic Apheresis and Dialysis</i> , 2019, 23, 507-517. | 0.9 | 4 |
| 17 | Age-related Hearing Loss Is Strongly Associated With Cognitive Decline Regardless of the APOE4 Polymorphism. <i>Otology and Neurotology</i> , 2019, 40, 1263-1267. | 1.3 | 17 |
| 18 | Emotional disturbance assessed by the Self-Rating Depression Scale test is associated with mortality among Japanese Hemodialysis patients. <i>Fukushima Journal of Medical Sciences</i> , 2018, 64, 23-29. | 0.4 | 3 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Premature mortality due to nephrotic syndrome and the trend in nephrotic syndrome mortality in Japan, 1995â€“2014. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 55-60. | 1.6 | 8 |
| 20 | Hip Fracture Trends in Japanese Dialysis Patients, 2008-2013. <i>American Journal of Kidney Diseases</i> , 2018, 71, 173-181. | 1.9 | 22 |
| 21 | Parathyroid Hormone and Bone in Dialysis Patients. <i>Therapeutic Apheresis and Dialysis</i> , 2018, 22, 229-235. | 0.9 | 9 |
| 22 | Self-reported Slower Eating Is Associated with a Lower Salt Intake: A Population-based Cross-sectional Study. <i>Internal Medicine</i> , 2018, 57, 1561-1567. | 0.7 | 1 |
| 23 | Hyperphosphatemia is not significantly associated with increased all-cause mortality in Korean hemodialysis patients. <i>Kidney Research and Clinical Practice</i> , 2018, 37, 420-421. | 2.2 | 0 |
| 24 | Association between dialysis treatment and cognitive decline: A study from the Project in Sado for Total Health (PROST), Japan. <i>Geriatrics and Gerontology International</i> , 2017, 17, 1584-1587. | 1.5 | 9 |
| 25 | Prevalence of Earlobe Creases and Their Association With History of Cardiovascular Disease in Patients Undergoing Hemodialysis: A Cross-sectional Study. <i>Therapeutic Apheresis and Dialysis</i> , 2017, 21, 478-484. | 0.9 | 6 |
| 26 | Association between Overall Lifestyle Changes and the Incidence of Proteinuria: A Population-based, Cohort Study. <i>Internal Medicine</i> , 2017, 56, 1475-1484. | 0.7 | 38 |
| 27 | Elevated C-Reactive Protein Is Associated with Cognitive Decline in Outpatients of a General Hospital: The Project in Sado for Total Health (PROST). <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2016, 6, 10-19. | 1.3 | 48 |
| 28 | Mortality trends among Japanese dialysis patients, 1988â€“2013: a joinpoint regression analysis. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, 1501-1507. | 0.7 | 44 |
| 29 | Modifiable Factors Associated with Cognitive Impairment in 1,143 Japanese Outpatients: The Project in Sado for Total Health (PROST). <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2016, 6, 341-349. | 1.3 | 29 |
| 30 | Age- and gender-specific incidence rates of renal replacement therapy in Japan: an international comparison. <i>Renal Replacement Therapy</i> , 2016, 2, . | 0.7 | 8 |
| 31 | Title is missing!. <i>Nihon Toseki Igakkai Zasshi</i> , 2016, 49, 792-794. | 0.1 | 0 |
| 32 | Associations between the Intake of Miso Soup and Japanese Pickles and the Estimated 24-hour Urinary Sodium Excretion: A Population-based Cross-sectional Study. <i>Internal Medicine</i> , 2015, 54, 903-910. | 0.7 | 19 |
| 33 | Higher Mortality Due to Intracerebral Hemorrhage in Dialysis Patients: A Comparison with the General Population in Japan. <i>Therapeutic Apheresis and Dialysis</i> , 2015, 19, 45-49. | 0.9 | 21 |
| 34 | Anticipated Increase in the Number of Patients Who Require Dialysis Treatment Among the Aging Population of Japan. <i>Therapeutic Apheresis and Dialysis</i> , 2015, 19, 201-206. | 0.9 | 27 |
| 35 | Autologous blood transfusion for hemodialysis patients: A case report and review of clinical reports and therapeutic features. <i>Transfusion and Apheresis Science</i> , 2015, 52, 204-207. | 1.0 | 1 |
| 36 | Association between Hypouricemia and Reduced Kidney Function: A Cross-Sectional Population-Based Study in Japan. <i>American Journal of Nephrology</i> , 2015, 41, 138-146. | 3.1 | 59 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Association between Combined Lifestyle Factors and Non-Restorative Sleep in Japan: A Cross-Sectional Study Based on a Japanese Health Database. PLoS ONE, 2014, 9, e108718. | 2.5 | 52 |
| 38 | High rates of death and hospitalization follow bone fracture among hemodialysis patients. Kidney International, 2014, 86, 649. | 5.2 | 5 |
| 39 | A hip fracture in a dialysis patient with β_2 M amyloidosis. Kidney International, 2014, 85, 214-215. | 5.2 | 4 |
| 40 | Regional Variation in Hip Fracture Incidence Among Japanese Hemodialysis Patients. Therapeutic Apheresis and Dialysis, 2014, 18, 162-166. | 0.9 | 15 |
| 41 | Association between warfarin use and incidence of ischemic stroke in Japanese hemodialysis patients with chronic sustained atrial fibrillation: a prospective cohort study. Clinical and Experimental Nephrology, 2014, 18, 662-669. | 1.6 | 53 |
| 42 | Intracerebral hemorrhage was the highest cause of mortality among stroke subtypes in Japanese dialysis patients. Hemodialysis International, 2014, 18, 848-849. | 0.9 | 0 |
| 43 | Increased risk of hip fracture among Japanese hemodialysis patients. Journal of Bone and Mineral Metabolism, 2013, 31, 315-321. | 2.7 | 106 |
| 44 | Cause-Specific Excess Mortality Among Dialysis Patients: Comparison With the General Population in Japan. Therapeutic Apheresis and Dialysis, 2013, 17, 298-304. | 0.9 | 54 |
| 45 | A combination of healthy lifestyle factors is associated with a decreased incidence of chronic kidney disease: a population-based cohort study. Hypertension Research, 2013, 36, 328-333. | 2.7 | 63 |
| 46 | 71/4 Ž ā,āĈĀ>1/2ā@éĈēžē,ēēĈ...ā«āšāā,ā,,ŸæŸ“ç—†æ»ā°jçŽ† āĈ• ā,Ĉē^-ā1/2æ°ā“ā@æ”ē1/4f āĈ. Nihon Toseki Igakkai Zasshi, 2013, 46, | | |
| 47 | Differences in the local and national prevalences of chronic kidney disease based on annual health check program data. Clinical and Experimental Nephrology, 2012, 16, 749-754. | 1.6 | 1 |
| 48 | Weight gain after 20 years of age is associated with prevalence of chronic kidney disease. Clinical and Experimental Nephrology, 2012, 16, 259-268. | 1.6 | 20 |
| 49 | Use of Japanese Society for Dialysis Therapy Dialysis Tables to Compare the Local and National Incidence of Dialysis. Therapeutic Apheresis and Dialysis, 2012, 16, 63-67. | 0.9 | 7 |
| 50 | High Mortality Rate of Infectious Diseases in Dialysis Patients: A Comparison With the General Population in Japan. Therapeutic Apheresis and Dialysis, 2012, 16, 226-231. | 0.9 | 60 |
| 51 | Clinical value of ^{99m}Tc -MIBI scintigraphy in patients with secondary hyperparathyroidism. A systematic review and meta-analysis of observational studies in Japan. Nihon Toseki Igakkai Zasshi, 2011, 44, 65-72. | 0.1 | 0 |
| 52 | A case of collagenous colitis on chronic hemodialysis. Nihon Toseki Igakkai Zasshi, 2010, 43, 999-1003. | 0.1 | 1 |
| 53 | Impact of adherence to CKD-MBD guidelines on mortality among Japanese dialysis patients: an ecological study. Journal of Bone and Mineral Metabolism, 0, , . | 2.7 | 0 |