

Emmanuel Itti

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7936763/emmanuel-itti-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

118
papers

4,236
citations

34
h-index

63
g-index

127
ext. papers

5,142
ext. citations

5
avg, IF

4.9
L-index

#	Paper	IF	Citations
118	[18F]fluoro-2-deoxy-D-glucose positron emission tomography (FDG-PET) in aggressive lymphoma: an early prognostic tool for predicting patient outcome. <i>Blood</i> , 2005 , 106, 1376-81	2.2	418
117	Efficacy and safety of selective internal radiotherapy with yttrium-90 resin microspheres compared with sorafenib in locally advanced and inoperable hepatocellular carcinoma (SARAH): an open-label randomised controlled phase 3 trial. <i>Lancet Oncology</i> , 2017 , 18, 1624-1636	21.7	400
116	Early 18F-FDG PET for prediction of prognosis in patients with diffuse large B-cell lymphoma: SUV-based assessment versus visual analysis. <i>Journal of Nuclear Medicine</i> , 2007 , 48, 1626-32	8.9	307
115	An international confirmatory study of the prognostic value of early PET/CT in diffuse large B-cell lymphoma: comparison between Deauville criteria and SUVmax. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2013 , 40, 1312-20	8.8	195
114	Prognostic value of interim 18F-FDG PET in patients with diffuse large B-Cell lymphoma: SUV-based assessment at 4 cycles of chemotherapy. <i>Journal of Nuclear Medicine</i> , 2009 , 50, 527-33	8.9	192
113	Pretherapy metabolic tumour volume is an independent predictor of outcome in patients with diffuse large B-cell lymphoma. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2014 , 41, 2017-22	8.8	135
112	Whole-body diffusion-weighted magnetic resonance imaging with apparent diffusion coefficient mapping for staging patients with diffuse large B-cell lymphoma. <i>European Radiology</i> , 2010 , 20, 2027-38 ⁸		134
111	Interim [18F]fluorodeoxyglucose positron emission tomography scan in diffuse large B-cell lymphoma treated with anthracycline-based chemotherapy plus rituximab. <i>Journal of Clinical Oncology</i> , 2012 , 30, 184-90	2.2	130
110	Metabolic tumour volumes measured at staging in lymphoma: methodological evaluation on phantom experiments and patients. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2014 , 41, 1113-22	8.8	114
109	Personalised versus standard dosimetry approach of selective internal radiation therapy in patients with locally advanced hepatocellular carcinoma (DOSISPHERE-01): a randomised, multicentre, open-label phase 2 trial. <i>The Lancet Gastroenterology and Hepatology</i> , 2021 , 6, 17-29	18.8	82
108	Metastatic renal cell carcinoma: relationship between initial metastasis hypoxia, change after 1 month's sunitinib, and therapeutic response: an 18F-fluoromisonidazole PET/CT study. <i>Journal of Nuclear Medicine</i> , 2011 , 52, 1048-55	8.9	76
107	Automatic lesion detection and segmentation of 18F-FET PET in gliomas: A full 3D U-Net convolutional neural network study. <i>PLoS ONE</i> , 2018 , 13, e0195798	3.7	73
106	Improvement of early 18F-FDG PET interpretation in diffuse large B-cell lymphoma: importance of the reference background. <i>Journal of Nuclear Medicine</i> , 2010 , 51, 1857-62	8.9	71
105	Optimization of aerosol deposition by pressure support in children with cystic fibrosis: an experimental and clinical study. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2000 , 162, 2265-71	10.2	69
104	Whole-body diffusion-weighted imaging with apparent diffusion coefficient mapping for treatment response assessment in patients with diffuse large B-cell lymphoma: pilot study. <i>Investigative Radiology</i> , 2011 , 46, 341-9	10.1	64
103	Lymph node imaging: basic principles. <i>European Journal of Radiology</i> , 2006 , 58, 338-44	4.7	64
102	Usefulness of (99m)Tc-HMDP scintigraphy for the etiologic diagnosis and prognosis of cardiac amyloidosis. <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis</i> , 2015 , 22, 210-20	2.7	63

101	FDG PET/CT imaging as a biomarker in lymphoma. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2015 , 42, 623-33	8.8	59
100	Bone marrow involvement in diffuse large B-cell lymphoma: correlation between FDG-PET uptake and type of cellular infiltrate. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2009 , 36, 745-50	8.8	59
99	Sentinel node localization should be interpreted with caution in midline vulvar cancer. <i>Gynecologic Oncology</i> , 2005 , 97, 151-4	4.9	59
98	Total metabolic tumor volume, circulating tumor cells, cell-free DNA: distinct prognostic value in follicular lymphoma. <i>Blood Advances</i> , 2018 , 2, 807-816	7.8	56
97	Validation of a standardized normalization template for statistical parametric mapping analysis of 123I-FP-CIT images. <i>Journal of Nuclear Medicine</i> , 2007 , 48, 1459-67	8.9	53
96	Phase II study of sirolimus in treatment-naive patients with advanced hepatocellular carcinoma. <i>Digestive and Liver Disease</i> , 2012 , 44, 610-6	3.3	49
95	Development and application of a real-time on-line blinded independent central review of interim PET scans to determine treatment allocation in lymphoma trials. <i>Journal of Clinical Oncology</i> , 2009 , 27, 2739-41	2.2	48
94	[18F]-NaF PET/CT imaging in cardiac amyloidosis. <i>Journal of Nuclear Cardiology</i> , 2016 , 23, 846-9	2.1	46
93	Whole-Body Diffusion-weighted Imaging in Hodgkin Lymphoma and Diffuse Large B-Cell Lymphoma. <i>Radiographics</i> , 2015 , 35, 747-64	5.4	42
92	Long-term follow-up of cognitive dysfunction in patients with aluminum hydroxide-induced macrophagic myofasciitis (MMF). <i>Journal of Inorganic Biochemistry</i> , 2011 , 105, 1457-63	4.2	42
91	Baseline Total Metabolic Tumor Volume Measured with Fixed or Different Adaptive Thresholding Methods Equally Predicts Outcome in Peripheral T Cell Lymphoma. <i>Journal of Nuclear Medicine</i> , 2017 , 58, 276-281	8.9	41
90	Changes in cerebral metabolism are detected prior to perfusion changes in early HIV-CMC: A coregistered (1)H MRS and SPECT study. <i>Journal of Magnetic Resonance Imaging</i> , 2000 , 12, 859-65	5.6	40
89	Interim 18F-fluorodeoxyglucose positron emission tomography in diffuse large B-cell lymphoma: qualitative or quantitative interpretation--where do we stand?. <i>Leukemia and Lymphoma</i> , 2009 , 50, 1753-6	1.9	39
88	Good clinical practice recommendations for the use of PET/CT in oncology. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2020 , 47, 28-50	8.8	37
87	The predictive value of treatment response using FDG PET performed on day 21 of chemoradiotherapy in patients with oesophageal squamous cell carcinoma. A prospective, multicentre study (RTEP3). <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2013 , 40, 1345-55	8.8	34
86	Use of Model-Based Iterative Reconstruction (MBIR) in reduced-dose CT for routine follow-up of patients with malignant lymphoma: dose savings, image quality and phantom study. <i>European Radiology</i> , 2015 , 25, 2362-70	8	32
85	An 18F-FDG-PET maximum standardized uptake value > 10 represents a novel valid marker for discerning Richter's Syndrome. <i>Leukemia and Lymphoma</i> , 2016 , 57, 1474-7	1.9	32
84	Whole-body diffusion magnetic resonance imaging in the assessment of lymphoma. <i>Cancer Imaging</i> , 2012 , 12, 403-8	5.6	31

83	Positron emission tomography/computed tomography with 18F-fluorocholine improve tumor staging and treatment allocation in patients with hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2018 , 69, 336-344	13.4	28
82	Clinical impact of contrast-enhanced computed tomography combined with low-dose (18)F-fluorodeoxyglucose positron emission tomography/computed tomography on routine lymphoma patient management. <i>Leukemia and Lymphoma</i> , 2014 , 55, 2887-92	1.9	25
81	Distribution of ventilation/perfusion ratios in pulmonary embolism: an adjunct to the interpretation of ventilation/perfusion lung scans. <i>Journal of Nuclear Medicine</i> , 2002 , 43, 1596-602	8.9	24
80	18F-FDG PET/CT predicts microvascular invasion and early recurrence after liver resection for hepatocellular carcinoma: A prospective observational study. <i>Hpb</i> , 2019 , 21, 739-747	3.8	23
79	Fully automatic segmentation of diffuse large B cell lymphoma lesions on 3D FDG-PET/CT for total metabolic tumour volume prediction using a convolutional neural network. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2021 , 48, 1362-1370	8.8	23
78	Early interim PET scans in diffuse large B-cell lymphoma: can there be consensus about standardized reporting, and can PET scans guide therapy choices?. <i>Current Hematologic Malignancy Reports</i> , 2012 , 7, 193-9	4.4	22
77	MRI and PET in monitoring response in lymphoma. <i>Cancer Imaging</i> , 2005 , 5 Spec No A, S106-12	5.6	22
76	Early-phase myocardial uptake intensity of Tc-HMDP vs Tc-DPD in patients with hereditary transthyretin-related cardiac amyloidosis. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 217-222	2.1	19
75	Functional imaging of hepatocellular carcinoma using diffusion-weighted MRI and (18)F-FDG PET/CT in patients on waiting-list for liver transplantation. <i>Cancer Imaging</i> , 2016 , 16, 4	5.6	19
74	Dopamine transporter imaging under high-dose transdermal nicotine therapy in Parkinson's disease: an observational study. <i>Nuclear Medicine Communications</i> , 2009 , 30, 513-8	1.6	19
73	Respective prognostic values of germinal center phenotype and early (18)fluorodeoxyglucose-positron emission tomography scanning in previously untreated patients with diffuse large B-cell lymphoma. <i>Haematologica</i> , 2007 , 92, 778-83	6.6	19
72	Assessment of ejection fraction with Tl-201 gated SPECT in myocardial infarction: Precision in a rest-redistribution study and accuracy versus planar angiography. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 31-9	2.1	19
71	Metabolic Tumour Burden Measured by 18F-FDG PET/CT Predicts Malignant Transformation in Patients with Neurofibromatosis Type-1. <i>PLoS ONE</i> , 2016 , 11, e0151809	3.7	19
70	Early Phase Tc-HMDP Scintigraphy for the Diagnosis and Typing of Cardiac Amyloidosis. <i>JACC: Cardiovascular Imaging</i> , 2017 , 10, 601-603	8.4	18
69	Neuropsychological Correlates of Brain Perfusion SPECT in Patients with Macrophagic Myofasciitis. <i>PLoS ONE</i> , 2015 , 10, e0128353	3.7	18
68	Whole body MRI and PET/CT in haematological malignancies. <i>Cancer Imaging</i> , 2007 , 7 Spec No A, S88-93	5.6	17
67	F-sodium fluoride PET/MRI myocardial imaging in patients with suspected cardiac amyloidosis. <i>Journal of Nuclear Cardiology</i> , 2021 , 28, 1586-1595	2.1	16
66	Apical sparing pattern of left ventricular myocardial Tc-HMDP uptake in patients with transthyretin cardiac amyloidosis. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 2072-2079	2.1	16

65	Obinutuzumab vs rituximab for advanced DLBCL: a PET-guided and randomized phase 3 study by LYSA. <i>Blood</i> , 2021 , 137, 2307-2320	2.2	16
64	Patients with plasma cell disorders examined at whole-body dynamic contrast-enhanced MR imaging: initial experience. <i>Radiology</i> , 2009 , 250, 905-15	20.5	15
63	[F-18]-Fluoro-2-deoxy-D: -glucose positron emission tomography as a tool for early detection of immunotherapy response in a murine B cell lymphoma model. <i>Cancer Immunology, Immunotherapy</i> , 2007 , 56, 1163-71	7.4	13
62	Brain F-FDG PET Metabolic Abnormalities in Patients with Long-Lasting Macrophagic Myofasciitis. <i>Journal of Nuclear Medicine</i> , 2017 , 58, 492-498	8.9	12
61	Prognostic value of anthropometric measures extracted from whole-body CT using deep learning in patients with non-small-cell lung cancer. <i>European Radiology</i> , 2020 , 30, 3528-3537	8	12
60	Breast radiotherapy (RT) using tangential fields (TgF): a prospective evaluation of the dose distribution in the sentinel lymph node (SLN) area as determined intraoperatively by clip placement. <i>Annals of Surgical Oncology</i> , 2014 , 21, 3758-65	3.1	12
59	Quantitative lung perfusion scan as a predictor of aerosol distribution heterogeneity and disease severity in children with cystic fibrosis. <i>Nuclear Medicine Communications</i> , 2004 , 25, 563-9	1.6	11
58	Quantitative CT analysis for assessing response in lymphoma (Cheson's criteria). <i>Cancer Imaging</i> , 2005 , 5 Spec No A, S102-6	5.6	11
57	Cognitive dysfunction associated with aluminum hydroxide-induced macrophagic myofasciitis: A reappraisal of neuropsychological profile. <i>Journal of Inorganic Biochemistry</i> , 2018 , 181, 132-138	4.2	10
56	Natural history and impact of treatment with tafamidis on major cardiovascular outcome-free survival time in a cohort of patients with transthyretin amyloidosis. <i>European Journal of Heart Failure</i> , 2021 , 23, 264-274	12.3	10
55	Assessment of myocardial reperfusion after myocardial infarction using automatic 3-dimensional quantification and template matching. <i>Journal of Nuclear Medicine</i> , 2004 , 45, 1981-8	8.9	10
54	Clinical routine use of dopamine transporter imaging in 516 consecutive patients. <i>Journal of Neurology</i> , 2015 , 262, 909-15	5.5	8
53	Can the interim fluorodeoxyglucose-positron emission tomography standardized uptake value be used to determine the need for residual mass biopsy after dose-dense immunochemotherapy for advanced diffuse large B-cell lymphoma?. <i>Journal of Clinical Oncology</i> , 2010 , 28, e719-20; author reply e721-2	2.2	8
52	Diagnostic value of 99mTc-HMDP bone scan in atypical osseous tuberculosis mimicking multiple secondary metastases. <i>Spine</i> , 2004 , 29, E85-7	3.3	7
51	Whole-Body Functional MRI and PET/MRI in Multiple Myeloma. <i>Cancers</i> , 2020 , 12,	6.6	7
50	Extracardiac soft tissue uptake, evidenced on early Tc-HMDP SPECT/CT, helps typing cardiac amyloidosis and demonstrates high prognostic value. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2020 , 47, 2396-2406	8.8	6
49	FDG-PET/CT Brain Findings in a Patient With Macrophagic Myofasciitis. <i>Nuclear Medicine and Molecular Imaging</i> , 2016 , 50, 80-4	1.9	6
48	Subacute dopa-responsive Parkinsonism after successful surgical treatment of aqueductal stenosis. <i>Movement Disorders</i> , 2009 , 24, 2438-40	7	6

47	Early 18fluorodeoxyglucose PET Scan as a Prognostic Tool in Diffuse Large B-Cell Lymphoma Patients Treated with An Anthracycline-Based Chemotherapy Plus Rituximab.. <i>Blood</i> , 2009 , 114, 98-98	2.2	6
46	Cerebral 18F-FDG PET in macrophagic myofasciitis: An individual SVM-based approach. <i>PLoS ONE</i> , 2017 , 12, e0181152	3.7	6
45	Macrophagic myofasciitis-associated dysfunctioning: An update of neuropsychological and neuroimaging features. <i>Best Practice and Research in Clinical Rheumatology</i> , 2018 , 32, 640-650	5.3	6
44	Brain F-FDG PET Metabolic Abnormalities in Macrophagic Myofasciitis: Are They Stable?. <i>Journal of Nuclear Medicine</i> , 2017 , 58, 1532-1533	8.9	5
43	Hepatobiliary MR contrast agent uptake as a predictive biomarker of aggressive features on pathology and reduced recurrence-free survival in resectable hepatocellular carcinoma: comparison with dual-tracer 18F-FDG and 18F-FCH PET/CT. <i>European Radiology</i> , 2020 , 30, 5348-5357	8	5
42	Predictive value of brain 18F-FDG PET/CT in macrophagic myofasciitis?: A case report. <i>Medicine (United States)</i> , 2017 , 96, e8134	1.8	4
41	(99m)Tc-HMDP scintigraphy rectifies wrong diagnosis of AL amyloidosis. <i>Journal of Nuclear Cardiology</i> , 2015 , 22, 853-7	2.1	4
40	Baseline Metabolic Tumor Volume Is Predictive of Patient Outcome in Diffuse Large B Cell Lymphoma. <i>Blood</i> , 2012 , 120, 1598-1598	2.2	4
39	Low Suvmax Measured on Baseline FDG-PET/CT and Elevated κ Microglobulin Are Negative Predictors of Outcome in High Tumor Burden Follicular Lymphoma Treated By Immunochemotherapy: A Pooled Analysis of Three Prospective Studies. <i>Blood</i> , 2016 , 128, 1101-1101	2.2	4
38	Clinical impact of dual-tracer FDOPA and FDG PET/CT for the evaluation of patients with parkinsonian syndromes. <i>Medicine (United States)</i> , 2020 , 99, e23060	1.8	4
37	Characterization of 18-Fluorodeoxyglucose Uptake Pattern in Infective Endocarditis After Transcatheter Aortic Valve Implantation. <i>JACC: Cardiovascular Imaging</i> , 2019 , 12, 930-932	8.4	4
36	Renal Infarction and Its Consequences for Renal Function in Patients With Cardiac Amyloidosis. <i>Mayo Clinic Proceedings</i> , 2019 , 94, 961-975	6.4	3
35	Postinfectious encephalitis: a coregistered SPECT and magnetic resonance imaging study. <i>Clinical Nuclear Medicine</i> , 2002 , 27, 129-30	1.7	3
34	A rare presentation of skull-base osteomyelitis with neurovascular sheath extension following external otitis resolved by PET/MRI. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2018 , 45, 2025	8.8	3
33	Dropped head syndrome with proximal myopathy revealing AL amyloidosis. <i>Joint Bone Spine</i> , 2018 , 85, 779-781	2.9	2
32	Whole-Body Diffusion-weighted MR Imaging of Iron Deposits in Hodgkin, Follicular, and Diffuse Large B-Cell Lymphoma. <i>Radiology</i> , 2018 , 286, 560-567	20.5	2
31	Brain (18)F-FDG, (18)F-Florbetaben PET/CT, (123)I-FP-CIT SPECT and Cardiac (123)I-MIBG Imaging for Diagnosis of a "Cerebral Type" of Lewy Body Disease. <i>Nuclear Medicine and Molecular Imaging</i> , 2016 , 50, 258-60	1.9	2
30	Scintimammography for the Diagnosis of Breast Cancer. <i>Journal of Women's Imaging</i> , 2002 , 4, 66-72		2

29	Early Response Evaluation with 18FDG-PET Scanning, but Not Phenotypic Profile, Are Predictive of Outcome in Diffuse Large B-Cell Lymphoma.. <i>Blood</i> , 2005 , 106, 1914-1914	2.2	2
28	Assessing Cardiac Amyloidosis Subtypes by Unsupervised Phenotype Clustering Analysis. <i>Journal of the American College of Cardiology</i> , 2021 , 78, 2177-2192	15.1	2
27	History of extracardiac/cardiac events in cardiac amyloidosis: prevalence and time from initial onset to diagnosis. <i>ESC Heart Failure</i> , 2021 ,	3.7	2
26	Rituximab plus gemcitabine and oxaliplatin (R-GemOx) in refractory/relapsed diffuse large B-cell lymphoma: a real-life study in patients ineligible for autologous stem-cell transplantation. <i>Leukemia and Lymphoma</i> , 2021 , 62, 2161-2168	1.9	2
25	Health-related quality of life in locally advanced hepatocellular carcinoma treated by either radioembolisation or sorafenib (SARAH trial). <i>European Journal of Cancer</i> , 2021 , 154, 46-56	7.5	2
24	FDG PET/CT of Cervical Gout With Spinal Cord Compression. <i>Clinical Nuclear Medicine</i> , 2020 , 45, e29-e31	1.7	1
23	[18F]fluorodeoxyglucose triple-head coincidence imaging as an adjunct to 131I scanning for follow-up of papillary thyroid carcinoma. <i>Endocrine Practice</i> , 2003 , 9, 273-9	3.2	1
22	Prognostic Value of Baseline Quantitative PET Metrics for Patients with Unfavourable Early Stage Hodgkin Lymphoma Enrolled in the Standard Arm of the EORTC/Lysa/FIL H10 Trial. <i>Blood</i> , 2016 , 128, 184-184	2.2	1
21	Optimization of whole-body 2-[F]FDG-PET/MRI imaging protocol for the initial staging of patients with myeloma. <i>European Radiology</i> , 2021 , 1	8	1
20	Effectiveness and Safety of Subcutaneous Rituximab for Patients With Gastric MALT Lymphoma: A Case-Control Comparison With Intravenous Rituximab. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2021 , 21, e32-e38	2	1
19	Impact of the F-FDG-PET/MRI on Metastatic Staging in Patients with Hepatocellular Carcinoma: Initial Results from 104 Patients. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1
18	Cost-Utility Analysis of Transarterial Radioembolization With Yttrium-90 Resin Microspheres Compared With Sorafenib in Locally Advanced and Inoperable Hepatocellular Carcinoma. <i>Clinical Therapeutics</i> , 2021 , 43, 1201-1212	3.5	0
17	18F-FDG Whole-body PET/MRI of a 30-Week Pregnant Woman With Breast Cancer Revealed Interesting Fetal Findings. <i>Clinical Nuclear Medicine</i> , 2019 , 44, 818-820	1.7	0
16	Recommandations et rffentiels. <i>Medecine Nucleaire</i> , 2019 , 43, 1-4	0.1	
15	Amylose AL rvlè par un syndrome de la tte tombante avec myopathie proximale. <i>Revue Du Rhumatisme (Edition Francaise)</i> , 2019 , 86, 529-531	0.1	
14	Functional Imaging in Lymphoma 2014 , 1311-1334		
13	THE UNVEILED HEART teaching program in cardiovascular nuclear medicine. <i>Nuclear Instruments and Methods in Physics Research, Section A: Accelerators, Spectrometers, Detectors and Associated Equipment</i> , 2004 , 527, 130-133	1.2	
12	Prolonged Remissions After Nivolumab Plus Gemcitabine/Oxaliplatin in Relapsed/Refractory T-cell Lymphoma.. <i>HemaSphere</i> , 2022 , 6, e672	0.3	

11	A Molecular Classifier Increased the Accuracy of Lymphoma Diagnosis By Expert Pathologists in the Diffuse Large B-Cell Lymphoma Gained Trial from the Lysa. <i>Blood</i> , 2021 , 138, 3495-3495	2.2
10	Breast Implant Associated-Anaplastic Large-Cell Lymphoma (BIA-ALCL): Data Based on the Lymphoma Study Association (LYSA) Registry. Promising Results of Brentuximab Vedotin Combined with Cyclophosphamide, Doxorubicin and Prednisone (BV-CHP) As First Line Treatment for Patient Requiring Chemotherapy. <i>Blood</i> , 2021 , 138, 1387-1387	2.2
9	Integrative Study of a High-Grade B-Cell Lymphoma Cohort of 45 Patients: A Single Institution Real Life Study. <i>Blood</i> , 2021 , 138, 4576-4576	2.2
8	Shunt imaging in a complex congenital cardiomyopathy. <i>Clinical Nuclear Medicine</i> , 2000 , 25, 827-8	1.7
7	Response Assessment after an Inductive CHOP or CHOP-Like Regimen with or without Rituximab in 103 Patients with Diffuse Large B-Cell Lymphoma (DLBCL): Integrating 18Fluorodeoxyglucose Positron Emission Tomography (PET) to the International Workshop Criteria (IWC).. <i>Blood</i> , 2006 , 108, 2735-2735	2.2
6	Breast Implant Associated-Anaplastic Large Cell Lymphoma (BIA-ALCL): The French Lymphoma Study Association (LYSA) Registry Data. <i>Blood</i> , 2018 , 132, 1641-1641	2.2
5	Breast Implant Associated-Anaplastic Large Cell Lymphoma (BIA-ALCL): The Lymphoma Study Association (LYSA) Registry Data. <i>Blood</i> , 2019 , 134, 4021-4021	2.2
4	Rituximab Plus Gemcitabine and Oxaliplatin (R-GemOx) in Refractory/Relapsed (R/R) DLBCL. a Real Life Study in Patients Ineligible for Autologous Transplantation. <i>Blood</i> , 2019 , 134, 4115-4115	2.2
3	Number of Circulating t(14;18) Tumor Cells at Diagnosis Is Related to, but Add to the Prognostic Value of Metabolic Tumor Burden in Follicular Lymphoma. <i>Blood</i> , 2015 , 126, 3872-3872	2.2
2	Prognostic Value of Early 18fluorodeoxyglucose PET Scan in Patients with Diffuse Large B Cell Lymphoma Treated with Rituximab Plus CHOP or High-Dose CHOP.. <i>Blood</i> , 2008 , 112, 2006-2006	2.2
1	Meningeal Metastasis Relapse With Focal Involvement of Cranial Bone Flap: A Case Resolved by 18F-DOPA PET/MRI. <i>Clinical Nuclear Medicine</i> , 2019 , 44, e315-e317	1.7