

# Jeong Woo Kim

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7937568/publications.pdf>

Version: 2024-02-01

15  
papers

153  
citations

1478505

6  
h-index

1199594

12  
g-index

15  
all docs

15  
docs citations

15  
times ranked

358  
citing authors

#	ARTICLE	IF	CITATIONS
1	Ultrasonographic index for the diagnosis of non-alcoholic steatohepatitis in patients with non-alcoholic fatty liver disease. <i>Quantitative Imaging in Medicine and Surgery</i> , 2022, 12, 1815-1829.	2.0	3
2	MRE-based NASH score for diagnosis of nonalcoholic steatohepatitis in patients with nonalcoholic fatty liver disease. <i>Hepatology International</i> , 2022, 16, 316-324.	4.2	5
3	Ectopic lesions in the abdomen and pelvis: a multimodality pictorial review. <i>Abdominal Radiology</i> , 2022, 47, 2254-2276.	2.1	1
4	Recent Updates of Abbreviated MRI for Hepatocellular Carcinoma Screening. <i>Journal of the Korean Society of Radiology</i> , 2021, 82, 280.	0.2	0
5	Multiparametric MR Index for the Diagnosis of Non-Alcoholic Steatohepatitis in Patients with Non-Alcoholic Fatty Liver Disease. <i>Scientific Reports</i> , 2020, 10, 2671.	3.3	28
6	Multiparametric MR Is a Valuable Modality for Evaluating Disease Severity of Nonalcoholic Fatty Liver Disease. <i>Clinical and Translational Gastroenterology</i> , 2020, 11, e00157.	2.5	16
7	Abbreviated Gadoxetic Acid-enhanced MRI with Second-Shot Arterial Phase Imaging for Liver Metastasis Evaluation. <i>Radiology Imaging Cancer</i> , 2019, 1, e190006.	1.6	4
8	Second shot arterial phase to overcome degraded hepatic arterial phase in liver MR imaging. <i>European Radiology</i> , 2019, 29, 2821-2829.	4.5	5
9	The value of paradoxical uptake of hepatocellular carcinoma on the hepatobiliary phase of gadoxetic acid-enhanced liver magnetic resonance imaging for the prediction of lipiodol uptake after transcatheter arterial chemoembolization. <i>European Journal of Radiology</i> , 2017, 89, 169-176.	2.6	3
10	Use of hepatobiliary phase images in Gd-EOB-DTPA-enhanced MRI of breast cancer hepatic metastasis to predict response to chemotherapy. <i>Clinical Imaging</i> , 2017, 43, 127-131.	1.5	6
11	Washout appearance in Gd-EOB-DTPA-enhanced MR imaging: A differentiating feature between hepatocellular carcinoma with paradoxical uptake on the hepatobiliary phase and focal nodular hyperplasia-like nodules. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 45, 1599-1608.	3.4	18
12	Differentiation of hepatocellular carcinoma from its various mimickers in liver magnetic resonance imaging: What are the tips when using hepatocyte-specific agents?. <i>World Journal of Gastroenterology</i> , 2016, 22, 284.	3.3	28
13	The Short Breath-Hold Technique, Controlled Aliasing in Parallel Imaging Results in Higher Acceleration, Can Be the First Step to Overcoming a Degraded Hepatic Arterial Phase in Liver Magnetic Resonance Imaging. <i>Investigative Radiology</i> , 2016, 51, 440-446.	6.2	26
14	A Patient With Left Upper Abdominal Pain Who Had Petit Sac. <i>Gastroenterology</i> , 2016, 150, e12-e13.	1.3	0
15	Intravoxel incoherent motion magnetic resonance imaging to predict vesicoureteral reflux in children with urinary tract infection. <i>European Radiology</i> , 2016, 26, 1670-1677.	4.5	10