

Santiago Gutiérrez Martínez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/793990/publications.pdf>

Version: 2024-02-01

124
papers

4,969
citations

66234

42
h-index

110170

64
g-index

125
all docs

125
docs citations

125
times ranked

3220
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | The penicillin gene cluster is amplified in tandem repeats linked by conserved hexanucleotide sequences.. Proceedings of the National Academy of Sciences of the United States of America, 1995, 92, 6200-6204. | 3.3 | 174 |
| 2 | The ThPG1 Endopolygalacturonase Is Required for the <i>Trichoderma harzianum</i>–Plant Beneficial Interaction. Molecular Plant-Microbe Interactions, 2009, 22, 1021-1031. | 1.4 | 173 |
| 3 | The contribution of <i>Trichoderma</i> to balancing the costs of plant growth and defense. International Microbiology, 2013, 16, 69-80. | 1.1 | 173 |
| 4 | Transgenic expression of the <i>Trichoderma harzianum</i> hsp70 gene increases <i>Arabidopsis</i> resistance to heat and other abiotic stresses. Journal of Plant Physiology, 2010, 167, 659-665. | 1.6 | 161 |
| 5 | Involvement of <i>Trichoderma</i> Trichothecenes in the Biocontrol Activity and Induction of Plant Defense-Related Genes. Applied and Environmental Microbiology, 2012, 78, 4856-4868. | 1.4 | 143 |
| 6 | Evolution of structural diversity of trichothecenes, a family of toxins produced by plant pathogenic and entomopathogenic fungi. PLoS Pathogens, 2018, 14, e1006946. | 2.1 | 141 |
| 7 | Identification of Loci and Functional Characterization of Trichothecene Biosynthesis Genes in Filamentous Fungi of the Genus <i>Trichoderma</i> . Applied and Environmental Microbiology, 2011, 77, 4867-4877. | 1.4 | 124 |
| 8 | Influence of <i>Rhizoctonia solani</i> and <i>Trichoderma</i> spp. in growth of bean (<i>Phaseolus vulgaris</i> L.) and in the induction of plant defense-related genes. Frontiers in Plant Science, 2015, 6, 685. | 1.7 | 116 |
| 9 | Age-Related Clinical, Serological, and Histopathological Features of Celiac Disease. American Journal of Gastroenterology, 2008, 103, 2360-2365. | 0.2 | 114 |
| 10 | Overexpression of the trichodiene synthase gene <i>tri5</i> increases trichodermin production and antimicrobial activity in <i>Trichoderma brevicompactum</i> . Fungal Genetics and Biology, 2011, 48, 285-296. | 0.9 | 110 |
| 11 | Phylogenomic Analysis of a 55.1-kb 19-Gene Dataset Resolves a Monophyletic <i>Fusarium</i> that Includes the <i>Fusarium solani</i> Species Complex. Phytopathology, 2021, 111, 1064-1079. | 1.1 | 107 |
| 12 | The <i>cefG</i> gene of <i>Cephalosporium acremonium</i> is linked to the <i>cefEF</i> gene and encodes a deacetylcephalosporin C acetyltransferase closely related to homoserine O-acetyltransferase. Journal of Bacteriology, 1992, 174, 3056-3064. | 1.0 | 92 |
| 13 | Functional Analysis of the <i>Trichoderma harzianum nox1</i> Gene, Encoding an NADPH Oxidase, Relates Production of Reactive Oxygen Species to Specific Biocontrol Activity against <i>Pythium ultimum</i> . Applied and Environmental Microbiology, 2011, 77, 3009-3016. | 1.4 | 92 |
| 14 | Mutants blocked in penicillin biosynthesis show a deletion of the entire penicillin gene cluster at a specific site within a conserved hexanucleotide sequence. Applied Microbiology and Biotechnology, 1996, 44, 597-604. | 1.7 | 90 |
| 15 | Relevance of trichothecenes in fungal physiology: Disruption of <i>tri5</i> in <i>Trichoderma arundinaceum</i> . Fungal Genetics and Biology, 2013, 53, 22-33. | 0.9 | 89 |
| 16 | Resolution of four large chromosomes in penicillin-producing filamentous fungi: the penicillin gene cluster is located on chromosome II (9.6 Mb) in <i>Penicillium notatum</i> and chromosome 1 (10.4 Mb) in <i>Penicillium chrysogenum</i> . Molecular Genetics and Genomics, 1993, 241-241, 573-578. | 2.4 | 80 |
| 17 | The isopenicillin-N acyltransferase of <i>Penicillium chrysogenum</i> has isopenicillin-N amidohydrolase, 6-aminopenicillanic acid acyltransferase and penicillin amidase activities, all of which are encoded by the single <i>penDE</i> gene. FEBS Journal, 1993, 215, 323-332. | 0.2 | 80 |
| 18 | Cloning and characterization of the <i>erg1</i> gene of <i>Trichoderma harzianum</i> : Effect of the <i>erg1</i> silencing on ergosterol biosynthesis and resistance to terbinafine. Fungal Genetics and Biology, 2006, 43, 164-178. | 0.9 | 77 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | The Cerato-Platanin protein Epl-1 from <i>Trichoderma harzianum</i> is involved in mycoparasitism, plant resistance induction and self cell wall protection. <i>Scientific Reports</i> , 2016, 5, 17998. | 1.6 | 77 |
| 20 | Expression of the <i>cefC</i> gene is limiting for cephalosporin biosynthesis in <i>Acremonium chrysogenum</i> . <i>Applied Microbiology and Biotechnology</i> , 1997, 48, 606-614. | 1.7 | 74 |
| 21 | The <i>cefT</i> gene of <i>Acremonium chrysogenum</i> C10 encodes a putative multidrug efflux pump protein that significantly increases cephalosporin C production. <i>Molecular Genetics and Genomics</i> , 2002, 267, 673-683. | 1.0 | 73 |
| 22 | A Novel Epimerization System in Fungal Secondary Metabolism Involved in the Conversion of Isopenicillin N into Penicillin N in <i>Acremonium chrysogenum</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 46216-46225. | 1.6 | 71 |
| 23 | Overexpression of a <i>Trichoderma</i> HSP70 gene increases fungal resistance to heat and other abiotic stresses. <i>Fungal Genetics and Biology</i> , 2008, 45, 1506-1513. | 0.9 | 68 |
| 24 | Autonomously replicating plasmids carrying the <i>AMA1</i> region in <i>Penicillium chrysogenum</i> . <i>Current Genetics</i> , 1996, 29, 482-489. | 0.8 | 65 |
| 25 | Production of trichodiene by <i>Trichoderma harzianum</i> alters the perception of this biocontrol strain by plants and antagonized fungi. <i>Environmental Microbiology</i> , 2015, 17, 2628-2646. | 1.8 | 64 |
| 26 | Expression of the <i>penDE</i> gene of <i>Penicillium chrysogenum</i> encoding isopenicillin N acyltransferase in <i>Cephalosporium acremonium</i> : production of benzylpenicillin by the transformants. <i>Molecular Genetics and Genomics</i> , 1991, 225, 56-64. | 2.4 | 63 |
| 27 | Penicillin and cephalosporin biosynthesis: mechanism of carbon catabolite regulation of penicillin production. <i>Antonie Van Leeuwenhoek</i> , 1999, 75, 21-31. | 0.7 | 63 |
| 28 | Cloning, characterization of the acyl-CoA : 6-amino penicillanic acid acyltransferase gene of <i>Aspergillus nidulans</i> and linkage to the isopenicillin N synthase gene. <i>Molecular Genetics and Genomics</i> , 1990, 221, 322-330. | 2.4 | 62 |
| 29 | Gene organization and plasticity of the beta-lactam genes in different filamentous fungi. <i>Antonie Van Leeuwenhoek</i> , 1999, 75, 81-94. | 0.7 | 61 |
| 30 | Generation, annotation and analysis of ESTs from <i>Trichoderma harzianum</i> CECT 2413. <i>BMC Genomics</i> , 2006, 7, 193. | 1.2 | 60 |
| 31 | Partial silencing of a hydroxy-methylglutaryl-CoA reductase-encoding gene in <i>Trichoderma harzianum</i> CECT 2413 results in a lower level of resistance to lovastatin and lower antifungal activity. <i>Fungal Genetics and Biology</i> , 2007, 44, 269-283. | 0.9 | 60 |
| 32 | Secondary Metabolism and Antimicrobial Metabolites of <i>Trichoderma</i> . , 2014, , 125-137. | | 59 |
| 33 | Novel aspinolide production by <i>Trichoderma arundinaceum</i> with a potential role in <i>Botrytis cinerea</i> antagonistic activity and plant defence priming. <i>Environmental Microbiology</i> , 2015, 17, 1103-1118. | 1.8 | 56 |
| 34 | Exogenous methionine increases levels of mRNAs transcribed from <i>pcbAB</i> , <i>pcbC</i> , and <i>cefEF</i> genes, encoding enzymes of the cephalosporin biosynthetic pathway, in <i>Acremonium chrysogenum</i> . <i>Journal of Bacteriology</i> , 1994, 176, 985-991. | 1.0 | 53 |
| 35 | Thaumatococin Production in <i>Aspergillus awamori</i> by Use of Expression Cassettes with Strong Fungal Promoters and High Gene Dosage. <i>Applied and Environmental Microbiology</i> , 1999, 65, 1168-1174. | 1.4 | 53 |
| 36 | Silencing of the Aspergillopepsin B (<i>pepB</i>) Gene of <i>Aspergillus awamori</i> by Antisense RNA Expression or Protease Removal by Gene Disruption Results in a Large Increase in Thaumatococin Production. <i>Applied and Environmental Microbiology</i> , 2002, 68, 3550-3559. | 1.4 | 49 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Resolution of chromosomes III and VI of <i>Aspergillus nidulans</i> by pulsed-field gel electrophoresis shows that the penicillin biosynthetic pathway genes <i>pcbAB</i> , <i>pcbC</i> , and <i>penDE</i> are clustered on chromosome VI (3.0 megabases). <i>Journal of Bacteriology</i> , 1992, 174, 7063-7067. | 1.0 | 47 |
| 38 | Screening of antimicrobial activities in <i>Trichoderma</i> isolates representing three <i>Trichoderma</i> sections. <i>Mycological Research</i> , 2005, 109, 1397-1406. | 2.5 | 47 |
| 39 | Development of a qPCR Strategy to Select Bean Genes Involved in Plant Defense Response and Regulated by the <i>Trichoderma velutinum</i> × <i>Rhizoctonia solani</i> Interaction. <i>Frontiers in Plant Science</i> , 2016, 7, 1109. | 1.7 | 46 |
| 40 | Overexpression of the <i>Trichoderma brevicompactum tri5</i> Gene: Effect on the Expression of the Trichodermin Biosynthetic Genes and on Tomato Seedlings. <i>Toxins</i> , 2011, 3, 1220-1232. | 1.5 | 45 |
| 41 | A comparison of the phenotypic and genetic stability of recombinant <i>Trichoderma</i> spp. generated by protoplast- and <i>Agrobacterium</i> -mediated transformation. <i>Journal of Microbiology</i> , 2006, 44, 383-95. | 1.3 | 43 |
| 42 | A Novel Heptameric Sequence (TTAGTAA) Is the Binding Site for a Protein Required for High Level Expression of <i>pcbAB</i> , the First Gene of the Penicillin Biosynthesis in <i>Penicillium chrysogenum</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 2423-2430. | 1.6 | 42 |
| 43 | Stable transformants of the azaphilone pigment-producing <i>Monascus purpureus</i> obtained by protoplast transformation and <i>Agrobacterium</i> -mediated DNA transfer. <i>Current Genetics</i> , 2003, 43, 447-452. | 0.8 | 42 |
| 44 | Transcription of the <i>pcbAB</i> , <i>pcbC</i> and <i>penDE</i> genes of <i>Penicillium chrysogenum</i> AS-P-78 is repressed by glucose and the repression is not reversed by alkaline pHs. <i>Microbiology (United Kingdom)</i> , 1999, 145, 317-324. | 0.7 | 41 |
| 45 | Detection of putative peptide synthetase genes in <i>Trichoderma</i> species: Application of this method to the cloning of a gene from <i>T. harzianum</i> CECT 2413. <i>FEMS Microbiology Letters</i> , 2005, 244, 139-148. | 0.7 | 41 |
| 46 | ThPTR2, a di/tri-peptide transporter gene from <i>Trichoderma harzianum</i> . <i>Fungal Genetics and Biology</i> , 2006, 43, 234-246. | 0.9 | 41 |
| 47 | Involvement of <i>Trichoderma harzianum</i> Epl-1 Protein in the Regulation of Botrytis Virulence- and Tomato Defense-Related Genes. <i>Frontiers in Plant Science</i> , 2017, 8, 880. | 1.7 | 40 |
| 48 | The heterologous overexpression of <i>hsp23</i> , a small heat-shock protein gene from <i>Trichoderma virens</i> , confers thermotolerance to <i>T. harzianum</i> . <i>Current Genetics</i> , 2007, 52, 45-53. | 0.8 | 39 |
| 49 | Characterization of the <i>lys2</i> gene of <i>Penicillium chrysogenum</i> encoding $\hat{\Gamma}$ -amino adipic acid reductase. <i>Molecular Genetics and Genomics</i> , 1998, 259, 549-556. | 2.4 | 38 |
| 50 | Targeted Inactivation of the <i>mecB</i> Gene, Encoding Cystathionine- $\hat{\Gamma}$ -Lyase, Shows that the Reverse Transsulfuration Pathway Is Required for High-Level Cephalosporin Biosynthesis in <i>Acremonium chrysogenum</i> C10 but Not for Methionine Induction of the Cephalosporin Genes. <i>Journal of Bacteriology</i> , 2001, 183, 1765-1772. | 1.0 | 38 |
| 51 | Trichodiene Production in a <i>Trichoderma harzianum</i> <i>erg1</i> -Silenced Strain Provides Evidence of the Importance of the Sterol Biosynthetic Pathway in Inducing Plant Defense-Related Gene Expression. <i>Molecular Plant-Microbe Interactions</i> , 2015, 28, 1181-1197. | 1.4 | 38 |
| 52 | Genes for β -lactam antibiotic biosynthesis. <i>Antonie Van Leeuwenhoek</i> , 1995, 67, 181-200. | 0.7 | 37 |
| 53 | Characterization and nitrogen-source regulation at the transcriptional level of the <i>gdh A</i> gene of <i>Aspergillus awamori</i> encoding an NADP-dependent glutamate dehydrogenase. <i>Current Genetics</i> , 1998, 34, 50-59. | 0.8 | 37 |
| 54 | Effects of Trichothecene Production on the Plant Defense Response and Fungal Physiology: Overexpression of the <i>Trichoderma arundinaceum tri4</i> Gene in <i>T. harzianum</i> . <i>Applied and Environmental Microbiology</i> , 2015, 81, 6355-6366. | 1.4 | 37 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Effect of <i>Trichoderma velutinum</i> and <i>Rhizoctonia solani</i> on the Metabolome of Bean Plants (<i>Phaseolus vulgaris</i> L.). <i>International Journal of Molecular Sciences</i> , 2019, 20, 549. | 1.8 | 36 |
| 56 | Nitrogen Metabolism and Growth Enhancement in Tomato Plants Challenged with <i>Trichoderma harzianum</i> Expressing the <i>Aspergillus nidulans</i> Acetamidase <i>amdS</i> Gene. <i>Frontiers in Microbiology</i> , 2016, 7, 1182. | 1.5 | 34 |
| 57 | Intrachromosomal recombination between direct repeats in <i>Penicillium chrysogenum</i> : gene conversion and deletion events. <i>Molecular Genetics and Genomics</i> , 1999, 261, 994-1000. | 2.4 | 33 |
| 58 | Overexpression and lack of degradation of thaumatin in an aspergillopepsin A-defective mutant of <i>Aspergillus awamori</i> containing an insertion in the <i>pepA</i> gene. <i>Applied Microbiology and Biotechnology</i> , 2000, 54, 772-777. | 1.7 | 33 |
| 59 | Inhibitory activity of <i>Beauveria bassiana</i> and <i>Trichoderma</i> spp. on the insect pests <i>Xylotrechus arvicola</i> (Coleoptera: Cerambycidae) and <i>Acanthoscelides obtectus</i> (Coleoptera: Chrisomelidae). <i>Trends in Microbiology</i> , 2014, 22, 104-114. | 1.4 | 31 |
| 60 | Expression of a synthetic copy of the bovine chymosin gene in <i>Aspergillus awamori</i> from constitutive and pH-regulated promoters and secretion using two different pre-pro sequences. <i>Biotechnology and Bioengineering</i> , 2003, 83, 249-259. | 1.7 | 29 |
| 61 | Cloning and characterization of the <i>Thcut1</i> gene encoding a cutinase of <i>Trichoderma harzianum</i> T34. <i>Current Genetics</i> , 2008, 54, 301-312. | 0.8 | 29 |
| 62 | Colonization of <i>Vitis vinifera</i> L. by the Endophyte <i>Trichoderma</i> sp. Strain T154: Biocontrol Activity Against <i>Phaeoacremonium minimum</i> . <i>Frontiers in Plant Science</i> , 2020, 11, 1170. | 1.7 | 29 |
| 63 | Molecular characterization of three loss-of-function mutations in the isopenicillin N-acyltransferase gene (<i>penDE</i>) of <i>Penicillium chrysogenum</i> . <i>Journal of Bacteriology</i> , 1994, 176, 4941-4948. | 1.0 | 28 |
| 64 | The importance of chorismate mutase in the biocontrol potential of <i>Trichoderma parareesei</i> . <i>Frontiers in Microbiology</i> , 2015, 6, 1181. | 1.5 | 28 |
| 65 | Effect of deletion of a trichothecene toxin regulatory gene on the secondary metabolism transcriptome of the saprotrophic fungus <i>Trichoderma arundinaceum</i> . <i>Fungal Genetics and Biology</i> , 2018, 119, 29-46. | 0.9 | 27 |
| 66 | Molecular characterization of the <i>Acremonium chrysogenum</i> <i>cefG</i> gene product: the native deacetylcephalosporin C acetyltransferase is not processed into subunits. <i>Biochemical Journal</i> , 1999, 337, 379-385. | 1.7 | 26 |
| 67 | Trichothecenes and aspinolides produced by <i>Trichoderma arundinaceum</i> regulate expression of <i>Botrytis cinerea</i> genes involved in virulence and growth. <i>Environmental Microbiology</i> , 2016, 18, 3991-4004. | 1.8 | 25 |
| 68 | Characterization and lysine control of expression of the <i>lys1</i> gene of <i>Penicillium chrysogenum</i> encoding homocitrate synthase. <i>Gene</i> , 1999, 226, 51-59. | 1.0 | 23 |
| 69 | Effect of trichodiene production by <i>Trichoderma harzianum</i> on <i>Acanthoscelides obtectus</i> . <i>Journal of Stored Products Research</i> , 2018, 77, 231-239. | 1.2 | 23 |
| 70 | Overexpression of <i>erg1</i> gene in <i>Trichoderma harzianum</i> CECT 2413: effect on the induction of tomato defence-related genes. <i>Journal of Applied Microbiology</i> , 2014, 117, 812-823. | 1.4 | 22 |
| 71 | Genetic bases for variation in structure and biological activity of trichothecene toxins produced by diverse fungi. <i>Applied Microbiology and Biotechnology</i> , 2020, 104, 5185-5199. | 1.7 | 21 |
| 72 | Involvement of the Transcriptional Coactivator <i>ThMBF1</i> in the Biocontrol Activity of <i>Trichoderma harzianum</i> . <i>Frontiers in Microbiology</i> , 2017, 8, 2273. | 1.5 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Investigations of <i>Trichoderma</i> spp. and <i>Beauveria bassiana</i> as biological control agent for <i>Xylotrechus arvicola</i> , a major insect pest in Spanish vineyards. <i>Journal of Economic Entomology</i> , 2018, 111, 2585-2591. | 0.8 | 20 |
| 74 | Expression of genes and processing of enzymes for the biosynthesis of penicillins and cephalosporins. <i>Antonie Van Leeuwenhoek</i> , 1994, 65, 227-243. | 0.7 | 19 |
| 75 | Characterization of the bip gene of <i>Aspergillus awamori</i> encoding a protein with an HDEL retention signal homologous to the mammalian BiP involved in polypeptide secretion. <i>Current Genetics</i> , 1997, 32, 139-146. | 0.8 | 19 |
| 76 | The isopenicillin N acyltransferases of <i>Aspergillus nidulans</i> and <i>Penicillium chrysogenum</i> differ in their ability to maintain the 40-kDa alphabeta heterodimer in an undissociated form. <i>FEBS Journal</i> , 2003, 270, 1958-1968. | 0.2 | 19 |
| 77 | Overexpression of the lys1 gene in <i>Penicillium chrysogenum</i> : homocitrate synthase levels, β -aminoacidic acid pool and penicillin production. <i>Applied Microbiology and Biotechnology</i> , 2000, 54, 69-77. | 1.7 | 18 |
| 78 | Detection of peptaibols and partial cloning of a putative peptaibol synthetase gene from <i>T. harzianum</i> CECT 2413. <i>Folia Microbiologica</i> , 2006, 51, 114-120. | 1.1 | 18 |
| 79 | TvDim1 of <i>Trichoderma virens</i> is involved in redox-processes and confers resistance to oxidative stresses. <i>Current Genetics</i> , 2010, 56, 63-73. | 0.8 | 18 |
| 80 | Relevance of the deletion of the <i>Tatri4</i> gene in the secondary metabolome of <i>Trichoderma arundinaceum</i> . <i>Organic and Biomolecular Chemistry</i> , 2018, 16, 2955-2965. | 1.5 | 18 |
| 81 | Three genes hrdB, hrdD and hrdT of <i>Streptomyces griseus</i> IMRU 3570, encoding sigma factor-like proteins, are differentially expressed under specific nutritional conditions. <i>Gene</i> , 1995, 153, 41-48. | 1.0 | 17 |
| 82 | Cloning and characterization of the gene cah B encoding a cephalosporin C acetylhydrolase from <i>Acremonium chrysogenum</i> . <i>Applied Microbiology and Biotechnology</i> , 2001, 57, 350-356. | 1.7 | 16 |
| 83 | Role of <i>Trichoderma arundinaceum</i> tri10 in regulation of terpene biosynthetic genes and in control of metabolic flux. <i>Fungal Genetics and Biology</i> , 2019, 122, 31-46. | 0.9 | 16 |
| 84 | Identification of plant genes putatively involved in the perception of fungal ergosterol \rightarrow squalene. <i>Journal of Integrative Plant Biology</i> , 2020, 62, 927-947. | 4.1 | 16 |
| 85 | Genome-wide analysis of differentially expressed genes from <i>Penicillium chrysogenum</i> grown with a repressing or a non-repressing carbon source. <i>Current Genetics</i> , 2006, 49, 85-96. | 0.8 | 15 |
| 86 | Co-transformation with autonomous replicating and integrative plasmids in <i>Penicillium chrysogenum</i> is highly efficient and leads in some cases to rescue of the intact integrative plasmid. <i>Fungal Genetics and Biology</i> , 2003, 40, 83-92. | 0.9 | 14 |
| 87 | Botrydial and botcinins produced by <i>Botrytis cinerea</i> regulate the expression of <i>Trichoderma arundinaceum</i> genes involved in trichothecene biosynthesis. <i>Molecular Plant Pathology</i> , 2016, 17, 1017-1031. | 2.0 | 14 |
| 88 | Subcellular localization of the homocitrate synthase in <i>Penicillium chrysogenum</i> . <i>Molecular Genetics and Genomics</i> , 2002, 266, 711-719. | 1.0 | 13 |
| 89 | A cytochrome P450 monooxygenase gene required for biosynthesis of the trichothecene toxin harzianum A in <i>Trichoderma</i> . <i>Applied Microbiology and Biotechnology</i> , 2019, 103, 8087-8103. | 1.7 | 13 |
| 90 | Requirement of Two Acyltransferases for 4-O-Acylation during Biosynthesis of Harzianum A, an Antifungal Trichothecene Produced by <i>Trichoderma arundinaceum</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2019, 67, 723-734. | 2.4 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Molecular characterization of the <i>Acremonium chrysogenum</i> cefG gene product: the native deacetylcephalosporin C acetyltransferase is not processed into subunits. <i>Biochemical Journal</i> , 1999, 337, 379. | 1.7 | 10 |
| 92 | Characterization of the reverse transsulfuration gene <i>mecB</i> of <i>Acremonium chrysogenum</i> , which encodes a functional cystathionine- β -lyase. <i>Molecular Genetics and Genomics</i> , 2001, 264, 746-754. | 1.0 | 10 |
| 93 | Spores of <i>Trichoderma</i> strains sprayed over <i>Acanthoscelides obtectus</i> and <i>Phaseolus vulgaris</i> L. beans: Effects in the biology of the bean weevil. <i>Journal of Stored Products Research</i> , 2020, 88, 101666. | 1.2 | 10 |
| 94 | Use of the volatile trichodiene to reduce <i>Fusarium</i> head blight and trichothecene contamination in wheat. <i>Microbial Biotechnology</i> , 2022, 15, 513-527. | 2.0 | 10 |
| 95 | Volatile Organic Compound Chamber: A Novel Technology for Microbiological Volatile Interaction Assays. <i>Journal of Fungi (Basel, Switzerland)</i> , 2021, 7, 248. | 1.5 | 10 |
| 96 | The Influence of Temperature on the Growth, Sporulation, Colonization, and Survival of <i>Trichoderma</i> spp. in Grapevine Pruning Wounds. <i>Agronomy</i> , 2021, 11, 1771. | 1.3 | 10 |
| 97 | Distribution, Function, and Evolution of a Gene Essential for Trichothecene Toxin Biosynthesis in <i>Trichoderma</i> . <i>Frontiers in Microbiology</i> , 2021, 12, 791641. | 1.5 | 10 |
| 98 | Effect of trichodiene synthase encoding gene expression in <i>Trichoderma</i> strains on their effectiveness in the control of <i>Acanthoscelides obtectus</i> . <i>Journal of Stored Products Research</i> , 2019, 83, 275-280. | 1.2 | 9 |
| 99 | Synthesis of Trichodermin Derivatives and Their Antimicrobial and Cytotoxic Activities. <i>Molecules</i> , 2019, 24, 3811. | 1.7 | 9 |
| 100 | Mutants blocked in penicillin biosynthesis show a deletion of the entire penicillin gene cluster at a specific site within a conserved hexanucleotide sequence. <i>Applied Microbiology and Biotechnology</i> , 1996, 44, 597-604. | 1.7 | 8 |
| 101 | The specific transport system for lysine is fully inhibited by ammonium in <i>Penicillium chrysogenum</i> : an ammonium-insensitive system allows uptake in carbon-starved cells. <i>Antonie Van Leeuwenhoek</i> , 2000, 77, 91-100. | 0.7 | 6 |
| 102 | Influence of Fungicide Application and Vine Age on <i>Trichoderma</i> Diversity as Source of Biological Control Agents. <i>Agronomy</i> , 2021, 11, 446. | 1.3 | 6 |
| 103 | Influence of Physicochemical Characteristics of Bean Crop Soil in <i>Trichoderma</i> spp. Development. <i>Agronomy</i> , 2021, 11, 274. | 1.3 | 6 |
| 104 | Fungal Horizontal Gene Transfer: A History Beyond the Phylogenetic Kingdoms. , 2019, , 315-336. | | 5 |
| 105 | Self-Inhibitory Activity of <i>Trichoderma</i> Soluble Metabolites and Their Antifungal Effects on <i>Fusarium oxysporum</i> . <i>Journal of Fungi (Basel, Switzerland)</i> , 2020, 6, 176. | 1.5 | 5 |
| 106 | Characterization of the <i>gdhA</i> Gene from the Phytopathogen <i>Botrytis cinerea</i> . <i>Fungal Genetics and Biology</i> , 2001, 34, 193-206. | 0.9 | 4 |
| 107 | <i>Trichoderma</i> Transformation Methods. <i>Fungal Biology</i> , 2015, , 41-48. | 0.3 | 4 |
| 108 | <i>Trichoderma</i> trichothecenes. , 2020, , 281-301. | | 4 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Influence of Substrates in the Development of Bean and in Pathogenicity of <i>Rhizoctonia solani</i> JG KÄ¼hn. <i>Agronomy</i> , 2020, 10, 707. | 1.3 | 4 |
| 110 | Evaluation of substrates and additives to <i>Trichoderma harzianum</i> development by qPCR. <i>Agronomy Journal</i> , 2020, 112, 3188-3194. | 0.9 | 4 |
| 111 | Autonomously replicating plasmids carrying the AMA1 region in <i>Penicillium chrysogenum</i> . <i>Current Genetics</i> , 1996, 29, 482-489. | 0.8 | 4 |
| 112 | Novel culture chamber to evaluate in vitro plant-microbe volatile interactions: Effects of <i>Trichoderma harzianum</i> volatiles on wheat plantlets. <i>Plant Science</i> , 2022, 320, 111286. | 1.7 | 4 |
| 113 | Organic and Conventional Bean Pesticides in Development of Autochthonous <i>Trichoderma</i> Strains. <i>Journal of Fungi</i> (Basel, Switzerland), 2022, 8, 603. | 1.5 | 4 |
| 114 | Volatile-mediated interactions between <i>Trichoderma harzianum</i> and <i>Acanthoscelides obtectus</i> : A novel in vitro methodology to evaluate the impact of microbial volatile compounds on dry grain storage pests. <i>Biological Control</i> , 2022, 169, 104868. | 1.4 | 3 |
| 115 | Molecular genetics as a tool to remove bottlenecks in the biosynthesis of β -lactam antibiotics. <i>World Journal of Microbiology and Biotechnology</i> , 1996, 12, 517-523. | 1.7 | 2 |
| 116 | Intrachromosomal recombination after targeted monocopy integration in <i>Penicillium chrysogenum</i> : stabilization of the direct repeats to prevent loss of the inserted gene. <i>Current Genetics</i> , 2001, 39, 231-236. | 0.8 | 2 |
| 117 | Insecticidal activity of <i>Trichoderma harzianum</i> against <i>Xylotrechus arvicola</i> and <i>Acanthoscelides obtectus</i> immature stages. <i>Planta Medica</i> , 2016, 81, S1-S381. | 0.7 | 2 |
| 118 | Germination and Agronomic Traits of <i>Phaseolus vulgaris</i> L. Beans Sprayed with <i>Trichoderma</i> Strains and Attacked by <i>Acanthoscelides obtectus</i> . <i>Agronomy</i> , 2021, 11, 2130. | 1.3 | 2 |
| 119 | Editorial: Designing Bio-Formulations Based on Organic Amendments, Beneficial Microbes and Their Metabolites. <i>Frontiers in Microbiology</i> , 2021, 12, 832149. | 1.5 | 2 |
| 120 | Effect of Farnesol, a compound produced by <i>Trichoderma</i> when growing on bean (<i>Phaseolus vulgaris</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf | 0.9 | 1 |
| 121 | Use of VOC Chambers to evaluate the impact of microbial volatile compounds on dry grain insect pests. <i>MethodsX</i> , 2022, 9, 101734. | 0.7 | 1 |
| 122 | USO DE NUTRAGREEN® COMO TRANSPORTADOR COLOIDAL PARA REDUCIR EL USO DE FERTILIZANTES Y PESTICIDAS EN PERAL. , 0, , 220-227. | | 0 |
| 123 | Fungal Secondary Metabolism. , 2021, , 54-63. | | 0 |
| 124 | Lippia spp. Essential Oil as a Control Agent against Acanthoscelides obtectus, an Insect Pest in Phaseolus vulgaris Beans. , 0, , . | | 0 |