

# C Maleita

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7954838/publications.pdf>

Version: 2024-02-01

29  
papers

443  
citations

759233  
12  
h-index

794594  
19  
g-index

30  
all docs

30  
docs citations

30  
times ranked

373  
citing authors

#	ARTICLE	IF	CITATIONS
1	First Report of Root Knot Nematodes <i>&lt; i&gt;Meloidogyne incognita&lt;/i&gt;</i> and <i>&lt; i&gt;M. javanica&lt;/i&gt;</i> Parasitizing Sweet Potato, <i>&lt; i&gt;Ipomoea batatas&lt;/i&gt;</i> , in Portugal. <i>Plant Disease</i> , 2022, 106, 2536.	1.4	1
2	Susceptibility of crop plants to the root-knot nematode <i>Meloidogyne luci</i> , a threat to agricultural productivity. <i>Phytopathologia Mediterranea</i> , 2022, 61, 169-179.	1.3	3
3	Juglone and 1,4-Naphthoquinoneâ€”Promising Nematicides for Sustainable Control of the Root Knot Nematode <i>Meloidogyne luci</i> . <i>Frontiers in Plant Science</i> , 2022, 13, .	3.6	7
4	Species-Specific Molecular Detection of the Root Knot Nematode <i>Meloidogyne luci</i> . <i>Biology</i> , 2021, 10, 775.	2.8	9
5	<i>Meloidogyne graminicola</i> â€”A Threat to Rice Production: Review Update on Distribution, Biology, Identification, and Management. <i>Biology</i> , 2021, 10, 1163.	2.8	13
6	Tomato Mi-1.2 gene confers resistance to <i>Meloidogyne luci</i> and <i>M. ethiopica</i> . <i>European Journal of Plant Pathology</i> , 2020, 156, 571-580.	1.7	14
7	<i>Ficus microcarpa</i> Bonsai â€œTiger barkâ€•Parasitized by the Root-Knot Nematode <i>Meloidogyne javanica</i> and the Spiral Nematode <i>Helicotylenchus dihystera</i> , a New Plant Host Record for Both Species. <i>Plants</i> , 2020, 9, 1085.	3.5	3
8	The quarantine rootâ€knot nematode <i>&lt; i&gt;Meloidogyne enterolobii&lt;/i&gt;</i> â€“ a potential threat to Portugal and Europe. <i>Plant Pathology</i> , 2019, 68, 1607-1615.	2.4	25
9	In vitro reproduction of <i>Pratylenchus neglectus</i> on carrot discs and investigation of its interaction with <i>Meloidogyne hispanica</i> on potato. <i>European Journal of Plant Pathology</i> , 2019, 153, 957-963.	1.7	0
10	New Hosts and Records in Portugal for the Root-Knot Nematode <i>&lt; i&gt;Meloidogyne luci&lt;/i&gt;</i> . <i>Journal of Nematology</i> , 2019, 51, 1-4.	0.9	15
11	<i>Laimaphelenchus suberensis</i> sp. nov. associated with <i>Quercus suber</i> in Portugal. <i>European Journal of Plant Pathology</i> , 2018, 150, 747-758.	1.7	7
12	On the species status of <i>Laimaphelenchus hyrcanus</i> and <i>L.Âbelgradiensis</i> . <i>Forest Pathology</i> , 2018, 48, e12425.	1.1	1
13	<i>&lt; i&gt;Meloidogyne luci&lt;/i&gt;</i> , a new rootâ€knot nematode parasitizing potato in Portugal. <i>Plant Pathology</i> , 2018, 67, 366-376.	2.4	32
14	First report of <i>Meloidogyne graminis</i> on golf courses turfgrass in Brazil. <i>PLoS ONE</i> , 2018, 13, e0192397.	2.5	9
15	Significant effects of RNAi silencing of the venom allergenâ€like protein ( <i>&lt; i&gt;Mhiâ€vapâ€1&lt;/i&gt;</i> ) of the rootâ€knot nematode <i>&lt; i&gt;Meloidogyne hispanica&lt;/i&gt;</i> in the early events of infection. <i>Plant Pathology</i> , 2017, 66, 1329-1337.	2.4	10
16	Naphthoquinones from Walnut Husk Residues Show Strong Nematicidal Activities against the Root-knot Nematode <i>&lt; i&gt;Meloidogyne hispanica&lt;/i&gt;</i> . <i>ACS Sustainable Chemistry and Engineering</i> , 2017, 5, 3390-3398.	6.7	38
17	Toxicity of the bionematicide 1,4-naphthoquinone on non-target soil organisms. <i>Chemosphere</i> , 2017, 181, 579-588.	8.2	21
18	Molecular characterization of putative parasitism genes in the plant-parasitic nematode <i>&lt; i&gt;Meloidogyne hispanica&lt;/i&gt;</i> . <i>Journal of Helminthology</i> , 2016, 90, 28-38.	1.0	6

#	ARTICLE	IF	CITATIONS
19	First report on <i>Meloidogyne chitwoodi</i> hatching inhibition activity of essential oils and essential oils fractions. <i>Journal of Pest Science</i> , 2016, 89, 207-217.	3.7	46
20	Root-lesion and root-knot nematodes parasitizing potato. <i>European Journal of Plant Pathology</i> , 2015, 141, 397-406.	1.7	19
21	First report of <i>Laimaphelenchus heidelbergi</i> (Nematoda: Aphelenchoididae) in Europe. <i>Forest Pathology</i> , 2015, 45, 76-81.	1.1	7
22	Characterization of the venom allergen-like protein (vap-1) and the fatty acid and retinol binding protein (far-1) genes in <i>Meloidogyne hispanica</i> . <i>European Journal of Plant Pathology</i> , 2014, 139, 825-836.	1.7	14
23	In vitro co-culture of <i>Solanum tuberosum</i> hairy roots with <i>Meloidogyne chitwoodi</i> : structure, growth and production of volatiles. <i>Plant Cell, Tissue and Organ Culture</i> , 2014, 118, 519-530.	2.3	12
24	Characterization of structure, growth and production of volatiles of <i>Solanum tuberosum</i> hairy roots and <i>Solanum tuberosum</i> hairy roots/ <i>Meloidogyne chitwoodi</i> in vitro co-cultures. <i>Planta Medica</i> , 2014, 80,	1.3	0
25	Biometrical, Biochemical, and Molecular Diagnosis of Portuguese <i>Meloidogyne hispanica</i> isolates. <i>Plant Disease</i> , 2012, 96, 865-874.	1.4	31
26	Host status of cultivated plants to <i>Meloidogyne hispanica</i> . <i>European Journal of Plant Pathology</i> , 2012, 133, 449-460.	1.7	33
27	Thermal requirements for the embryonic development and life cycle of <i>Meloidogyne hispanica</i> . <i>Plant Pathology</i> , 2012, 61, 1002-1010.	2.4	21
28	Effect of the Mi gene on reproduction of <i>Meloidogyne hispanica</i> on tomato genotypes. <i>Nematology</i> , 2011, 13, 939-949.	0.6	14
29	Molecular Characterization of <i>Meloidogyne hispanica</i> (Nematoda, Meloidogynidae) by Phylogenetic Analysis of Genes Within the rDNA in <i>Meloidogyne</i> spp.. <i>Plant Disease</i> , 2008, 92, 1104-1110.	1.4	29