

Yujiro Nakano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7956917/publications.pdf>

Version: 2024-02-01

24
papers

728
citations

686830

13
h-index

642321

23
g-index

24
all docs

24
docs citations

24
times ranked

1284
citing authors

#	ARTICLE	IF	CITATIONS
1	A case of ezetimibe-effective hypercholesterolemia with a novel heterozygous variant in <i>ABCG5</i> . <i>Endocrine Journal</i> , 2020, 67, 1099-1105.	0.7	3
2	A Case of Refractory Hypothyroidism due to Poor Compliance Treated with the Weekly Intravenous and Oral Levothyroxine Administration. <i>Case Reports in Endocrinology</i> , 2019, 2019, 1-6.	0.2	6
3	Incidence and predictive factors of hypoglycemia after pheochromocytoma resection. <i>International Journal of Urology</i> , 2019, 26, 273-277.	0.5	11
4	miRNA299 involvement in CYP11B2 expression in aldosterone-producing adenoma. <i>European Journal of Endocrinology</i> , 2019, 181, 69-78.	1.9	8
5	Ratio of visceral to subcutaneous fat area predicts cardiovascular events in patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2018, 9, 396-402.	1.1	32
6	Dipeptidyl peptidase 4 inhibitors attenuates the decline of skeletal muscle mass in patients with type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2018, 34, e2957.	1.7	33
7	Effect of Eplerenone on the Glomerular Filtration Rate (GFR) in Primary Aldosteronism: Sequential Changes in the GFR During Preoperative Eplerenone Treatment to Subsequent Adrenalectomy. <i>Internal Medicine</i> , 2018, 57, 2459-2466.	0.3	9
8	Luseogliflozin reduces epicardial fat accumulation in patients with type 2 diabetes: a pilot study. <i>Cardiovascular Diabetology</i> , 2017, 16, 32.	2.7	128
9	Insulin Treatment Attenuates Decline of Muscle Mass in Japanese Patients with Type 2 Diabetes. <i>Calcified Tissue International</i> , 2017, 101, 1-8.	1.5	43
10	Ipragliflozin Reduces Epicardial Fat Accumulation in Non-Obese Type 2 Diabetic Patients with Visceral Obesity: A Pilot Study. <i>Diabetes Therapy</i> , 2017, 8, 851-861.	1.2	84
11	Association of sarcopenia with both latent autoimmune diabetes in adults and type 2 diabetes: a cross-sectional study. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 992-996.	1.2	23
12	Molecular characteristics of the KCNJ5 mutated aldosterone-producing adenomas. <i>Endocrine-Related Cancer</i> , 2017, 24, 531-541.	1.6	16
13	Association of diabetic retinopathy with both sarcopenia and muscle quality in patients with type 2 diabetes: a cross-sectional study. <i>BMJ Open Diabetes Research and Care</i> , 2017, 5, e000404.	1.2	31
14	Reduction of visceral fat by liraglutide is associated with ameliorations of hepatic steatosis, albuminuria, and micro-inflammation in type 2 diabetic patients with insulin treatment: a randomized control trial. <i>Endocrine Journal</i> , 2017, 64, 269-281.	0.7	81
15	Gender difference in the impact of gynoid and android fat masses on the progression of hepatic steatosis in Japanese patients with type 2 diabetes. <i>BMC Obesity</i> , 2017, 4, 27.	3.1	5
16	Indirect measure of visceral adiposity (A Body Shape Index™ (ABSI)) is associated with arterial stiffness in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2016, 4, e000188.	1.2	64
17	Expression of inflammation-related genes in aldosterone-producing adenomas with KCNJ5 mutation. <i>Biochemical and Biophysical Research Communications</i> , 2016, 476, 614-619.	1.0	3
18	Retrograde pyelonephritis and lumbar spondylitis as a result of <i>Salmonella typhi</i> in a type 2 diabetes patient with neurogenic bladder. <i>Journal of Diabetes Investigation</i> , 2016, 7, 436-439.	1.1	10

#	ARTICLE	IF	CITATIONS
19	Increased visceral adiposity with normal weight is associated with the prevalence of nonalcoholic fatty liver disease in Japanese patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2016, 7, 607-614.	1.1	13
20	Is visceral adiposity a modifier for the impact of blood pressure on arterial stiffness and albuminuria in patients with type 2 diabetes?. <i>Cardiovascular Diabetology</i> , 2016, 15, 10.	2.7	9
21	Clinical relevance of dual-energy X-ray absorptiometry (DXA) as a simultaneous evaluation of fatty liver disease and atherosclerosis in patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2016, 15, 64.	2.7	25
22	A Case of Cushing's Syndrome with Multiple Adrenocortical Adenomas Composed of Compact Cells and Clear Cells. <i>Endocrine Pathology</i> , 2016, 27, 136-141.	5.2	0
23	High visceral fat with low subcutaneous fat accumulation as a determinant of atherosclerosis in patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2015, 14, 136.	2.7	61
24	Impact of increased visceral adiposity with normal weight on the progression of arterial stiffness in Japanese patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2015, 3, e000081.	1.2	30