Samuel Niedermann

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/7959674/samuel-niedermann-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60
papers

1,930
citations

26
h-index

g-index

43
g-index

4.52
ext. papers

ext. citations

avg, IF

L-index

#	Paper	IF	Citations
60	The SPICE project: Calibrated cosmogenic 26Al production rates and cross-calibrated 26Al /10Be, 26Al/14C, and 26Al/21Ne ratios in quartz from the SP basalt flow, AZ, USA. <i>Quaternary Geochronology</i> , 2022 , 67, 101218	2.7	O
59	Quaternary landscape evolution in a tectonically active rift basin (paleo-lake Mweru, south-central Africa). <i>Geomorphology</i> , 2021 , 381, 107669	4.3	1
58	Nature Does the AveragingIh-Situ Produced 10Be, 21Ne, and 26Al in a Very Young River Terrace. <i>Geosciences (Switzerland)</i> , 2020 , 10, 237	2.7	4
57	Age and source of the Kle/ldemilarsenopyrite-gold mineralization (Menderes Massif, Turkey) determined by Re-Os-He isotopes. <i>Ore Geology Reviews</i> , 2020 , 118, 103333	3.2	
56	The SPICE project: Production rates of cosmogenic 21Ne, 10Be, and 14C in quartz from the 72 ka SP basalt flow, Arizona, USAu. <i>Quaternary Geochronology</i> , 2019 , 54, 101019	2.7	5
55	Formation of epithermal Sn-Ag-(Zn) vein-type mineralization at the Pirquitas deposit, NW Argentina: Fluid inclusion and noble gas isotopic constraints. <i>Chemical Geology</i> , 2019 , 508, 78-91	4.2	5
54	Monitoring of helium and carbon isotopes in the western Eger Rift area (Czech Republic): Relationships with the 2014 seismic activity and indications for recent (2000 2 016) magmatic unrest. <i>Chemical Geology</i> , 2018 , 482, 131-145	4.2	16
53	Applying stable cosmogenic 21Ne to understand surface processes in deep geological time (107🛮 08 yr). <i>Earth and Planetary Science Letters</i> , 2018 , 498, 266-274	5.3	3
52	Glacial chronology and production rate cross-calibration of five cosmogenic nuclide and mineral systems from the southern Central Andean Plateau. <i>Earth and Planetary Science Letters</i> , 2018 , 500, 242	-253	12
51	Regional distribution pattern of carbon and helium isotopes from different volcanic fields in the French Massif Central: Evidence for active mantle degassing and water transport. <i>Chemical Geology</i> , 2017 , 469, 4-18	4.2	7
50	Subaqueous hot springs in Kycelż Lake, Dalyan Channel and Fethiye-Glek Bay (SW Turkey): Locations, chemistry and origins. <i>Journal of Volcanology and Geothermal Research</i> , 2017 , 345, 81-97	2.8	3
49	Dating the incision of the Yangtze River gorge at the First Bend using three-nuclide burial ages. <i>Geophysical Research Letters</i> , 2016 , 43, 101-110	4.9	38
48	Surface exposure dating of Holocene basalt flows and cinder cones in the Kula volcanic field (Western Turkey) using cosmogenic 3He and 10Be. <i>Quaternary Geochronology</i> , 2016 , 34, 81-91	2.7	18
47	He, Ne and Ar isotope signatures of mid-ocean ridge basalts and their implications for upper mantle structure: A case study from the Mid-Atlantic Ridge at 4112°S. <i>Geochimica Et Cosmochimica Acta</i> , 2016 , 183, 94-105	5.5	6
46	The fluid budget of a continental plate boundary fault: Quantification from the Alpine Fault, New Zealand. <i>Earth and Planetary Science Letters</i> , 2016 , 445, 125-135	5.3	30
45	The cosmogenic record of mountain erosion transmitted across a foreland basin: Source-to-sink analysis of in situ 10 Be, 26 Al and 21 Ne in sediment of the Po river catchment. <i>Earth and Planetary Science Letters</i> , 2016 , 452, 258-271	5.3	31
44	The Touissit-Bou Beker Mississippi Valley-Type District of Northeastern Morocco: Relationships to the Messinian Salinity Crisis, Late Neogene-Quaternary Alkaline Magmatism, and Buoyancy-Driven Fluid Convection. <i>Economic Geology</i> , 2015 , 110, 1455-1484	4.3	17

43	Interlaboratory comparison of cosmogenic 21 Ne in quartz. <i>Quaternary Geochronology</i> , 2015 , 26, 20-28	2.7	31
42	Noble gases in sulfide deposits of modern deep-sea hydrothermal systems: Implications for heat fluxes and hydrothermal fluid processes. <i>Chemical Geology</i> , 2015 , 409, 1-11	4.2	17
41	Erosion during extreme flood events dominates Holocene canyon evolution in northeast Iceland. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 2355-60	11.5	70
40	Noble gas residence times of saline waters within crystalline bedrock, Outokumpu Deep Drill Hole, Finland. <i>Geochimica Et Cosmochimica Acta</i> , 2014 , 145, 159-174	5.5	25
39	Surface exposure dating of young basalts (1월00 ka) in the San Francisco volcanic field (Arizona, USA) using cosmogenic 3He and 21Ne. <i>Quaternary Geochronology</i> , 2014 , 19, 87-105	2.7	20
38	Tectonic implications of fluvial incision and pediment deformation at the northern margin of the Central Anatolian Plateau based on multiple cosmogenic nuclides. <i>Tectonics</i> , 2013 , 32, 1107-1120	4.3	27
37	40Ar/39Ar dating of the SP and Bar Ten lava flows AZ, USA: Laying the foundation for the SPICE cosmogenic nuclide production-rate calibration project. <i>Quaternary Geochronology</i> , 2013 , 18, 158-172	2.7	14
36	Indications for the existence of different magmatic reservoirs beneath the Eifel area (Germany): A multi-isotope (C, N, He, Ne, Ar) approach. <i>Chemical Geology</i> , 2013 , 356, 193-208	4.2	30
35	Neon identifies two billion year old fluid component in Kaapvaal Craton. <i>Chemical Geology</i> , 2011 , 283, 287-296	4.2	50
34	Regional 10Be production rate calibration for the past 12 ka deduced from the radiocarbon-dated GrElandsura and Russenes rock avalanches at 69° N, Norway. <i>Quaternary Geochronology</i> , 2011 , 6, 437-45	5 2 -7	99
33	Inter-comparison of cosmogenic in-situ 3He, 21Ne and 36Cl at low latitude along an altitude transect on the SE slope of Kilimanjaro volcano (3°S, Tanzania). <i>Quaternary Geochronology</i> , 2011 , 6, 425-	-4376	18
32	Late PleistoceneBarly Holocene paleoseismicity deduced from lake sediment deformation and coeval landsliding in the CalchaquBs valleys, NW Argentina 2011 ,		5
31	Landslide Dams in the Central Andes of Argentina (Northern Patagonia and the Argentine Northwest). <i>Lecture Notes in Earth Sciences</i> , 2011 , 147-176		16
30	Late Pleistocene landscape evolution in south-central Chile constrained by luminescence and stable cosmogenic nuclide dating. <i>Bulletin of the Geological Society of America</i> , 2010 , 122, 1235-1247	3.9	10
29	Eruption ages of Las Tres VEgenes volcano (Baja California): A tale of two helium isotopes. <i>Quaternary Geochronology</i> , 2010 , 5, 503-511	2.7	54
28	Multiple sources for mineralizing fluids in the Charmitan gold(-tungsten) mineralization (Uzbekistan). <i>Mineralium Deposita</i> , 2010 , 45, 667-682	4.8	30
27	Evaluation of cosmogenic 3He and 21Ne production rates in olivine and pyroxene from two Pleistocene basalt flows, western Grand Canyon, AZ, USA. <i>Quaternary Geochronology</i> , 2009 , 4, 475-492	2.7	24
26	Natural laboratory NW Bohemia: Comprehensive fluid studies between 1992 and 2005 used to trace geodynamic processes. <i>Geochemistry, Geophysics, Geosystems</i> , 2008 , 9, n/a-n/a	3.6	46

25	Plumefidge interaction revisited: Evidence for melt mixing from He, Ne and Ar isotope and abundance systematics. <i>Earth and Planetary Science Letters</i> , 2008 , 268, 424-432	5.3	10
24	Seismically induced changes of the fluid signature detected by a multi-isotope approach (He, CO2, CH4, N2) at the Wettinquelle, Bad Brambach (central Europe). <i>Journal of Geophysical Research</i> , 2007 , 112,		25
23	Neon and helium isotopes as tracers of mantle reservoirs and mantle dynamics. <i>Earth and Planetary Science Letters</i> , 2007 , 258, 334-344	5.3	13
22	Climatic versus tectonic control on river incision at the margin of NE Tibet: 10Be exposure dating of river terraces at the mountain front of the Qilian Shan. <i>Journal of Geophysical Research</i> , 2006 , 111, n/a-	n/a	62
21	Origin of ore fluids in the Muruntau gold system: Constraints from noble gas, carbon isotope and halogen data. <i>Geochimica Et Cosmochimica Acta</i> , 2006 , 70, 5356-5370	5.5	64
20	Evidence for ascending upper mantle-derived melt beneath the Cheb basin, central Europe. <i>Geophysical Research Letters</i> , 2005 , 32,	4.9	29
19	Correction to E vidence for ascending upper mantle-derived melt beneath the Cheb basin, central Europe [Geophysical Research Letters, 2005 , 32, n/a-n/a	4.9	3
18	Implications of the fault scaling law for the growth of topography: mountain ranges in the broken foreland of north-east Tibet. <i>Terra Nova</i> , 2004 , 16, 157-162	3	61
17	Rock avalanching into a landslide-dammed lake causing multiple dam failure in Las Conchas valley (NW Argentina) Levidence from surface exposure dating and stratigraphic analyses. <i>Landslides</i> , 2004 , 1, 113	6.6	92
16	Late Pleistocene/Holocene slip rate of the Zhangye thrust (Qilian Shan, China) and implications for the active growth of the northeastern Tibetan Plateau. <i>Tectonics</i> , 2004 , 23, n/a-n/a	4.3	105
15	Evidence for a nitrogen flux directly derived from the European subcontinental mantle in the Western Eger Rift, Central Europe. <i>Geochimica Et Cosmochimica Acta</i> , 2004 , 68, 4935-4947	5.5	47
14	Noble gases in olivine phenocrysts from drill core samples of the Hawaii Scientific Drilling Project (HSDP) pilot and main holes (Mauna Loa and Mauna Kea, Hawaii). <i>Geochemistry, Geophysics, Geosystems</i> , 2003 , 4,	3.6	15
13	16. Cosmic-Ray-Produced Noble Gases in Terrestrial Rocks: Dating Tools for Surface Processes 2002 , 731-784		11
12	Low slip rates and long-term preservation of geomorphic features in Central Asia. <i>Nature</i> , 2002 , 417, 428-32	50.4	166
11	21Ne versus 10Be and 26Al exposure ages of fluvial terraces: the influence of crustal Ne in quartz. <i>Earth and Planetary Science Letters</i> , 2002 , 201, 575-591	5.3	48
10	Neotectonics and catastrophic failure of mountain fronts in the southern intra-Andean Puna Plateau, Argentina. <i>Geology</i> , 2001 , 29, 619	5	75
9	Tephrochronologic Constraints on Temporal Distribution of Large Landslides in Northwest Argentina. <i>Journal of Geology</i> , 2000 , 108, 35-52	2	52
8	The 21Ne production rate in quartz revisited. <i>Earth and Planetary Science Letters</i> , 2000 , 183, 361-364	5.3	50

LIST OF PUBLICATIONS

7	Atmospheric noble gases in volcanic glasses from the southern Lau Basin: origin from the subducting slab?. <i>Earth and Planetary Science Letters</i> , 1998 , 160, 297-309	5.3	36
6	Anomalously nucleogenic neon in North Chile Ridge basalt glasses suggesting a previously degassed mantle source. <i>Earth and Planetary Science Letters</i> , 1998 , 160, 447-462	5.3	26
5	Noble gas evidence for a lower mantle component in MORBs from the southern East Pacific Rise: Decoupling of helium and neon isotope systematics. <i>Geochimica Et Cosmochimica Acta</i> , 1997 , 61, 2697-2	2715	155
4	Origin and Evolution of Gas in Salt Beds of a Potash Mine. <i>Advances in Geosciences</i> ,54, 15-21		2
3	The Schlaining quartz-stibnite deposit, Eastern Alps, Austria: constraints from conventional and infrared microthermometry and isotope and crush-leach analyses of fluid inclusions. <i>Mineralium Deposita</i> ,1	4.8	0
2	Microthermometry and noble gas isotope analysis of magmatic fluid inclusions in the Kerman porphyry Cu deposits, Iran: constraints on the source of ore-forming fluids. <i>Mineralium Deposita</i> ,1	4.8	1
1	A geochemical study of the Sweet Home mine, Colorado Mineral Belt, USA: formation of deep hydrothermal veinEype molybdenum greisen and base metal mineralization. <i>Mineralium Deposita</i> ,1	4.8	О