

Marguerite Hatch

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/796355/publications.pdf>

Version: 2024-02-01

56
papers

1,646
citations

279701

23
h-index

302012

39
g-index

57
all docs

57
docs citations

57
times ranked

995
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | The anion exchanger PAT-1 (Slc26a6) does not participate in oxalate or chloride transport by mouse large intestine. <i>Pflugers Archiv European Journal of Physiology</i> , 2021, 473, 95-106. | 1.3 | 7 |
| 2 | The role of NHE3 (Slc9a3) in oxalate and sodium transport by mouse intestine and regulation by cAMP. <i>Physiological Reports</i> , 2021, 9, e14828. | 0.7 | 3 |
| 3 | Extracellular Vesicle Analysis by Paper Spray Ionization Mass Spectrometry. <i>Metabolites</i> , 2021, 11, 308. | 1.3 | 9 |
| 4 | Forty Years of <i>Oxalobacter formigenes</i> , a Gutsy Oxalate-Degrading Specialist. <i>Applied and Environmental Microbiology</i> , 2021, 87, e0054421. | 1.4 | 32 |
| 5 | Oxalate Flux Across the Intestine: Contributions from Membrane Transporters. , 2021, 12, 2835-2875. | | 3 |
| 6 | Induction of enteric oxalate secretion by <i>Oxalobacter formigenes</i> in mice does not require the presence of either apical oxalate transport proteins Slc26A3 or Slc26A6. <i>Urolithiasis</i> , 2020, 48, 1-8. | 1.2 | 12 |
| 7 | <i>Oxalobacter formigenes</i> produces metabolites and lipids undetectable in oxalotrophic <i>Bifidobacterium animalis</i> . <i>Metabolomics</i> , 2020, 16, 122. | 1.4 | 7 |
| 8 | Metabolomic Alteration in the Mouse Distal Colonic Mucosa after Oral Gavage with <i>Oxalobacter formigenes</i> . <i>Metabolites</i> , 2020, 10, 405. | 1.3 | 6 |
| 9 | Oxalate transport by the mouse intestine in vitro is not affected by chronic challenges to systemic acid-base homeostasis. <i>Urolithiasis</i> , 2019, 47, 243-254. | 1.2 | 5 |
| 10 | Metabolomic profiling of oxalate-degrading probiotic <i>Lactobacillus acidophilus</i> and <i>Lactobacillus gasseri</i> . <i>PLoS ONE</i> , 2019, 14, e0222393. | 1.1 | 36 |
| 11 | ¹²⁵ Iodide as a surrogate tracer for epithelial chloride transport by the mouse large intestine in vitro. <i>Experimental Physiology</i> , 2019, 104, 334-344. | 0.9 | 5 |
| 12 | Metabolomic and lipidomic characterization of <i>Oxalobacter formigenes</i> strains HC1 and OxWR by UHPLC-HRMS. <i>Analytical and Bioanalytical Chemistry</i> , 2019, 411, 4807-4818. | 1.9 | 20 |
| 13 | Absence of the sulfate transporter SAT-1 has no impact on oxalate handling by mouse intestine and does not cause hyperoxaluria or hyperoxalemia. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 316, G82-C94. | 1.6 | 16 |
| 14 | ¹²⁵ Iodide as a Surrogate for ³⁶ Chloride in Tracing Transepithelial Intestinal Chloride Transport. <i>FASEB Journal</i> , 2019, 33, 575.13. | 0.2 | 0 |
| 15 | Title is missing!. , 2019, 14, e0222393. | | 0 |
| 16 | Title is missing!. , 2019, 14, e0222393. | | 0 |
| 17 | Title is missing!. , 2019, 14, e0222393. | | 0 |
| 18 | Title is missing!. , 2019, 14, e0222393. | | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Oxalobacter formigenes colonization normalizes oxalate excretion in a gastric bypass model of hyperoxaluria. <i>Surgery for Obesity and Related Diseases</i> , 2017, 13, 1152-1157. | 1.0 | 24 |
| 20 | The role of intestinal oxalate transport in hyperoxaluria and the formation of kidney stones in animals and man. <i>Urolithiasis</i> , 2017, 45, 89-108. | 1.2 | 68 |
| 21 | Genome Sequence of Oxalobacter formigenes Strain OXCC13. <i>Genome Announcements</i> , 2017, 5, . | 0.8 | 4 |
| 22 | Loss of the anion exchanger DRA (Slc26a3), or PAT1 (Slc26a6), alters sulfate transport by the distal ileum and overall sulfate homeostasis. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 313, G166-G179. | 1.6 | 17 |
| 23 | Genome Sequence of Oxalobacter formigenes Strain HC-1. <i>Genome Announcements</i> , 2017, 5, . | 0.8 | 3 |
| 24 | Gut microbiota and oxalate homeostasis. <i>Annals of Translational Medicine</i> , 2017, 5, 36-36. | 0.7 | 48 |
| 25 | The mechanistic basis of hyperoxaluria following gastric bypass in obese rats. <i>Urolithiasis</i> , 2016, 44, 221-230. | 1.2 | 11 |
| 26 | Bifidobacterium animalis subsp. lactis decreases urinary oxalate excretion in a mouse model of primary hyperoxaluria. <i>Urolithiasis</i> , 2015, 43, 107-117. | 1.2 | 41 |
| 27 | Effects of acid-base variables and the role of carbonic anhydrase on oxalate secretion by the mouse intestine in vitro. <i>Physiological Reports</i> , 2015, 3, e12282. | 0.7 | 7 |
| 28 | Chronic metabolic acidosis reduces urinary oxalate excretion and promotes intestinal oxalate secretion in the rat. <i>Urolithiasis</i> , 2015, 43, 489-499. | 1.2 | 11 |
| 29 | Kidney stone incidence and metabolic urinary changes after modern bariatric surgery: review of clinical studies, experimental models, and prevention strategies. <i>Surgery for Obesity and Related Diseases</i> , 2014, 10, 734-742. | 1.0 | 53 |
| 30 | Intestinal adaptations in chronic kidney disease and the influence of gastric bypass surgery. <i>Experimental Physiology</i> , 2014, 99, 1163-1167. | 0.9 | 17 |
| 31 | A human strain of Oxalobacter (HC-1) promotes enteric oxalate secretion in the small intestine of mice and reduces urinary oxalate excretion. <i>Urolithiasis</i> , 2013, 41, 379-384. | 1.2 | 64 |
| 32 | Steatorrhea and Hyperoxaluria Occur after Gastric Bypass Surgery in Obese Rats Regardless of Dietary Fat or Oxalate. <i>Journal of Urology</i> , 2013, 190, 1102-1109. | 0.2 | 40 |
| 33 | Transcellular oxalate and Cl ⁻ absorption in mouse intestine is mediated by the DRA anion exchanger Slc26a3, and DRA deletion decreases urinary oxalate. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 305, G520-G527. | 1.6 | 56 |
| 34 | Sulfate secretion and chloride absorption are mediated by the anion exchanger DRA (Slc26a3) in the mouse cecum. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 305, G172-G184. | 1.6 | 21 |
| 35 | Hyperoxaluric rats do not exhibit alterations in renal expression patterns of Slc26a1 (SAT1) mRNA or protein. <i>Urological Research</i> , 2012, 40, 647-654. | 1.5 | 7 |
| 36 | Enteric oxalate elimination is induced and oxalate is normalized in a mouse model of primary hyperoxaluria following intestinal colonization with Oxalobacter. <i>American Journal of Physiology - Renal Physiology</i> , 2011, 300, G461-G469. | 1.6 | 127 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Parsing apical oxalate exchange in Caco-2BBE1 monolayers: siRNA knockdown of SLC26A6 reveals the role and properties of PAT-1. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 297, G918-G929. | 1.6 | 32 |
| 38 | Enteric oxalate secretion is not directly mediated by the human CFTR chloride channel. <i>Urological Research</i> , 2008, 36, 127-131. | 1.5 | 8 |
| 39 | The Roles and Mechanisms of Intestinal Oxalate Transport in Oxalate Homeostasis. <i>Seminars in Nephrology</i> , 2008, 28, 143-151. | 0.6 | 71 |
| 40 | Increased colonic sodium absorption in rats with chronic renal failure is partially mediated by AT ₁ receptor agonism. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, G348-G356. | 1.6 | 20 |
| 41 | Ileal oxalate absorption and urinary oxalate excretion are enhanced in Slc26a6 null mice. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 290, G719-G728. | 1.6 | 142 |
| 42 | Lipid peroxidation is not the underlying cause of renal injury in hyperoxaluric rats. <i>Kidney International</i> , 2005, 68, 2629-2638. | 2.6 | 29 |
| 43 | Intestinal transport of an obdurate anion: oxalate. <i>Urological Research</i> , 2005, 33, 1-16. | 1.5 | 99 |
| 44 | Ethylene glycol induces hyperoxaluria without metabolic acidosis in rats. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 289, F536-F543. | 1.3 | 57 |
| 45 | Serum oxalate in human beings and rats as determined with the use of ion chromatography. <i>Translational Research</i> , 2004, 144, 45-52. | 2.4 | 22 |
| 46 | Angiotensin II involvement in adaptive enteric oxalate excretion in rats with chronic renal failure induced by hyperoxaluria. <i>Urological Research</i> , 2003, 31, 426-432. | 1.5 | 24 |
| 47 | Renal and Intestinal Handling of Oxalate following Oxalate Loading in Rats. <i>American Journal of Nephrology</i> , 2003, 23, 18-26. | 1.4 | 43 |
| 48 | Muscarinic down-regulation of cAMP-stimulated potassium ion secretion by rabbit distal colon. <i>Pflugers Archiv European Journal of Physiology</i> , 2000, 440, 243-252. | 1.3 | 5 |
| 49 | Conductive pathways for chloride and oxalate in rabbit ileal brush-border membrane vesicles. <i>American Journal of Physiology - Cell Physiology</i> , 1998, 275, C748-C757. | 2.1 | 51 |
| 50 | Local upregulation of colonic angiotensin II receptors enhances potassium excretion in chronic renal failure. <i>American Journal of Physiology - Renal Physiology</i> , 1998, 274, F275-F282. | 1.3 | 22 |
| 51 | Mechanisms of oxalate absorption and secretion across the rabbit distal colon. <i>Pflugers Archiv European Journal of Physiology</i> , 1994, 426, 101-109. | 1.3 | 50 |
| 52 | Enhanced Enteric Excretion of Urate in Rats with Chronic Renal Failure. <i>Clinical Science</i> , 1994, 86, 511-516. | 1.8 | 62 |
| 53 | Oxalate status in stone-formers. <i>Urological Research</i> , 1993, 21, 55-59. | 1.5 | 33 |
| 54 | Characteristics of the transport of oxalate and other ions across rabbit proximal colon. <i>Pflugers Archiv European Journal of Physiology</i> , 1993, 423, 206-212. | 1.3 | 36 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Spectrophotometric determination of oxalate in whole blood. <i>Clinica Chimica Acta</i> , 1990, 193, 199-202. | 0.5 | 18 |
| 56 | Oxalate transport across the isolated rat colon. A re-examination. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1980, 600, 838-843. | 1.4 | 42 |