

Sergio T RodrÃ-iguez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7963879/publications.pdf>

Version: 2024-02-01

11
papers

154
citations

1307594

7
h-index

1281871

11
g-index

11
all docs

11
docs citations

11
times ranked

228
citing authors

#	ARTICLE	IF	CITATIONS
1	Klotho expression in peripheral blood circulating cells is associated with vascular and systemic inflammation in atherosclerotic vascular disease. <i>Scientific Reports</i> , 2022, 12, 8422.	3.3	22
2	High serum caspase-3 levels in hepatocellular carcinoma prior to liver transplantation and high mortality risk during the first year after liver transplantation. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 635-640.	3.1	2
3	Low Serum Melatonin Levels Prior to Liver Transplantation in Patients with Hepatocellular Carcinoma are Associated with Lower Survival after Liver Transplantation. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1696.	4.1	3
4	FGF23 and Klotho Levels are Independently Associated with Diabetic Foot Syndrome in Type 2 Diabetes Mellitus. <i>Journal of Clinical Medicine</i> , 2019, 8, 448.	2.4	15
5	High serum soluble CD40L levels previously to liver transplantation in patients with hepatocellular carcinoma are associated with mortality at one year. <i>Journal of Critical Care</i> , 2018, 43, 316-320.	2.2	7
6	Serum total antioxidant capacity prior to liver transplantation for hepatocellular carcinoma is associated with 1-year liver transplantation survival. <i>Journal of International Medical Research</i> , 2018, 46, 2641-2649.	1.0	3
7	Patients with high serum substance P levels previously to liver transplantation for hepatocellular carcinoma have higher risk of one-year liver transplantation mortality. <i>Oncotarget</i> , 2018, 9, 21552-21559.	1.8	3
8	Soluble levels and endogenous vascular gene expression of <i>KLOTHO</i> are related to inflammation in human atherosclerotic disease. <i>Clinical Science</i> , 2017, 131, 2601-2609.	4.3	37
9	Association between Pre-Transplant Serum Malondialdehyde Levels and Survival One Year after Liver Transplantation for Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2016, 17, 500.	4.1	23
10	Prognostic Value of Serum Caspase-Cleaved Cytokeratin-18 Levels before Liver Transplantation for One-Year Survival of Patients with Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1524.	4.1	16
11	Serum soluble CD40 Ligand levels are associated with severity and mortality of brain trauma injury patients. <i>Thrombosis Research</i> , 2014, 134, 832-836.	1.7	23