Maiko Fujimori

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7965786/publications.pdf

Version: 2024-02-01

51	1,236	15	34
papers	citations	h-index	g-index
54	54	54	1543
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Preferences of Cancer Patients Regarding Communication of Bad News: A Systematic Literature Review. Japanese Journal of Clinical Oncology, 2009, 39, 201-216.	0.6	196
2	Effect of Communication Skills Training Program for Oncologists Based on Patient Preferences for Communication When Receiving Bad News: A Randomized Controlled Trial. Journal of Clinical Oncology, 2014, 32, 2166-2172.	0.8	190
3	Preferences of cancer patients regarding the disclosure of bad news. Psycho-Oncology, 2007, 16, 573-581.	1.0	141
4	Good communication with patients receiving bad news about cancer in Japan. Psycho-Oncology, 2005, 14, 1043-1051.	1.0	59
5	A cross-sectional study of psychological distress, burnout, and the associated risk factors in hospital pharmacists in Japan. BMC Public Health, 2016, 16, 534.	1.2	55
6	Preferences of advanced cancer patients for communication on anticancer treatment cessation and the transition to palliative care. Cancer, 2015, 121, 4240-4249.	2.0	52
7	Patients' perception of the usefulness of a question prompt sheet for advanced cancer patients when deciding the initial treatment: a randomized, controlled trial. Psycho-Oncology, 2012, 21, 706-713.	1.0	47
8	Development and preliminary evaluation of communication skills training program for oncologists based on patient preferences for communicating bad news. Palliative and Supportive Care, 2014, 12, 379-386.	0.6	45
9	Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale – Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. Journal of Psychosomatic Research, 2020. 129. 109892.	1.2	33
10	Adding a Wider Range and "Hope for the Best, and Prepare for the Worst―Statement: Preferences of Patients with Cancer for Prognostic Communication. Oncologist, 2019, 24, e943-e952.	1.9	30
11	Prevalence and predictive factors of depression and anxiety in patients with pancreatic cancer: a longitudinal study. Japanese Journal of Clinical Oncology, 2016, 46, 71-77.	0.6	28
12	Cancer screening participation in schizophrenic outpatients and the influence of their functional disability on the screening rate: ⟨scp⟩A⟨/scp⟩ crossâ€sectional study in ⟨scp⟩J⟨/scp⟩apan. Psychiatry and Clinical Neurosciences, 2017, 71, 813-825.	1.0	28
13	Effectiveness of Japanese SHARE model in improving Taiwanese healthcare personnel's preference for cancer truth telling. Psycho-Oncology, 2014, 23, 259-265.	1.0	25
14	Suicide prevention strategies in Japan: A 15-year review (1998–2013). Journal of Public Health Policy, 2015, 36, 52-66.	1.0	19
15	Changes in Physicians' Intrapersonal Empathy After a Communication Skills Training in Japan. Academic Medicine, 2018, 93, 1821-1826.	0.8	19
16	Prevalence and associated factors of perceived cancer-related stigma in Japanese cancer survivors. Japanese Journal of Clinical Oncology, 2020, 50, 1325-1329.	0.6	19
17	Depression prevalence using the HADS-D compared to SCID major depression classification: An individual participant data meta-analysis. Journal of Psychosomatic Research, 2020, 139, 110256.	1.2	19
18	Breakthrough Cancer Pain Influences General Activities and Pain Management: A Comparison of Patients with and without Breakthrough Cancer Pain. Journal of Palliative Medicine, 2018, 21, 1636-1640.	0.6	17

#	Article	IF	CITATIONS
19	Unmet supportive care needs and associated factors among young adult cancer patients in Japan. BMC Cancer, 2021, 21, 17.	1.1	17
20	Smoking among adults with serious psychological distress: Analysis of anonymized data from a national cross-sectional survey in Japan. Journal of Affective Disorders, 2018, 239, 131-137.	2.0	14
21	Explicit prognostic disclosure to Asian women with breast cancer: A randomized, scripted videoâ€vignette study (Jâ€6UPPORT1601). Cancer, 2019, 125, 3320-3329.	2.0	14
22	Encouraging participation in colorectal cancer screening for people with schizophrenia: A randomized controlled trial. Acta Psychiatrica Scandinavica, 2021, 144, 318-328.	2.2	14
23	Factors associated with patient preferences for communication of bad news. Palliative and Supportive Care, 2017, 15, 328-335.	0.6	13
24	Truth telling in Taiwanese cancer care: patients' and families' preferences and their experiences of doctors' practices. Psycho-Oncology, 2017, 26, 999-1005.	1.0	12
25	Association between serious psychological distress and nonparticipation in cancer screening and the modifying effect of socioeconomic status: Analysis of anonymized data from a national crossâ€sectional survey in Japan. Cancer, 2018, 124, 555-562.	2.0	12
26	Low Cancer Screening Rates among Japanese People with Schizophrenia: A Cross-Sectional Study. Tohoku Journal of Experimental Medicine, 2018, 244, 209-218.	0.5	12
27	Comparison of Indicators for Achievement of Pain Control With a Personalized Pain Goal in a Comprehensive Cancer Center. Journal of Pain and Symptom Management, 2018, 55, 1159-1164.	0.6	11
28	Which Physicians' Behaviors on Death Pronouncement Affect Family-Perceived Physician Compassion? A Randomized, Scripted, Video-Vignette Study. Journal of Pain and Symptom Management, 2018, 55, 189-197.e4.	0.6	11
29	A nationally representative crossâ€sectional survey on health information access for consumers in Japan: A protocol for the INFORM Study. World Medical and Health Policy, 2022, 14, 225-275.	0.9	11
30	Integrated communication support program for oncologists, caregivers and patients with rapidly progressing advanced cancer to promote patient-centered communication: J-SUPPORT 1904 study protocol for a randomised controlled trial. BMJ Open, 2020, 10, e036745.	0.8	8
31	Exploring autistic-like traits relating to empathic attitude and psychological distress in hospital pharmacists. International Journal of Clinical Pharmacy, 2015, 37, 1258-1266.	1.0	7
32	Assessing the need for a question prompt list that encourages end-of-life discussions between patients with advanced cancer and their physicians: A focus group interview study. Palliative and Supportive Care, 2022, 20, 564-569.	0.6	7
33	Death by suicide, other externally caused injuries and cardiovascular diseases within 6Âmonths of cancer diagnosis (J-SUPPORT 1902). Japanese Journal of Clinical Oncology, 2021, 51, 744-752.	0.6	6
34	Change in smoking cessation stage over $1\hat{a}\in\%$ year in patients with schizophrenia: a follow up study in Japan. BMC Psychiatry, 2019, 19, 367.	1.1	5
35	Cancer care for people with mental disorders: A qualitative survey among cancer care and psychiatric care professionals in Japan. Psycho-Oncology, 2021, 30, 2060-2066.	1.0	5
36	Emotional Intelligence and its Effect on Pharmacists and Pharmacy Students with Autistic-like Traits. American Journal of Pharmaceutical Education, 2017, 81, 74.	0.7	5

3

#	Article	IF	CITATIONS
37	A randomised controlled trial of a case management approach to encourage participation in colorectal cancer screening for people with schizophrenia in psychiatric outpatient clinics: study protocol for the J-SUPPORT 1901 (ACCESS) study. BMJ Open, 2019, 9, e032955.	0.8	4
38	Cancer Survivors' Experiences of Changes in Employment and Income, and the Factors Affecting Their Experience, Quality of Life and Mental Health. Palliative Care Research, 2018, 13, 209-218.	0.0	4
39	The Effects of Adding Reassurance Statements: Cancer Patients' Preferences for Phrases in End-of-Life Discussions. Journal of Pain and Symptom Management, 2019, 57, 1121-1129.	0.6	3
40	Effectiveness of communication skill training on cancer truthâ€ŧelling for advanced practice nurses in Taiwan: A pilot study. Psycho-Oncology, 2021, 30, 765-772.	1.0	3
41	Patients' acceptability and implementation outcomes of a case management approach to encourage participation in colorectal cancer screening for people with schizophrenia: a qualitative secondary analysis of a mixed-method randomised clinical trial. BMJ Open, 2022, 12, e060621.	0.8	3
42	Characteristics of cancer patients who died by suicide in the Tokyo metropolitan area. Japanese Journal of Clinical Oncology, 2017, 47, 458-462.	0.6	2
43	Suicide, other externally caused injuries and cardiovascular death following a cancer diagnosis: study protocol for a nationwide population-based study in Japan (J-SUPPORT 1902). BMJ Open, 2019, 9, e030681.	0.8	2
44	Trends in smoking rates among individuals with serious psychological distress: Analysis of data from a Japanese national survey, 2007–2016. Psychiatry Research, 2020, 291, 113225.	1.7	2
45	Prevalence and associated factors of psychological distress among young adult cancer patients in Japan. Palliative and Supportive Care, 2023, 21, 93-99.	0.6	2
46	Factors affecting the implementation of guideline-based prophylactic antiemetic therapy for chemotherapy-induced nausea and vomiting in Japan: a protocol for a hospital-based qualitative study. BMJ Open, 2022, 12, e055473.	0.8	2
47	Issues of cancer care in people with mental disorders as perceived by cancer care providers: A quantitative questionnaire survey. Psycho-Oncology, 0, , .	1.0	2
48	Assessing the need for a question prompt list that encourages end-of-life discussions between patients with advanced cancer and their physicians: A focus group interview study $\hat{a} \in ERRATUM$. Palliative and Supportive Care, 2022, , 1-3.	0.6	1
49	Reply to B. Gyawali et al. Journal of Clinical Oncology, 2015, 33, 223-224.	0.8	0
50	Confidence in communicating with patients with cancer mediates the relationship between rehabilitation therapists' autistic-like traits and perceived difficulty in communication. Palliative and Supportive Care, 2019, 17, 186-194.	0.6	0
51	Validity and reliability of the Japanese version of the Peace, Equanimity, and Acceptance in the Cancer Experience (PEACE) questionnaire. Palliative and Supportive Care, 2021, , 1-7.	0.6	o