

Yuki Someya

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7974388/publications.pdf>

Version: 2024-02-01

13
papers

165
citations

1306789

7
h-index

1125271

13
g-index

13
all docs

13
docs citations

13
times ranked

198
citing authors

#	ARTICLE	IF	CITATIONS
1	An Investigation of Water Diffusivity Changes along the Perivascular Space in Elderly Subjects with Hypertension. <i>American Journal of Neuroradiology</i> , 2022, 43, 48-55.	1.2	28
2	Engagement in different sport disciplines during university years and risk of locomotive syndrome in older age: J-Fit+Study. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 36.	1.4	3
3	Short-Term SGLT2 Inhibitor Administration Does Not Alter Systemic Insulin Clearance in Type 2 Diabetes. <i>Biomedicines</i> , 2021, 9, 1154.	1.4	2
4	Association of physical fitness and motor ability at young age with locomotive syndrome risk in middle-aged and older men: J-Fit+ Study. <i>BMC Geriatrics</i> , 2021, 21, 89.	1.1	6
5	Associations of Exercise Habits in Adolescence and Old Age with Risk of Osteoporosis in Older Adults: The Bunkyo Health Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 5968.	1.0	6
6	Associations of Voluntary Exercise and Screen Time during the First Wave of COVID-19 Restrictions in Japan with Subsequent Grip Strength among University Students: J-Fit+ Study. <i>Sustainability</i> , 2021, 13, 13648.	1.6	2
7	Secular trends in the grip strength and body mass index of sport university students between 1973 and 2016: J-Fit+Study. <i>Journal of Exercise Science and Fitness</i> , 2020, 18, 21-30.	0.8	11
8	A body mass index over 22 kg/m ² at college age is a risk factor for future diabetes in Japanese men. <i>PLoS ONE</i> , 2019, 14, e0211067.	1.1	14
9	Slightly increased BMI at young age is a risk factor for future hypertension in Japanese men. <i>PLoS ONE</i> , 2018, 13, e0191170.	1.1	12
10	Muscle strength at young age is not associated with future development of type 2 diabetes in Japanese male athletes. <i>The Journal of Physical Fitness and Sports Medicine</i> , 2017, 6, 167-173.	0.2	4
11	Relationship between dietary patterns and risk factors for cardiovascular disease in patients with type 2 diabetes mellitus: a cross-sectional study. <i>Nutrition Journal</i> , 2015, 15, 15.	1.5	20
12	Poor sleep quality is associated with increased arterial stiffness in Japanese patients with type 2 diabetes mellitus. <i>BMC Endocrine Disorders</i> , 2015, 15, 29.	0.9	39
13	Cardiorespiratory fitness and the incidence of type 2 diabetes: a cohort study of Japanese male athletes. <i>BMC Public Health</i> , 2014, 14, 493.	1.2	18