Qiu-Yue Zhong

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7978880/publications.pdf

Version: 2024-02-01

713013 623188 21 683 14 21 citations g-index h-index papers 21 21 21 1100 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Suicide risk assessment: examining transitions in suicidal behaviors among pregnant women in Per $\tilde{\mathbb{A}}^{\circ}$. Archives of Women's Mental Health, 2019, 22, 65-73.	1.2	15
2	Associations of antepartum suicidal behaviour with adverse infant and obstetric outcomes. Paediatric and Perinatal Epidemiology, 2019, 33, 137-144.	0.8	3
3	Use of natural language processing in electronic medical records to identify pregnant women with suicidal behavior: towards a solution to the complex classification problem. European Journal of Epidemiology, 2019, 34, 153-162.	2.5	37
4	Causal Model of the Association of Social Support With Antepartum Depression: A Marginal Structural Modeling Approach. American Journal of Epidemiology, 2018, 187, 1871-1879.	1.6	18
5	The psychometric properties of the 16-item version of the Prodromal Questionnaire (PQ-16) as a screening instrument for perinatal psychosis. Archives of Women's Mental Health, 2018, 21, 563-572.	1.2	13
6	Childhood abuse and adult-onset asthma among Peruvian women. Journal of Asthma, 2018, 55, 430-436.	0.9	13
7	Trauma exposure and post-traumatic stress disorder in a cohort of pregnant Peruvian women. Archives of Women's Mental Health, 2018, 21, 193-202.	1.2	11
8	Association Between Migraine and Suicidal Behaviors: A Nationwide Study in the USA. Headache, 2018, 58, 371-380.	1.8	13
9	Adverse obstetric and neonatal outcomes complicated by psychosis among pregnant women in the United States. BMC Pregnancy and Childbirth, 2018, 18, 120.	0.9	43
10	Screening pregnant women for suicidal behavior in electronic medical records: diagnostic codes vs. clinical notes processed by natural language processing. BMC Medical Informatics and Decision Making, 2018, 18, 30.	1.5	46
11	Adverse obstetric outcomes during delivery hospitalizations complicated by suicidal behavior among US pregnant women. PLoS ONE, 2018, 13, e0192943.	1.1	23
12	Childhood physical and sexual abuse experiences associated with post-traumatic stress disorder among pregnant women. Annals of Epidemiology, 2017, 27, 716-723.e1.	0.9	23
13	Correlates of early pregnancy serum brain-derived neurotrophic factor in a Peruvian population. Archives of Women's Mental Health, 2017, 20, 777-785.	1.2	6
14	Psychometric Evaluation of the Ford Insomnia Response to Stress Test (FIRST) in Early Pregnancy. Journal of Clinical Sleep Medicine, 2016, 12, 579-587.	1.4	10
15	Childhood abuse and suicidal ideation in a cohort of pregnant Peruvian women. American Journal of Obstetrics and Gynecology, 2016, 215, 501.e1-501.e8.	0.7	28
16	Suicidal behavior-related hospitalizations among pregnant women in the USA, 2006–2012. Archives of Women's Mental Health, 2016, 19, 463-472.	1,2	21
17	Diagnostic Validity of the Generalized Anxiety Disorder - 7 (GAD-7) among Pregnant Women. PLoS ONE, 2015, 10, e0125096.	1.1	173
18	Association of poor subjective sleep quality with suicidal ideation among pregnant Peruvian women. General Hospital Psychiatry, 2015, 37, 441-447.	1.2	52

#	Article	IF	CITATIONS
19	Childhood abuse is associated with stress-related sleep disturbance and poor sleep quality in pregnancy. Sleep Medicine, 2015, 16, 1274-1280.	0.8	30
20	Using the Patient Health Questionnaire (PHQ-9) and the Edinburgh Postnatal Depression Scale (EPDS) to assess suicidal ideation among pregnant women in Lima, Peru. Archives of Women's Mental Health, 2015, 18, 783-792.	1.2	66
21	Association of decreased serum brain-derived neurotrophic factor (BDNF) concentrations in early pregnancy with antepartum depression. BMC Psychiatry, 2015, 15, 43.	1.1	39