

Xiao-Yu Qu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7987892/publications.pdf>

Version: 2024-02-01

14
papers

318
citations

840119

11
h-index

1058022

14
g-index

14
all docs

14
docs citations

14
times ranked

412
citing authors

#	ARTICLE	IF	CITATIONS
1	Calycosin ameliorates doxorubicin-induced cardiotoxicity by suppressing oxidative stress and inflammation via the sirtuin 1 "NOD-like receptor protein 3 pathway. <i>Phytotherapy Research</i> , 2020, 34, 649-659.	2.8	59
2	Dysregulation of BSEP and MRP2 May Play an Important Role in Isoniazid-Induced Liver Injury & via the SIRT1/FXR Pathway in Rats and HepG2 Cells. <i>Biological and Pharmaceutical Bulletin</i> , 2018, 41, 1211-1218.	0.6	47
3	XingNaoJing injection ameliorates cerebral ischaemia/reperfusion injury via SIRT1-mediated inflammatory response inhibition. <i>Pharmaceutical Biology</i> , 2020, 58, 16-24.	1.3	31
4	Autophagy inhibition enhanced assembly of the NLRP3 inflammasome is associated with cisplatin-induced acute injury to the liver and kidneys in rats. <i>Journal of Biochemical and Molecular Toxicology</i> , 2019, 33, e22208.	1.4	26
5	Xingnaojing Injection Protects against Cerebral Ischemia Reperfusion Injury via PI3K/Akt-Mediated eNOS Phosphorylation. <i>Evidence-based Complementary and Alternative Medicine</i> , 2018, 2018, 1-13.	0.5	25
6	Quercetin protected against isoniazide-induced HepG2 cell apoptosis by activating the SIRT1/ERK pathway. <i>Journal of Biochemical and Molecular Toxicology</i> , 2019, 33, e22369.	1.4	25
7	Quercetin attenuates NLRP3 inflammasome activation and apoptosis to protect INH-induced liver injury via regulating SIRT1 pathway. <i>International Immunopharmacology</i> , 2020, 85, 106634.	1.7	25
8	The role of Ntcp, Oatp2, Bsep and Mrp2 in liver injury induced by <i>Dioscorea bulbifera</i> L. and Diosbulbin B in mice. <i>Environmental Toxicology and Pharmacology</i> , 2017, 51, 16-22.	2.0	16
9	Ginsenoside Rg3 attenuates cisplatin-induced kidney injury through inhibition of apoptosis and autophagy inhibited NLRP3. <i>Journal of Biochemical and Molecular Toxicology</i> , 2021, 35, e22896.	1.4	13
10	<i>Dioscorea bulbifera</i> L. delays the excretion of doxorubicin and aggravates doxorubicin-induced cardiotoxicity and nephrotoxicity by inhibiting the expression of P-glycoprotein in mice liver and kidney. <i>Xenobiotica</i> , 2019, 49, 1116-1125.	0.5	12
11	Salvianolic Acid B Attenuates Apoptosis of HUVEC Cells Treated with High Glucose or High Fat via Sirt1 Activation. <i>Evidence-based Complementary and Alternative Medicine</i> , 2019, 2019, 1-11.	0.5	12
12	A meta-analysis of naldemedine for the treatment of opioid-induced constipation. <i>Expert Review of Clinical Pharmacology</i> , 2019, 12, 121-128.	1.3	12
13	Huaiqihuang (HQH) granule alleviates cyclophosphamide-induced nephrotoxicity via suppressing the MAPK/NF- κ B pathway and NLRP3 inflammasome activation. <i>Pharmaceutical Biology</i> , 2021, 59, 1423-1429.	1.3	9
14	Salvianolic acid inhibits the effects of high glucose on vascular endothelial dysfunction by modulating the Sirt1-eNOS pathway. <i>Journal of Biochemical and Molecular Toxicology</i> , 2019, 33, e22245.	1.4	6