

Chien-Hao Huang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7989636/publications.pdf>

Version: 2024-02-01

23
papers

236
citations

1307594

7
h-index

996975

15
g-index

23
all docs

23
docs citations

23
times ranked

378
citing authors

#	ARTICLE	IF	CITATIONS
1	Blockade of TNF- α signaling benefits cancer therapy by suppressing effector regulatory T cell expansion. <i>Oncimmunology</i> , 2015, 4, e1040215.	4.6	40
2	Recurrence of spontaneous bacterial peritonitis in cirrhotic patients non-prophylactically treated with norfloxacin: serum albumin as an easy but reliable predictive factor. <i>Liver International</i> , 2011, 31, 184-191.	3.9	28
3	Increased EMR2 expression on neutrophils correlates with disease severity and predicts overall mortality in cirrhotic patients. <i>Scientific Reports</i> , 2016, 6, 38250.	3.3	25
4	Differences in Liver Fibrosis Between Patients With Chronic Hepatitis B and C. <i>Journal of Ultrasound in Medicine</i> , 2015, 34, 813-821.	1.7	23
5	Memory Regulatory T cells Increase Only In Inflammatory Phase of Chronic Hepatitis B Infection and Related to Galectin-9/Tim-3 interaction. <i>Scientific Reports</i> , 2017, 7, 15280.	3.3	16
6	Increased regulatory T cells in patients with liver cirrhosis correlated with hyperbilirubinemia and predict bacterial complications. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2015, 30, 775-783.	2.8	13
7	Innate-like bystander-activated CD38+HLA-DR+CD8+T cells play a pathogenic role in patients with chronic hepatitis C. <i>Hepatology</i> , 2022, 76, 803-818.	7.3	11
8	Risk of cardiac arrhythmias in patients with chronic hepatitis B and C infections – A 13-year nationwide population-based study. <i>Journal of Cardiology</i> , 2019, 74, 333-338.	1.9	10
9	In HCV-related liver cirrhosis, local pulse wave velocity increases and in decompensated patients correlates with poorer survival. <i>PLoS ONE</i> , 2019, 14, e0212770.	2.5	10
10	Add-on sorafenib is beneficial for hepatocellular carcinoma patients with transarterial chemoembolization refractoriness: a real-world experience. <i>European Journal of Gastroenterology and Hepatology</i> , 2020, 32, 1192-1199.	1.6	9
11	Therapeutic Role of Inducible Nitric Oxide Synthase Expressing Myeloid-Derived Suppressor Cells in Acetaminophen-Induced Murine Liver Failure. <i>Frontiers in Immunology</i> , 2020, 11, 574839.	4.8	9
12	The incidence and predictors of post transarterial chemoembolization variceal bleeding in hepatocellular carcinoma patients. <i>Journal of the Formosan Medical Association</i> , 2020, 119, 635-643.	1.7	7
13	Interferon Is Superior to Direct Acting Antiviral Therapy in Tertiary Prevention of Early Recurrence of Hepatocellular Carcinoma. <i>Cancers</i> , 2020, 12, 23.	3.7	7
14	Fibrosis index predicts variceal bleeding and reduces need for nonselective beta-blocker in compensated cirrhosis with initial small esophageal varices without red-color sign. <i>Annals of Translational Medicine</i> , 2020, 8, 1223-1223.	1.7	6
15	Patients younger than forty years old with hepatitis C virus genotype 1 chronic infection had treatment responses similar to genotype 2 infection and not related to interleukin 28B polymorphism. <i>Annals of Hepatology</i> , 2013, 12, 62-69.	1.5	5
16	Timely eradication of HCV viremia by PegIFN/RBV is crucial in prevention of post RFA recurrence in CHC-HCC patients. <i>Journal of the Formosan Medical Association</i> , 2019, 118, 1239-1246.	1.7	4
17	Dynamic Change of Albumin-Bilirubin Score Is Good Predictive Parameter for Prognosis in Chronic Hepatitis C-hepatocellular Carcinoma Patients Receiving Transarterial Chemoembolization. <i>Diagnostics</i> , 2022, 12, 665.	2.6	4
18	Hepatic Encephalopathy and Spontaneous Bacterial Peritonitis Improve Cirrhosis Outcome Prediction: A Modified Seven-Stage Model as a Clinical Alternative to MELD. <i>Journal of Personalized Medicine</i> , 2020, 10, 186.	2.5	3

#	ARTICLE	IF	CITATIONS
19	Postembolization fever after transarterial chemoembolization is a sign of unfavorable therapeutic response in hepatocellular carcinoma patients. <i>Advances in Digestive Medicine</i> , 2020, 7, 83-92.	0.2	2
20	Complete response at first transarterial chemoembolization predicts favorable outcome in hepatocellular carcinoma. <i>American Journal of Cancer Research</i> , 2021, 11, 4956-4965.	1.4	2
21	Poor dietary intake improved by total excision of oral cavity metastases in a patient with hepatocellular carcinoma and elevated myeloid-derived suppressor cells. <i>Hepatobiliary Surgery and Nutrition</i> , 2020, 9, 558-561.	1.5	1
22	MELD score is the better predictor for 30-day mortality in patients with ruptured hepatocellular carcinoma treated by trans-arterial embolization. <i>American Journal of Cancer Research</i> , 2021, 11, 3726-3734.	1.4	1
23	<i>IL28B</i> polymorphism and early anemia predict the rapid null response in genotype 1 chronic hepatitis C with dual therapy. <i>Advances in Digestive Medicine</i> , 2015, 2, 128-134.	0.2	0